Temporal association between hospital admissions for asthma in Birmingham and ambient levels of sulphur dioxide and smoke

Sarah Walters, R K Griffiths, J G Ayres

Abstract

Background – A study was performed to determine whether daily and weekly variations in the levels of smoke and sulphur dioxide (SO₂) in Birmingham are related to hospital admissions for asthma and acute respiratory diseases.

Methods – Daily numbers of hospital admissions for asthma (ICD code 493) and acute respiratory conditions (ICD 466, 480–486, 490–496) for residents of Birmingham between 1988 and 1990 were obtained from West Midlands RHA Körner inpatient data. Average daily levels of sulphur dioxide and smoke were obtained from Birmingham City Council for the same period, together with daily meteorological summaries from the Department of Geography, University of Birmingham. With the exception of one day, all air pollution measurements remained within current EC guide levels. Data were divided into seasons and the relation between hospital admissions and pollutant levels were explored by stepwise least squares regression models. Meteorological variables (temperature, pressure, humidity) were entered into the model if they showed significant association with hospital admissions during the season in question. Analysis was undertaken for daily (same day and lagged by two days) and weekly pollutant levels. Admissions were lagged behind pollution levels to allow for delayed effects of pollutants.

Results – The mean daily level of smoke was 12.7 µg/m³ and of SO₂ was 39.1 µg/m³, with maxima of 188.3 µg/m³ and 126.3 µg/m³, respectively. Significant associations were found between hospital admissions for respiratory disease lagged by two days, and smoke and SO₂ levels during winter. Associations between admissions for asthma and smoke and SO₂ levels were significant at the 5% level. These were independent of temperature, pressure, and humidity. Stepwise regression including both pollutants showed that smoke, but not SO₂, was a significant independent predictor of hospital admissions for both asthma and all respiratory conditions. During winter a rise of 100 µg/m³ smoke might result in five (95% CI 0.6 to 9) more asthma admissions and 21.5 (95% CI 10 to 33) more acute respiratory admissions each day in Birmingham. A 100 µg/m³ rise in SO₂ might result in four

(0 to 7) more asthma admissions and 15.5 (6 to 25) more respiratory admissions each day. Independent associations were also found between weekly mean smoke and SO₂ levels and all respiratory admissions during autumn and winter. During summer, daily mean smoke and SO₂ levels were significantly associated with non-lagged daily admissions for all respiratory diseases ($p < 0.02$). There was no association between air pollution and hospital admissions during spring.

Conclusions – Daily variations in smoke and SO₂ levels are significantly associated with hospital admissions for asthma and respiratory disease during winter in Birmingham at levels of air pollutants within the EC guide levels. This association was independent of potential confounding effects of weather (temperature, pressure, humidity) and suggests that current levels of air pollution can still produce significant health effects.

(Thorax 1994;49:133–140)

The effects of air pollution on public health have been studied for many years. Studies of both short term health effects of temporal changes in air pollution levels, and long term effects on prevalence of disease and mortality, have been undertaken. Study methods used have included epidemiological studies – both ecological and planned observations – and controlled exposure experiments. Epidemiological studies have addressed a wide range of potential health outcomes including mortality, hospital attendances and admissions, and changes in symptoms and respiratory function. Despite all this work the exact nature and scale of the public health effects of air pollution, and the levels of pollutants at which health effects occur, remain unclear.

In 1952 a severe smog in London with exceptionally high levels of sulphur dioxide (SO₂) (4000 µg/m³) and smoke (estimated 7000–8000 µg/m³) resulted in more than 4000 deaths from respiratory and cardiovascular disease.¹ Following this episode, clean air legislation successfully reduced SO₂ and smoke levels throughout the UK.² With falling levels of these pollutants, interest and research into their effects on health waned in the UK. Studies from the USA suggest that current lower levels of particulates are still associated with mortality. A consistent independent relation has been shown between both total suspended particulates^{3,4} and particulates below

Institute of Public and Environmental Health, University of Birmingham Medical School, Birmingham B15 2TT
S Walters
R K Griffiths

Chest Research Institute, Birmingham Heartlands Hospital, Bordesley Green East, Birmingham B9 5ST
J G Ayres

Reprint requests to:
Dr S Walters.

Received 25 February 1993
Returned to authors
20 April 1993
Revised version received
9 August 1993
Accepted for publication
4 November 1993

10 μm in diameter (PM10)^{5,6} and daily mortality; the relation is strongest in patients with respiratory and cardiovascular conditions.

The effect of SO₂ and particulates on hospital admissions or attendances has also been studied. A series of Canadian studies showed significant relations between hospital admissions for asthma and respiratory diseases and air pollution. Admissions were significantly associated with levels of SO₂, coefficient of haze, and a composite air pollution index in southern Ontario.⁷ Bates and Sizto showed a relation between hospital admissions for asthma and respiratory diseases and levels of SO₂, ozone, oxides of nitrogen and sulphate during summer.⁸ In Vancouver, SO₂ was significantly associated with attendances at the emergency department for asthma and respiratory diseases during both summer and winter.⁹ This study also showed a significant association during summer between hospital attendances and sulphate and ozone levels. A recent study from Barcelona showed significant, though weak, relations between SO₂, smoke and carbon monoxide levels and hospital attendances in patients with chronic obstructive pulmonary disease (COPD).¹⁰

An important series of recent studies from Utah on the effects of PM10 pollution on various health outcomes has confirmed health effects of relatively low levels of particulates. Hospital admissions rose by 89% in children and by 47% in adults when PM10 levels exceeded 50 $\mu\text{g}/\text{m}^3$.¹¹ In a later study hospitalisation rates for Utah Valley were compared with those in two other valleys during a period of steel mill closure. Hospital admission rates were high in the polluted Utah Valley, but fell almost to those in the comparison valleys during closure of the mill, and rose again rapidly after reopening.¹² A study from Seattle confirmed the association between hospital attendances for asthma and PM10 levels.¹³ Further studies from Utah have shown PM10 to be associated with school absences, even when corrected for weather.¹⁴ Peak expiratory flow (PEF) in panels of children is significantly associated with PM10, even when days above the air quality standard of 150 $\mu\text{g}/\text{m}^3$ were excluded.¹⁵ The association was stronger for children with respiratory symptoms.

Early panel studies showed a relation between daily smoke and SO₂ levels in London and symptoms in patients with COPD¹⁶; however, pollution levels were considerably higher than at present. More recent studies in normal children in Holland showed an average 5% decrement in forced expiratory volume in one second (FEV₁) during two pollution episodes (SO₂ and total suspended particulate levels 200–300 $\mu\text{g}/\text{m}^3$).^{17,18} A large panel study in Holland in normal and asthmatic children showed an association between PEF, wheeze, bronchodilator use, and levels of PM10, smoke, and SO₂.¹⁹ A US study reported similar findings.²⁰ Levels of total suspended particulates and nitrogen dioxide (NO₂) were associated with daily reported cases of croup in Germany – a 27% rise in cases for a 60 $\mu\text{g}/\text{m}^3$ rise in particulates.²¹

Other pollutants, alone or in combination with SO₂ and particulates, have also been associated with health effects. In a panel study in Denver hydrogen ion concentration was significantly associated with several symptomatic indices of asthmatic status, as were fine particulates and sulphate.²² In Los Angeles, risk of asthma attacks was increased on days with high oxidant and particulate pollution.²³ Several studies in summer camps have confirmed that ozone levels can affect pulmonary function.^{24,25}

There is good experimental evidence to back up these epidemiological findings. SO₂ can cause bronchoconstriction at high levels in normal subjects,²⁶ but asthmatic subjects appear to react at one tenth of the concentration.²⁷ The effect is increased by exercise^{28,29} and may occur after exposure of five minutes or less.³⁰ Ozone^{31,32} and NO₂³³ have also been shown to cause bronchoconstriction, either directly or via their effects on bronchial sensitivity.³⁴

As well as studies of acute effects, there is some evidence that exposure to air pollution can produce longer term effects on symptom prevalence. The Harvard Six Cities study reported an association between prevalence of cough, bronchitis, and chest illness in children and ambient particulate and SO₂ levels.³⁵ This was subsequently confirmed in a second study which showed symptom prevalence to be more strongly associated with particulate levels than SO₂ or NO₂.³⁶ Associations were stronger in children with asthma, and for fine particulates (PM15).

Thus there is a substantial amount of published literature to show that particulate pollution and SO₂ is associated with acute health effects. However, there are no recent studies from the UK of health effects at current levels of particulate and SO₂ pollution. Hospital admissions for asthma are rising in the West Midlands, particularly in children.³⁷ This study sought to determine whether SO₂ and smoke (the only available measure of particulates) were associated with hospital admissions for asthma and respiratory disease in Birmingham, to determine the size and potential public health importance of any association seen, and to establish whether these health effects could be detected using routinely collected data.

Methods

DATA SOURCES

Data on hospital admissions for residents of Birmingham admitted to hospital within the West Midlands Region were obtained in the form of a Körner minimum inpatient data set from the information department at West Midlands Regional Health Authority. This included all patients admitted with a diagnosis of asthma or acute respiratory disease (ICD-9 codes 466, 480–486 and 490–496) for the period April 1988 to March 1990.

Air pollution data were obtained from the Environmental Protection Unit at Birmingham City Council for the same time period.

Particulate pollution was measured by the British standard black smoke method.³⁸ This method has the disadvantage that it measures only coloured particulates, which may constitute a variable proportion of total particulates, and in urban areas this may be a minority.³⁹ A high proportion of the smoke in urban areas comes from diesel exhaust emissions.⁴⁰ Use of the black smoke method in cities is likely to underestimate total particulate load, and particularly acid aerosols which may also cause health effects. Unfortunately, black smoke was the only measurement method used at the time of the study. It is recognised that use of black smoke will have underestimated total particulate exposure. Sulphur dioxide was measured by the hydrogen peroxide absorption method.³⁸ Both pollutants were measured at seven sites in Birmingham, and data were provided as 24 hour mean levels ($\mu\text{g}/\text{m}^3$). Full meteorological summaries including information about temperature, barometric pressure, and relative humidity were provided by the University of Birmingham Department of Geography for the period April 1988 to March 1990.

ANALYSIS

Data on daily total hospital admissions, air pollution levels, and meteorological conditions were stored and analysed on a personal computer using the Epi-Info and SPSS for Windows statistical packages, and Quattro Pro for Windows.

The data were divided into seasons as follows: spring (March, April, May), summer (June, July, August), autumn (September, October, November) and winter (December, January, February), since both hospital admissions and air pollution mixtures differ markedly by season. During summer, smoke and SO_2 are mixed with ozone, acid aerosols, and other secondary photochemical pollutants for which contemporary data were not available. There are also other potential environmental cofactors such as airborne pollen and spores which show seasonal patterns. Asthma admis-

sions show seasonal peaks in early summer and autumn. Because each season shows different atmospheric chemistry characteristics and different potential covariates such as pollen and respiratory infection, the relations between pollution and hospital admissions were examined seasonally. This approach has been taken by other authors.^{8,9}

Analysis was performed by least squares regression. For each season the bivariate relation between hospital admissions and three meteorological variables was determined by simple least squares regression. Those meteorological variables showing a significant ($p < 0.05$) association with hospital admissions in each season were entered into a least squares regression model along with each pollutant in turn. Stepwise multiple regression models were also constructed using each pollutant in turn and both pollutants together. Regression coefficients (β coefficients), 95% confidence intervals, and p values were calculated for the relation between each pollutant and hospital admissions for asthma and all respiratory conditions. This was repeated with hospital admissions lagged two days behind pollution variables to allow for delayed effects, as previously described.⁹ During the winter only, analysis was repeated using successive daily lags for hospital admissions, and the strongest associations obtained when admissions were lagged two to four days behind pollution peaks (fig 1). Because of this, regression models were also developed using three day moving average levels of pollutant. For all models the degree of autocorrelation in the dependent variable was estimated by the Durbin-Watson statistic.

Similar regression models were developed for the association between weekly mean pollutant levels and weekly total hospital admissions for asthma and all respiratory diseases, without lagging of hospital admissions, to examine the effect of more prolonged exposure to high levels of pollutant over a period of several days.

In total, 60 multiple regression models were constructed, and seven additional bivariate regressions were carried out to assess the degree to which hospital admission lagged behind pollutant levels. For this reason the level of statistical significance used in this study was 1%.

Results

TEMPORAL PATTERNS IN AIR POLLUTION AND HOSPITAL ADMISSIONS

Temporal variations in weekly mean smoke and SO_2 levels are shown in fig 2, and variations in weekly total hospital admissions for asthma and all respiratory conditions are shown in fig 3. Levels of smoke showed consistent winter peaks, but SO_2 peaked at other times of year as well. All respiratory admissions showed consistent winter peaks with smaller peaks during spring. Asthma admissions showed peaks during early summer and autumn, but also at other times of year. The mean levels and 10th to 90th centiles of smoke and SO_2 are given in table 1, and similar

Figure 1 Correlation coefficient for relation between lagged hospital admissions, Birmingham, winter 1988-90.

figures for asthma and respiratory admissions are presented in table 2. The mean daily smoke level was $12.7 \mu\text{g}/\text{m}^3$ over the two year period, and the mean level of SO_2 was $39.1 \mu\text{g}/\text{m}^3$. The maxima were $188.3 \mu\text{g}/\text{m}^3$ and $126.3 \mu\text{g}/\text{m}^3$ respectively.

Figure 2 Mean weekly levels of smoke and sulphur dioxide, Birmingham, April 1988 to March 1990.

Figure 3 Weekly total admissions for asthma and respiratory disease, Birmingham residents, April 1988 to March 1990.

Table 1 Mean (SE) and centile points for levels of smoke and sulphur dioxide ($\mu\text{g}/\text{m}^3$) in Birmingham, 1988–90

	Mean (SE)	Centiles				
		10th	25th	50th	75th	90th
Smoke						
Spring	10.15 (0.48)	4	5	9	14	19
Summer	7.19 (0.29)	3	4	6	10	12.5
Autumn	20.53 (1.74)	4	7	13	24	45
Winter	12.56 (0.91)	4	5	8	15	29
All year	12.74 (0.54)	4	5	8	15	25
Sulphur dioxide						
Spring	42.87 (1.35)	27	31	37	49	64.5
Summer	37.77 (1.17)	23	27	34	45	55.5
Autumn	40.91 (1.24)	26	29	36	48	68
Winter	34.18 (1.03)	22	25.2	30	37.7	52
All year	39.06 (0.62)	24	28	34	46	62

RELATION BETWEEN HOSPITAL ADMISSIONS AND AIR POLLUTION

Daily pollution levels and hospital admission

Results of multiple least squares regression models for daily levels of air pollution and daily hospital admissions are shown by season, and for zero and two day lags in hospital admissions in table 3. During winter daily levels of smoke and SO_2 were significantly associated with hospital admissions for all respiratory conditions at the 1% level. This was independent of temperature, pressure and humidity, all of which were entered into the regression model. The association was seen for hospital admissions without and with a two day lag. Daily level of smoke was associated with hospital admissions for asthma, lagged by two days at the 3% level, and the association between SO_2 and asthma fell just below the 5% significance level. Stepwise regression models revealed smoke and SO_2 to be significant independent determinants of hospital admissions for both asthma and all respiratory conditions at or below the 5% level. Stepwise multiple regression including both pollutants revealed smoke, but not SO_2 to be a significant independent determinant of wintertime hospital admissions for asthma and all respiratory conditions, lagged by two days, x-coefficient 2.22 (95% CI 1.23 to 3.22), $p < 0.00001$. There was no significant association between hospital admissions and daily air pollution levels during spring or autumn. During summer daily levels of smoke were significantly related to daily unlagged admissions for all respiratory diseases at the 1% level, and SO_2 was significantly associated at the 2% level. The results of regression analysis using three day moving average levels of pollutants and daily hospital admissions were very similar to those using daily averages with hospital admissions lagged by two days (table 4).

A scatter plot of daily levels of smoke and SO_2 and hospital admissions for all respiratory conditions lagged by two days during winter is shown in fig 4. There is a broad scatter of points but no clear suggestion of a threshold for effect.

These results suggest that over the range of air pollution tested during winter a rise of $100 \mu\text{g}/\text{m}^3$ smoke might result in five (95% CI 0.6 to 9) more asthma admissions and 21.5 (95% CI 10 to 33) more acute respiratory admissions a day in Birmingham. A $100 \mu\text{g}/\text{m}^3$ rise in SO_2 might result in four (0 to 7) more asthma admissions and 15.5 (6 to 25) more respiratory admissions a day.

Weekly pollution levels and hospital admissions

Results of multiple least squares regression using weekly mean pollution levels and weekly total hospital admissions are shown in table 5. All regression models included meteorological variables which were significantly associated with hospital admissions during the season in question.

Significant associations ($p < 0.01$) were shown between all respiratory admissions and smoke and SO_2 levels during winter, and

Table 2 Mean (SE) and centile points for daily hospital admissions for respiratory disease and asthma in Birmingham, 1988–90

	Mean (SE)	Centiles				
		10th	25th	50th	75th	90th
All respiratory diseases						
Spring	11.52 (0.40)	4	8	11	15	19.5
Summer	12.32 (0.38)	6.5	8	12	16	20
Autumn	17.42 (0.48)	10	13	17	21	27
Winter	17.48 (0.64)	9	12	16	21	30
All year	14.68 (0.26)	7	10	14	18	23
Asthma						
Spring	1.38 (0.18)	1	3	4	6	8
Summer	4.88 (0.20)	2	3	4	6	9
Autumn	7.34 (0.25)	3	5	7	10	12
Winter	5.27 (0.23)	2	3	5	7	9
All year	5.47 (0.12)	2	3	5	7	10

between all respiratory admissions and SO₂ levels only during autumn. All respiratory admissions were associated with weekly smoke levels in autumn but only at the 3% level. Asthma admissions were not significantly associated with weekly levels of smoke or SO₂ during any season. Stepwise multiple regres-

sion models including both pollutants showed that, during winter, weekly SO₂, but not smoke, levels were significantly associated with weekly hospital admissions for asthma ($p < 0.005$). All respiratory admissions were significantly associated with weekly smoke levels ($p < 0.001$), but not SO₂. During autumn SO₂ was significantly associated with weekly admissions for all respiratory diseases ($p < 0.001$). No significant associations between weekly total hospital admissions and weekly pollutant levels were found during spring or summer. During winter a rise of 25 µg/m³ in weekly mean smoke levels might result in 83 (95% CI 44 to 122) more respiratory admissions during the same week, and 43 (15 to 72) more asthma admissions.

Association between meteorological conditions and air pollution levels

The association between meteorological conditions (temperature, barometric pressure, and

Table 3 Multiple regression analysis: daily mean levels of sulphur dioxide and smoke (µg/m³) and daily total hospital admissions by season in Birmingham, 1988–90

	Smoke v asthma	Smoke v all respiratory diseases	SO ₂ v asthma	SO ₂ v all respiratory diseases
No lag in hospital admissions				
Spring				
x-coefficient (95% CI)	0.006 (−0.005 to 0.06)	0.05 (−0.07 to 0.17)	−0.001 (0.19 to −0.0008)	−0.008 (−0.005 to 0.04)
p value	NS	NS	NS	NS
Variables significant in stepwise model	None	Pressure	None	Pressure
Summer				
x-coefficient (95% CI)	0.057 (−0.04 to 0.16)	0.29 (0.10 to 0.48)	0.014 (−0.10 to 0.39)	0.059 (0.011 to 0.106)
p value	NS	<0.005	NS	<0.02
Variables significant in stepwise model	None	Smoke	None	SO ₂
Autumn				
x-coefficient (95% CI)	−0.018 (−0.04 to 0.01)	−0.001 (−0.04 to 0.4)	−0.019 (−0.05 to 0.13)	0.04 (−0.18 to 0.102)
p value	NS	NS	NS	NS
Variables significant in stepwise model	None	Temperature	None	Temperature
Winter				
x-coefficient (95% CI)	0.028 (−0.014 to 0.07)	0.20 (0.094 to 0.313)	0.027 (−0.008 to 0.061)	0.18 (0.088 to 0.268)
p value	NS	<0.0005	NS	<0.0002
Variables significant in stepwise model	Smoke	Smoke, temperature	SO ₂ , pressure	SO ₂ , temperature, humidity
Hospital admissions lagged by two days				
Spring				
x-coefficient (95% CI)	0.009 (−0.05 to 0.07)	−0.005 (−0.13 to 0.12)	−0.006 (−0.03 to 0.14)	−0.03 (−0.08 to 0.10)
p value	NS	NS	NS	NS
Variables significant in stepwise model	None	Pressure	None	Pressure
Summer				
x-coefficient (95% CI)	0.077 (−0.022 to 0.18)	0.15 (−0.04 to 0.35)	0.012 (−0.013 to 0.37)	0.029 (−0.018 to 0.08)
p value	<0.2	<0.2	NS	NS
Variables significant in stepwise model	None	None	None	None
Autumn				
x-coefficient (95% CI)	−0.005 (−0.03 to 0.02)	0.007 (−0.35 to 0.49)	−0.011 (−0.04 to 0.22)	0.04 (−0.02 to 0.10)
p value	NS	NS	NS	<0.2
Variables significant in stepwise model	None	Temperature	None	Temperature
Winter				
x-coefficient (95% CI)	0.049 (0.006 to 0.093)	0.215 (0.102 to 0.327)	0.036 (0.000 to 0.073)	0.155 (0.059 to 0.251)
p value	<0.05	0.0002	<0.1	<0.002
Variables significant in stepwise model	Smoke	Smoke, humidity	SO ₂	SO ₂ , temperature, humidity

Table 4 Multiple regression analysis: three day moving average levels of smoke and sulphur dioxide v daily total hospital admissions in Birmingham, 1988–90

Three day moving average pollution levels	Smoke v asthma	Smoke v all respiratory diseases	SO ₂ v asthma	SO ₂ v all respiratory diseases
Spring				
x-coefficient (95% CI)	−0.007 (−0.09 to 0.07)	−0.02 (−0.19 to 0.16)	−0.007 (−0.03 to 0.16)	−0.03 (−0.08 to 0.02)
p value	NS	NS	NS	NS
Variables significant in stepwise model	None	Pressure	None	Pressure
Summer				
x-coefficient (95% CI)	0.021 (−0.11 to 0.15)	0.25 (−0.008 to 0.50)	0.016 (−0.001 to 0.04)	0.069 (0.01 to 0.12)
p value	NS	0.05	NS	<0.02
Variables significant in stepwise model	Humidity	None	Humidity	SO ₂
Autumn				
x-coefficient (95% CI)	−0.03 (−0.06 to −0.002)	0.013 (−0.4 to 0.7)	−0.26 (−0.7 to 0.13)	0.03 (−0.04 to 0.11)
p value	<0.05	NS	NS	NS
Variables significant in stepwise model	None	Temperature	None	Temperature
Winter				
x-coefficient (95% CI)	0.03 (−0.02 to 0.078)	0.29 (0.17 to 0.42)	0.039 (0 to 0.78)	0.23 (0.14 to 0.33)
p value	NS	<0.0001	<0.05	<0.0001
Variables significant in stepwise model	Pressure, temperature	Smoke, temperature	SO ₂ , pressure	SO ₂ , temperature

Figure 4 Scatter plot between daily mean smoke level and daily hospital admissions for respiratory disease, Birmingham, winter 1988–9 and 1989–90.

relative humidity) and levels of smoke and SO_2 are shown by season in table 6.

Discussion

This is the first recent study of the acute health effects of air pollutants in the UK on hospital admissions for respiratory diseases. We have shown that there is a significant association between hospital admissions for asthma and all respiratory conditions and air pollution levels during the winter at levels of air pollution which remained within EC guide levels, with

the exception of one day. Associations at other times of year were present but weaker. These associations were independent of significant confounding meteorological variables (temperature, pressure, relative humidity). Smoke, and not SO_2 , was the important independent predictor of daily hospital admissions during winter, and of weekly admissions for all respiratory conditions. This confirms the potential importance of health effects of particulates from the studies of Pope *et al* in Utah where PM_{10} was found to be a significant predictor of a wide range of respiratory morbidity at very low concurrent levels of SO_2 .^{11 12 14 15} SO_2 was a significant predictor of weekly asthma admissions.

These associations were weak, but statistically significant, and consistent in strength with those found in other studies.^{8–10} However, the North American studies^{8,9} found significant associations between hospital admissions and attendances and air pollution levels during summer, whereas this study has found significant associations mainly during winter. This may reflect climatic differences resulting in differences in seasonality of air pollution peaks. In the study of COPD admissions in Barcelona, seasonal breakdown revealed strongest associations between black smoke and emergency room admissions for COPD in winter,¹⁰ while associations between SO_2 levels and admissions were strongest in summer. Both pollutants were significantly associated with pollutants in all seasons except autumn. However, mean levels of both SO_2 and, particularly, black smoke were higher in Barcelona than in Birmingham ($56.5 \mu\text{g}/\text{m}^3$ and $72.9 \mu\text{g}/\text{m}^3$ respectively). The results of this

Table 5 Multiple regression analysis: weekly mean smoke and sulphur dioxide levels v weekly total hospital admissions in Birmingham, 1988–90

Weekly average pollution levels	Smoke v asthma	Smoke v all respiratory diseases	SO_2 v asthma	SO_2 v all respiratory diseases
Spring				
x-coefficient (95% CI)	0.103 (–1.24 to 1.44)	–0.13 (–3.7 to 3.44)	–0.096 (–0.39 to 0.19)	–0.39 (–1.15 to 0.37)
p value	NS	NS	NS	NS
Variables significant in stepwise model	None	None	None	None
Summer				
x-coefficient (95% CI)	1.14 (–0.88 to 3.17)	3.42 (0.08 to 6.77)	0.167 (–0.12 to 0.54)	0.59 (–0.023 to 1.22)
p value	NS	<0.05	NS	<0.1
Variables significant in stepwise model	None	None	None	None
Autumn				
x-coefficient (95% CI)	0.176 (–0.41 to 0.76)	1.07 (0.11 to 2.03)	0.46 (–0.05 to 0.96)	1.38 (0.596 to 2.16)
p value	NS	<0.05	<0.1	<0.002
Variables significant in stepwise model	None	Smoke	None	SO_2
Winter				
x-coefficient (95% CI)	0.5 (–0.25 to 1.24)	3.30 (1.75 to 4.87)	0.443 (0 to 0.89)	1.72 (0.58 to 2.87)
p value	NS	<0.0005	0.05	0.005
Variables significant in stepwise model	Smoke	Smoke, humidity	SO_2	SO_2 , humidity

Note: Stepwise regression models including meteorological variables of temperature, pressure and humidity, and both pollutants gave the following significant results:

Autumn: All respiratory SO_2 significant (x-coefficient 1.59, 0.91 to 2.27, $p < 0.001$).

Winter: All respiratory smoke significant (x-coefficient 3.37, 2.10 to 4.64, $p < 0.0001$). Asthma, SO_2 significant (x-coefficient 0.63, 0.25 to 1.01, $p < 0.005$).

Table 6 Association between levels of smoke and sulphur dioxide and meteorological variables (temperature, pressure, humidity) (x-coefficients and p values given)

	Spring	Summer	Autumn	Winter
Smoke				
Temperature	0.18 (NS)	0.27 ($p < 0.001$)	–2.28 ($p < 0.0001$)	–1.75 ($p < 0.0001$)
Pressure	0.87 ($p < 0.1$)	0.21 ($p < 0.0001$)	0.69 ($p < 0.0001$)	0.19 ($p < 0.001$)
Humidity	–0.02 (NS)	–0.05 ($p < 0.02$)	0.27 ($p < 0.05$)	0.229 ($p < 0.005$)
Sulphur dioxide				
Temperature	0.70 ($p < 0.05$)	0.31 ($p < 0.0001$)	–1.69 ($p < 0.0001$)	–1.53 ($p = 0.0001$)
Pressure	0.24 ($p < 0.1$)	0.71 ($p < 0.0001$)	0.69 ($p < 0.0001$)	0.195 ($p = 0.0007$)
Humidity	–0.14 (NS)	–0.19 ($p < 0.05$)	0.06 (NS)	0.193 ($p < 0.05$)

study are more consistent with those from Barcelona than the North American studies, and suggest that, in a European winter, particulate pollution below the EC guide level may be associated with significant health effects.

The associations between air pollution levels and hospital admissions observed in winter in this study may indicate that smoke and SO₂ are contributory causes of acute respiratory episodes, or that other confounding factors are operating. This analysis controlled for the effects of temperature, pressure, and humidity by including them in the regression model. However, it cannot control for other covariates which may have a confounding effect. This includes behavioural covariates. Cold weather is associated with higher levels of pollutants, but may also make people use more heating, and stay indoors more. The use of heating may result in higher exposure to indoor air pollutants from fuel including oxides of nitrogen which can influence the prevalence of respiratory symptoms.⁴¹ Staying indoors may also assist in the transmission of viral upper respiratory tract infections, and increase exposure of atopic subjects to indoor allergens such as house dust mite. All these factors, not directly associated with outdoor air pollution levels, may thus produce a confounding effect and could not be measured in this study. Air pollution may be acting as an indicator for other triggers as well as a direct trigger of respiratory episodes in itself.

The relative absence of associations between daily hospital admissions and air pollution during other seasons is interesting. Firstly, in this study we only measured SO₂ and smoke, but particularly during the summer these may covary with other pollutants present on hot sunny days with low winds and high barometric pressure, in particular ozone, oxides of nitrogen, and acid aerosols. They may also covary with grass pollen and other triggers for asthmatic attacks. The only significant association seen during summer was between all respiratory conditions and both smoke and SO₂ levels, but without any lagged effect. This suggests that the association may be due to covariation between these pollutants and other environmental triggers for respiratory disease. The British black smoke method of measuring particulates only measures coloured particulate matter, of which a large proportion comes from diesel vehicle exhausts.⁴⁰ However, the proportion of total particulates accounted for by the black smoke fraction may vary between seasons and with meteorological conditions. This means that black smoke may be a relatively poor index of total particulate exposure, and makes interpretation of negative results difficult. Finally, associations were noted during winter which is the season during which highest levels of smoke are seen. This may represent a threshold effect.

There are problems associated with the use of hospital admissions as an index of respiratory morbidity. Hospital admissions represent a complex social and behavioural process and may be influenced by factors not directly related to disease severity. If the behavioural

influences on hospital admission covary with pollution levels or weather conditions, this might influence the associations seen in this study. Factors such as pet ownership, smoking, heating and cooking methods, family history, and occupation might be expected to affect the prevalence of respiratory disease, but do not show rapid changes over time and are unlikely to act as confounding factors in this study.

This study was restricted by data availability to investigating the effects of SO₂ and smoke, both of which are now at low levels compared with 30 years ago and have not shown any tendency to rise over the previous decade.² It would be difficult to propose that these pollutants are the cause of rising levels of hospital admissions for asthma in the West Midlands Region. Recent concern about the health effects of air pollution has centred around pollutants associated with motor vehicles, including oxides of nitrogen, hydrocarbons, and diesel exhaust particulates.⁴² Although black smoke is quite a good measure of diesel exhaust particulates in an urban area, no measurements of oxides of nitrogen, hydrocarbons, or ozone were made at the time of this study. Evaluation of their health effects and covariance with smoke and SO₂ must await analysis of recently installed monitoring equipment as part of the extended urban air pollution monitoring network established by the Department of the Environment in Birmingham and other cities in the UK.

Although weather conditions are important determinants of air pollution levels in Birmingham, clean air legislation has proved that action to control emissions has been successful.² However, the cost of further reductions in emissions from motor vehicles and other important industrial sources must be balanced against the potential health improvements achieved. This study has shown that routinely collected data can be used, with certain limitations, to assess the potential health effects of air pollution and potential benefits of reducing levels. However, in view of potential confounding factors, further more specific population based and panel based studies are needed to establish the weight of effect of environmental factors, including air pollution, on respiratory health.

1 Ministry of Health. *Mortality and morbidity during the London fog of December 1952*. London: HMSO, 1954.

2 Quality of Urban Air Review Group. *Urban air quality in the United Kingdom*. London: Department of the Environment, 1993.

3 Schwartz J. Particulate air pollution and daily mortality in Detroit. *Environ Res* 1991;56:204-13.

4 Schwartz J, Dockery DW. Increased mortality in Philadelphia associated with daily air pollution concentrations. *Am Rev Respir Dis* 1992;145:600-4.

5 Schwartz J, Dockery DW. Particulate air pollution and daily mortality in Steubenville, Ohio. *Am J Epidemiol* 1992;135:12-9.

6 Pope CA, Schwartz J, Ransom MR. Daily mortality and PM10 pollution in Utah valley. *Arch Environ Health* 1992;47:211-7.

7 Levy D, Gent M, Newhouse MT. Relationship between acute respiratory illness and air pollution levels in an industrial city. *Am Rev Respir Dis* 1977;116:167-73.

8 Bates DV, Sizto R. Air pollution and hospital admissions in Southern Ontario: the acid summer haze effect. *Environ Res* 1987;43:317-31.

9 Bates DV, Baker-Anderson M, Sizto R. Asthma attack periodicity: a study of hospital emergency visits in Vancouver. *Environ Res* 1990;51:51-70.

- 10 Sunyer J, Anto JM, Murillo C, Saez M. Effects of urban air pollution on emergency room admissions for chronic obstructive pulmonary disease. *Am J Epidemiol* 1991;134:277-86.
- 11 Pope CA. Respiratory disease associated with community air pollution and a steel mill in Utah Valley. *Am J Public Health* 1989;79:623-8.
- 12 Pope CA. Respiratory hospital admissions associated with PM10 pollution in Utah, Salt Lake and Cache Valleys. *Arch Environ Health* 1991;46:90-7.
- 13 Schwartz J, Slater D, Larson TV, Pierson WE, Koenig WQ. Particulate air pollution and hospital emergency visits for asthma in Seattle. *Am Rev Respir Dis* 1992;145(4 part 2):A88.
- 14 Ransom MR, Pope CA. Elementary school absences and PM10 pollution in Utah Valley. *Environ Res* 1992;58:204-19.
- 15 Pope CA, Dockery DW. Acute health effects of PM10 pollution on symptomatic and asymptomatic children. *Am Rev Respir Dis* 1992;145:1123-8.
- 16 Waller RE. Air pollution and community health. *J R Coll Physicians Lond* 1971;5:362-8.
- 17 Dassen W, Brunekreef B, Hoek G, Hofschreuder P, Staasen B, de-Groote H, et al. Decline in children's lung function during an air pollution episode. *J Am Poll Control Assoc* 1986;36:1223-7.
- 18 Brunekreef B, Lumens M, Hoek G, Hofschreuder P, Fischer P, Biersteker K. Pulmonary function changes associated with an air pollution episode in January 1987. *J Am Poll Control Assoc* 1989;39:1444-7.
- 19 Hoek G, Brunekreef B, Roemer W. Acute effects of moderately elevated wintertime air pollution on respiratory health of children. *Am Rev Respir Dis* 1991;142(4 part 2):A88.
- 20 Neas LM, Dockery DW, Spengler JD, Speitzer FE, Tolend DJ. The association of ambient air pollution with twice daily peak respiratory flow measurements in children. *Am Rev Respir Dis* 1992;145(4 part 2):A429.
- 21 Schwartz J, Spix C, Wichmann HE, Malin E. Air pollution and acute respiratory illness in five German communities. *Environ Res* 1991;56:1-14.
- 22 Ostro BD, Lipsett MJ, Wiener MB, Selner JC. Asthmatic responses to airborne acid aerosols. *Am J Public Health* 1991;81:694-702.
- 23 Whittemore AS, Korn EL. Asthma and air pollution in the Los Angeles area. *Am J Public Health* 1980;70:687-97.
- 24 Spektor DM, Thurston GD, Mao J, He D, Hayes C, Lippman M. Effects of single and multiday ozone exposures on respiratory function in active normal children. *Environ Res* 1991;55:107-22.
- 25 Liou PJ, Vollmuth TA, Lipmann M. Persistence of peak flow decrement in children following ozone exposures exceeding the National Ambient Air Quality Standard. *J Am Poll Control Assoc* 1985;35:1068-71.
- 26 Lawther PJ, McFarlane AJ, Waller RE, Brooks AGF. Pulmonary function and sulfur dioxide - some preliminary findings. *Environ Res* 1975;10:355-67.
- 27 Sheppard D, Wong WS, Uehara CF, Nadel JA, Boushey HA. Lower threshold and greater bronchomotor responsiveness of asthmatic subjects to sulfur dioxide. *Am Rev Respir Dis* 1980;122:873-8.
- 28 Linn WS, Venet TG, Shamoo DA, Valencia LM, Anzar UT, Speir CE, et al. Respiratory effects of sulfur dioxide in heavily-exercising asthmatics: a dose-response study. *Am Rev Respir Dis* 1983;127:278-83.
- 29 Linn WS, Avol EL, Penc R-C, Shamoo SA, Hackney JD. Replicated dose-response study of sulfur dioxide effects in normal, atopic and asthmatic volunteers. *Am Rev Respir Dis* 1987;136:1127-34.
- 30 Balmes JR, Fine JM, Sheppard D. Symptomatic bronchoconstriction after short-term inhalation of sulfur dioxide. *Am Rev Respir Dis* 1987;136:1117-21.
- 31 Hazucha MJ. Relationship between ozone exposure and pulmonary function changes. *J Appl Physiol* 1987;62:1671-80.
- 32 Linn WS, Avol EL, Shamoo DA, Speir CE, Valencia LM, Venet TG, et al. Dose response study of healthy heavily-exercising men exposed to ozone at concentrations near the ambient air quality standard. *Toxicol Ind Health* 1986;2:99-112.
- 33 Orehek J, Massoni JP, Gattard C, Charpin J. Effect of short term low level nitrogen dioxide exposure on bronchial sensitivity of asthmatic patients. *J Clin Invest* 1986;56:301-7.
- 34 Molino NA, Wright SC, Katz I, Tarlo S, Silverman F, McClean PA, et al. Effect of low concentrations of ozone on inhaled allergen responses in asthmatic subjects. *Lancet* 1991;338:199-203.
- 35 Ware JH, Ferris BG Jr, Dockery DW, Spengler JD, Stram DO, Speizer FE. Effects of sulfur oxides and suspended particulates on respiratory health of preadolescent children. *Am Rev Respir Dis* 1986;133:834-42.
- 36 Dockery DW, Speizer FE, Stram DO, Ware JH, Spengler JD, Ferris BG Jr. Effects of inhalable particulates on respiratory health of children. *Am Rev Respir Dis* 1989;139:587-94.
- 37 West Midlands Regional Health Authority. *Partners in Health: Annual Report of the Regional Director of Public Health*. West Midlands RHA, 1992.
- 38 Harrison RM. Important air pollutants and their chemical analysis. In Harrison RM, ed. *Pollution: causes, effects and control*. Cambridge: Royal Society of Chemistry, 1990:127-55.
- 39 Clarke AG, Willison MJ, Ziki EM. A comparison of urban and rural aerosol composition using a dichotomous sampler. *Atmos Environ* 1984;23:1767-75.
- 40 Ball DJ, Caswell R. Smoke from diesel engined road vehicles - an investigation into the basis of British and European emission standards. *Atmos Environ* 1983;17:169-81.
- 41 Neas LM, Dockery DW, Ware JH, Spengler JD, Speizer FE, Ferris BG Jr. Association of indoor nitrogen dioxide with respiratory symptoms and pulmonary function in children. *Am J Epidemiol* 1991;134:204-19.
- 42 Godlee F. Air pollution: II. Road traffic and modern industry. *BMJ* 1991;303:1539-43.