Effects of *Aspergillus fumigatus* culture filtrate on antifungal activity of human phagocytes in vitro

Takako Murayama, Ryoichi Amitani, Yumiko Ikegami, Rieko Kawanami, Woon Joo Lee, Ryuhei Nawada

Abstract

Background—*Aspergillus fumigatus* can colonise the airways and the lungs with localised underlying conditions and occasionally invade the surrounding lung tissues even in subjects without systemic predisposing factors, presumably by escaping the local host defences. The aim of this study was to investigate the effects of *A. fumigatus* culture filtrate (ACF) on the activities of human phagocytes—**inhibition of germination of *A. fumigatus* spores by alveolar macrophages (AMs) and hyphal damage by polymorphonuclear leucocytes (PMNs)—which are the critical host defences against *A. fumigatus*.**

Methods—Spores were incubated with AMs at a ratio of 1:1 in a medium containing different concentrations of ACF for 10 hours at 37°C. Spore germination was visualised with light microscopy and the inhibition rate was calculated. The percentage of hyphal damage caused by PMNs pretreated with various concentrations of ACF was measured by a colorimetric tetrazolium metabolic assay.

Results—The inhibition rate of spore germination by AMs cultured with medium alone (control) was 90 (0.8)% whereas that by AMs cultured with the medium containing 10% ACF was significantly ($p < 0.05$) reduced to 41.7 (4.6)%. ACF suppressed the inhibition of spore germination in a dose dependent manner without altering the phagocytosing activity against the spores. The percentage of hyphal damage caused by PMNs pretreated with medium-199 (control) was 78.1 (2.3)% compared with 65.3 (2.8)% when PMNs were pretreated with 50% ACF ($p < 0.05$).

Conclusions—*A. fumigatus* releases biologically active substance(s) which suppress the inhibition of spore germination by AMs and also suppress PMN mediated hyphal damage, and thus may contribute to the pathogenicity of this fungus.

(Thorax 1998;53:975-978)

Keywords: *Aspergillus fumigatus*; human phagocytes; host defence

gatus frequently colonises damaged lungs and airways including healed tuberculosis, bronchiectasis and cystic fibrosis, and occasionally invades the surrounding lung tissues even in patients without apparently compromised systemic host defences.^{1,2} However, the pathogenic mechanisms and substances are still poorly understood.

It has been proposed that host defences against *A. fumigatus* consist of mucociliary clearance in the airways, alveolar macrophages (AMs), and polymorphonuclear leucocytes (PMNs). AMs can ingest the inhaled spores of *A. fumigatus*, inhibit them from germination, and kill them. The germinated spores and hyphae that escape the mucociliary clearance and AMs can be killed by PMNs, mainly by an oxidative mechanism.^{3,4} These lines of defence cooperate in elimination of the fungus.

We have previously reported that *A. fumigatus* culture filtrate (ACF) causes damage to respiratory ciliated epithelium by reducing ciliary beat frequency, and suppresses human phagocyte function including PMN migration towards formyl-methionyl-leucyl-phenylalanine and O_2^- release from PMNs in response to phorbol myristate acetate, and concluded that *A. fumigatus* produces several substances which suppress phagocyte function and mucociliary clearance.⁵⁻⁷

In the present study we have investigated the effects of ACF on inhibition of spore germination by human AMs and hyphal damage by human PMNs, in addition to the effects of ACF on ingestion of spores by AMs and the release of O_2^- from PMNs in response to *A. fumigatus* hyphae.

Methods

FUNGUS

A. fumigatus, YN strain, clinically isolated from the sputum of a patient with invasive pulmonary aspergillosis, was used in the present study. The strain was stored on potato dextrose agar (PDA) (Difco Laboratories, Detroit, Michigan, USA) slants at -80°C. Spores were obtained by scraping the surface of the slants, suspended in phosphate buffered saline (PBS), and kept at 4°C. To induce spore germination a suspension of 1.0×10^6 spores in RPMI-1640 (Gibco BRL, Paisley, UK) was incubated in polypropylene tubes (Falcon, Becton Dickinson, Lincoln, New Jersey, USA) for 16 hours at 30°C and for an additional two hours at 37°C.⁸ After incubation for 18 hours almost all spores developed into hyphae with a length of 10-30 µm, but not branched. The hyphae were washed with PBS, vigorously vortexed, and used in the PMN experiments.

Department of
Infection and
Inflammation, Chest
Disease Research
Institute, Kyoto
University, Sakyo-ku,
Kyoto 606-8397, Japan
T Murayama
R Amitani
Y Ikegami
R Kawanami
W J Lee
R Nawada

Correspondence to:
Dr T Murayama.

Received 15 September 1997
Returned to author
12 November 1997
Revised manuscript received
1 May 1998
Accepted for publication
6 May 1998

Aspergillus species, most commonly *Aspergillus fumigatus*, are responsible for a wide range of diseases including saprophytic colonisation, chronic necrotising aspergillosis, and rapidly invasive diseases.^{1,2} Although acute progressive and invasive diseases occur most commonly in systemic immunocompromised hosts, *A. fumi-*

PREPARATION OF *A. FUMIGATUS* CULTURE FILTRATE

A. fumigatus culture filtrate (ACF) was prepared as described previously.⁵⁻⁷ Spores (2×10^7) of YN strain were cultured in flasks containing 300 ml of medium-199 (Gibco BRL) at 37°C. After five days of culture the fungi were removed by passing the medium through gauze, and ACF was sterilised by passage through Millex-GV 0.22 µm millipore filters (Millipore Products, Bedford, Massachusetts, USA) and stored at -80°C.

PHAGOCYTES

Human AMs were isolated from broncho-alveolar lavage (BAL) fluid of six healthy adult volunteers. BAL was performed with 4 × 50 ml sequential volumes of saline at 37°C. The recovery rate of BAL fluid was 60–65%, and Giemsa staining of cytopsin preparations of BAL suspensions showed that more than 90% comprised AMs.

PMNs were prepared from the heparinised venous blood obtained from eight healthy adult volunteers by dextran sedimentation followed by centrifugation over Ficoll gradient and hypotonic lysis of erythrocytes. After treatment with ACF, the viability of cells was more than 95% as assessed by the trypan blue exclusion test. Cell damage was checked by release of lactate dehydrogenase (LDH) and no difference was found in LDH concentration whether or not they had been treated with ACF.

INGESTION OF SPORES BY AM

Ingestion of spores was assessed as described previously.⁷ Spores of YN strain were added to the AM suspensions (2×10^6 /ml) in medium-199 containing ACF (0%, 5%, and 10%) at an AM:spore ratio of 1:5. Each mixture was incubated in a water bath with shaking at 37°C for 120 minutes. Immediately after incubation, all tubes were washed with cold 0.1% EDTA-PBS. Cells were attached to a glass slide by the cytopsin procedure and stained with May-Grunwald-Giemsa. Both percentage phagocytosis and phagocytic index were calculated by counting 500 AMs per glass slide under a light microscope. Percentage phagocytosis was defined as the percentage of AMs with one or more spores ingested or attached. The phagocytic index was the mean number of spores that had been ingested by or attached to 100 AMs.

INHIBITION OF SPORE GERMINATION BY AM

The ability of AMs to inhibit spore germination was determined by a modification of the procedure of Waldorf *et al.*⁹ One millilitre aliquots of cell suspension in culture medium (RPMI-1640 with 5% heat inactivated AB serum) were seeded into multiwell tissue culture plates (Falcon) and incubated at 37°C in a 5% CO₂ incubator for two hours. Non-adherent cells were discarded by washing with RPMI-1640 and the culture medium containing various concentrations of ACF (0%, 1%, 3%, 5%, and 10%) was added to each well to form monolayers of AMs at a concentration of 5×10^5 /well. AMs were challenged with spores in a ratio of 1:1 and centri-

fuged for five minutes at 100g to facilitate contact between the spores and AMs. Incubation was continued for a total of 10 hours at 37°C in the CO₂ incubator, after which the plates were centrifuged, the supernatant discarded, and 0.5 ml of 2.5% deoxycholate (Sigma, St Louis, Missouri, USA) added to each well to lyse AMs. The plates were stained with lactophenol cotton blue and an inverted microscope (Olympus Optical Co, Tokyo, Japan) was used to visualise spore germination. The inhibition rate was calculated from the percentage of spores which did not germinate out of a total of 500 examined per well.

RELEASE OF O₂⁻ FROM PMNS IN RESPONSE TO HYPHAE

Release of O₂⁻ from PMNs in response to *A. fumigatus* hyphae was assessed by superoxide dismutase (SOD) inhibitable reduction of cytochrome C.¹⁰ Briefly, PMNs were preincubated with medium-199 containing various concentrations of ACF (0%, 10%, 30%, and 50%) for one hour at 37°C and then washed with PBS. PMNs were incubated with hyphae at a ratio of 1:1 in 1 ml of Hank's balanced salt solution (HBSS) containing cytochrome C (120 µmol) (Sigma), with or without 50 µg SOD (Sigma), in a shaking water bath at 37°C for one hour. These reaction tubes were centrifuged and O₂⁻ production was assessed in the supernatants as the difference in absorption from the control at 550 nm measured on a spectrophotometer (Hitachi U-3210, Hitachi Ltd., Tokyo, Japan), and amount of O₂⁻ in nanomoles released per 10⁶ PMNs was calculated. All experiments were performed in quadruplicate.

PMN-INDUCED HYPHAL DAMAGE

Damage of the hyphae caused by PMNs was assessed by a colorimetric assay.⁸ PMNs were pretreated with various concentrations of ACF (0%, 10%, 30%, and 50%) for one hour and washed with PBS. PMNs (6×10^6) were resuspended in HBSS and added to hyphae in a final volume of 1 ml containing 5% heat inactivated AB serum at a PMN:hyphae ratio of 4:1. After one hour of incubation at 37°C in a CO₂ incubator the PMNs were lysed with distilled water. The hyphae were then washed with distilled water and resuspended in 0.4 ml of 0.5 mg/ml 2,3-bis-(2-methoxy-4-nitro-5-sulphenyl)-(2H)-tetrazolium-5-carboxanilide (XTT; Sigma) supplemented with 40 µg/ml coenzyme Q (Sigma). Assay tubes were further incubated for one hour at 37°C in a water bath and 100 µl from each supernatant was transferred to the wells of microtitre plates. Absorption at 450 nm was determined using an ELISA plate reader (Flow Laboratories). The percentage of hyphal damage was defined as $[1 - (A_{450} \text{ of fungi incubated with PMN} - A_{450} \text{ of PMN alone}) / A_{450} \text{ of fungi incubated without PMN}] \times 100\%$.

ANALYSIS OF DATA

All data are presented as mean (SE) values. AM experiments were carried out in duplicate and PMN experiments in quadruplicate on four or five separate occasions. The raw data

Figure 1 Effect of *A. fumigatus* culture filtrate (ACF) on inhibition of spore germination by human alveolar macrophages (AMs). AMs co-cultured with different concentrations of ACF were challenged with spores at 37°C for 10 hours. The AM:spore ratio was 1:1. The percentage of inhibition was determined by calculating the number of spores which did not germinate. The results represent the mean (SE) of four separate experiments, each performed in duplicate. * $p < 0.05$ compared with control (medium alone).

were subjected to repeated measures analysis of variance (ANOVA) and the Dunnett's pairwise multiple comparison t test. A p value of less than 0.05 was considered to be significant between control and ACF treatment.

Results

EFFECT OF ACF ON INHIBITION OF SPORE GERMINATION BY AM

AM cultured with medium alone (control) inhibited germination of *A. fumigatus* spores effectively with an inhibition rate of 90 (0.8)%. On the other hand, treatment with 5% and 10% ACF led to reductions of 29% and 54%, respectively, in the inhibition rate compared with the control ($p < 0.05$; fig 1). The number of AMs recovered from culture supernatant in each well after 10 hours of incubation was always less than 4×10^4 which indicated that the suppressive effect of ACF was not a consequence of loss of AMs. To determine whether ACF has a direct effect on spore germination, spores were incubated in AM-free medium containing ACF at concentrations similar to those used in the AM experiments. ACF did not show any direct effects on spore germination, and germination rates incubated with or without ACF were always >99%.

The effect of ACF on the phagocytic activity of AMs against spores was also evaluated. Percentage phagocytosis and the phagocytic index were 69.1 (7.0)% and 155.6 (45.8) in medium alone, 73.6 (5.0)% and 179.5 (54.1) when pretreated with 5% ACF, and 72.0 (4.4)% and 172.8 (52.1) when pretreated with 10% ACF. Thus, 5–10% ACF had no significant effect on either AM mediated percentage phagocytosis or percentage phagocytic index ($p > 0.05$). It is therefore very likely that the suppressive effect of ACF on the inhibition of spore germination described above was not associated with its effect on the ingestion of spores.

EFFECT OF ACF ON O_2^- RELEASE FROM PMN STIMULATED WITH HYPHAE

The effect of ACF on the oxidative metabolic burst of PMN, measured as the release of O_2^- in response to non-opsonised *A. fumigatus* hyphae,

Figure 2 Effect of *A. fumigatus* culture filtrate (ACF) on O_2^- release by human polymorphonuclear leucocytes (PMNs) in response to *A. fumigatus* hyphae. PMNs were pretreated with various concentrations of ACF at 37°C for one hour. O_2^- release was assessed by cytochrome C reduction assay at a PMN:hyphae ratio of 1:1 and expressed as nmol/10⁶ PMNs/h. The results represent the mean (SE) of five separate experiments, each performed in quadruplicate. * $p < 0.05$ compared with control (medium-199).

Figure 3 Effects of *A. fumigatus* culture filtrate (ACF) on hyphal damage by human PMNs. PMN suspensions pretreated with various concentrations of ACF were co-cultured with *A. fumigatus* hyphae at 37°C for one hour. The PMN:hyphae ratio used in this experiment was 4:1. The percentage of hyphal damage was examined by a colorimetric assay using (2,3)-bis-(2-methoxy-4-nitro-5-sulphenyl)-(2H)-tetrazolium-5-carboxamylide (XTT). The results represent the mean (SE) of five separate experiments, each performed in quadruplicate. * $p < 0.05$ compared with control (medium-199).

was examined. As shown in fig 2, pretreatment with 30% and 50% ACF for one hour significantly suppressed O_2^- release in response to hyphae by 22% and 29%, respectively ($p < 0.05$). Since spontaneous O_2^- release was not affected by ACF, and ACF itself did not stimulate O_2^- release of PMNs,⁷ this suppressive effect of ACF was not due to exhaustion of PMNs by ACF.

EFFECT OF ACF ON PMN INDUCED HYPHAL DAMAGE

We assessed the hyphal damage using the XTT colorimetric metabolic assay, a rapid and quantitative method for assessment of fungal cell damage.⁸ The effects of ACF on PMN induced damage of *A. fumigatus* hyphae are shown in fig 3. Pretreatment with 30% and 50% ACF for one hour significantly suppressed the percentage of hyphal damage by 12% and 16%, respectively ($p < 0.05$). When PMNs and hyphae were incubated with ACF throughout

the one hour experimental period without pretreatment, ACF showed the same suppressive effect on the PMN induced hyphal damage (data not shown). When hyphae were incubated in PMN-free medium containing ACF at the same concentrations as used in the PMN experiments, ACF did not affect the metabolic activity of the hyphae as determined by XTT assays (data not shown).

Discussion

The results of this study show that ACF suppresses inhibition of spore germination of *A. fumigatus* by human AMs without altering their phagocytosing activity against the spores, and that ACF also suppresses human PMN induced *A. fumigatus* hyphal damage apparently parallel with O_2^- release in response to hyphae. Although the present study was performed using a single strain of *A. fumigatus*, limited experiments carried out with other clinical isolates of *A. fumigatus* have shown similar results (data not shown).

Roilides *et al* reported that PMNs can respond to and damage *Aspergillus* hyphae irrespective of opsonisation, and that serum independent receptors of PMNs may contribute to attachment to *Aspergillus* hyphae and stimulation of the oxidative burst in the alveoli where serum opsonisation is poor.¹⁰ Warren *et al* found that PMNs can damage hyphae by releasing reactive oxygen intermediates and that the degree of the damage correlates with the duration and the levels of oxidant production.¹ In this study we found that ACF suppresses O_2^- release from PMNs in response to non-opsonised hyphae and also suppresses PMN mediated hyphal damage at similar concentrations. We assume that *A. fumigatus* derived products circumvent the host defence network, thus enhancing the chance of colonisation, proliferation, and invasion of *A. fumigatus* in human tissues.

It has been suggested that secondary metabolites, especially gliotoxin and extracellular proteases, may contribute to the pathogenic potential of *A. fumigatus*.^{6, 11–13} Müllbacher *et al* reported that a three day culture of *A. fumigatus* generated metabolites with antiphagocytic activity (inhibition of phagocytosis by mouse peritoneal exudate cells, inhibition of adherence to plastic surfaces by rodent peritoneal macrophages and AMs, reduction of the basal rate of H_2O_2 production by human PMNs), and they identified one of the biologically active components as gliotoxin.^{11, 12} Slight *et al* reported that *A. fumigatus* spore diffusates obtained from three hour culture supernatants reduced phagocytic activity (inhibition of oxidative burst by rat AMs and inhibition of production of tumour necrosis factor α by rat AMs). They suggested that the active component of spore diffusate was of small molecular weight (<10 kD) and might have a carbohydrate component.^{14, 15} However, to our knowledge there have been very few studies to date that have reported the suppressive effects of *Aspergillus* products on the antifungal activities

of human phagocytes. This study may be the first to demonstrate the suppressive effects of *Aspergillus* products on antifungal host defences by both human AMs and PMNs.

Amitani *et al* isolated the factors responsible for causing damage to the human respiratory ciliated epithelium from ACF and identified one of the most potent substances as gliotoxin.⁶ We have previously suggested that gliotoxin should be a candidate for the antiphagocytic substances in ACF.⁷ Richard *et al* detected gliotoxin in the lung tissue of turkey poult infected with *A. fumigatus* and suggested the involvement of gliotoxin in the pathogenesis of aspergillosis.¹⁶ These findings suggest that gliotoxin should be one of the most important substances contributing to fungal growth in human tissues.

Additional studies are in progress to clarify which of various *A. fumigatus* derived substances including gliotoxin are the most important in suppressing human phagocytic function.

We thank Professor R D Diamond, Section of Infectious Diseases, University Hospital, Boston University for his helpful advice regarding the assay of hyphal damage and Dr S Ishimura, Tsurumi University for his critical advice regarding the statistical analysis.

- Warren BG. The spectrum of pulmonary aspergillosis. *J Thorac Imaging* 1992;7:56–74.
- Binder RE, Faling LJ, Pugatch RD, *et al*. Chronic necrotizing pulmonary aspergillosis: discrete clinical entity. *Medicine* 1982;61:109–24.
- Calderone R, Diamond RD, Senet J-M, *et al*. Host cell-fungal interactions. *J Med Vet Mycol* 1994;32:151–68.
- Schaffner A. Macrophage-*Aspergillus* interactions. *Immunol Ser* 1994;60:545–52.
- Amitani R, Murayama T, Nawada R, *et al*. *Aspergillus* culture filtrates and sputum sols from pulmonary aspergillosis cause damage to human respiratory ciliated epithelium in vitro. *Eur Respir J* 1995;8:1681–7.
- Amitani R, Taylor G, Elezis E-N, *et al*. Purification and characterization of factors produced by *Aspergillus fumigatus* which affect human ciliated respiratory epithelium. *Infect Immun* 1995;63:3266–71.
- Murayama T, Amitani R, Ikegami Y, *et al*. Suppressive effects of *Aspergillus fumigatus* culture filtrates on human alveolar macrophages and polymorphonuclear leukocytes. *Eur Respir J* 1996;9:293–300.
- Meshulam T, Levitz SM, Christin L, *et al*. A simplified new assay for assessment of fungal cell damage with the tetrazolium dye, (2,3)-bis-(2-methoxy-4-nitro-5-sulphenyl)-(2H)-tetrazolium-5-carboxanilide (XTT). *J Infect Dis* 1995;172:1153–6.
- Waldorf AR, Levitz SM, Diamond RD. In vivo bronchoalveolar macrophage defense against *Rhizopus oryzae* and *Aspergillus fumigatus*. *J Infect Dis* 1984;150:752–60.
- Roilides E, Mertins S, Eddy J, *et al*. Enhancement of oxidative response and damage caused by human neutrophils to *Aspergillus fumigatus* hyphae by granulocyte colony-stimulating factor and gamma interferon. *Infect Immun* 1993;61:1185–93.
- Eichner RD, Salami MAL, Wood PR, *et al*. The effect of gliotoxin upon macrophage function. *Int J Immunopharmacol* 1986;8:789–97.
- Müllbacher A, Waring P, Eichner RD. Identification of an agent in cultures of *Aspergillus fumigatus* displaying anti-phagocytic and immunomodulating activity in vitro. *J Gen Microbiol* 1985;31:1251–8.
- Robinson BWS, Venaille TJV, Mendis AHW, *et al*. Allergens as proteases: an *Aspergillus fumigatus* proteinase directly induces human epithelial cell detachment. *J Allergy Clin Immunol* 1990;85:726–31.
- Nicholson WJ, Slight J, Donaldson K. Inhibition of the transcription factor NF- κ B and AP-1 underlies loss of cytokine gene expression in rat alveolar macrophages treated with a diffusible product from the spores of *Aspergillus fumigatus*. *Am J Respir Cell Mol Biol* 1996;15:88–96.
- Slight J, Nicholson WJ, Mitchell CG, *et al*. Inhibition of the alveolar macrophage oxidative burst by diffusible component from the surface of the spores of the fungus *Aspergillus fumigatus*. *Thorax* 1996;51:389–96.
- Richard JL, DeBey MC. Production of gliotoxin during the pathogenic states in turkey poult by *Aspergillus fumigatus* Fresenius. *Mycopathologia* 1995;129:111–5.