Prediction of pulmonary complications after a lobectomy in patients with non-small cell lung cancer

H Uramoto, R Nakanishi, Y Fujino, H Imoto, M Takenoyama, T Yoshimatsu, T Oyama, T Osaki, K Yasumoto

Abstract

Background—Although the preoperative prediction of pulmonary complications after lung major surgery has been reported in various papers, it still remains unclear.

Methods—Eighty nine patients with stage I-IIIa non-small cell lung cancer (NSCLC) who underwent a complete resection at our institute from 1994-8 were evaluated for the feasibility of making a preoperative prediction of pulmonary complications. All had either a predicted postoperative forced vital capacity (FVC) of $>800 \text{ ml/m}^2$ or forced expiratory volume in one second (FEV₁) of $>600 \text{ ml/m}^2$.

Results—Postoperative complications occurred in 37 patients (41.2%) but no patients died during the 30 day period after the operation. Pulmonary complications occurred in 20 patients (22.5%). Univariate analysis indicated that the factors significantly related to pulmonary complications were FVC $<80\%$, serum lactate dehydrogenase (LDH) level $\geq 230 \text{ U/l}$, and arterial oxygen tension (Pao_2) $<10.6 \text{ kPa}$ (80 mm Hg). In a multivariate analysis the three independent predictors of pulmonary complications were serum LDH $\geq 230 \text{ U/l}$ (odds ratio (OR) 10.5, 95% CI 1.4 to 77.3), residual volume (RV)/total lung capacity (TLC) $\geq 30\%$ (OR 6.0, 95% CI 1.1 to 33.7), and Pao_2 $<10.6 \text{ kPa}$ (OR 5.6, 95% CI 1.4 to 22.2).

Conclusions—The above findings indicate that three factors (serum LDH levels of $\geq 230 \text{ U/l}$, RV/TLC $\geq 30\%$, and Pao_2 $<10.6 \text{ kPa}$) may be associated with pulmonary complications in patients undergoing a lobectomy for NSCLC, even though the patient group was relatively small for statistical analysis of such a diverse subject as pulmonary complications.

(Thorax 2001;56:59-61)

Keywords: non-small cell lung cancer; pulmonary complications; morbidity

Lung cancer is the leading cause of death among cancer patients.¹ The necessity for surgical treatment is increasing because complete surgical removal of a tumour offers the best chance for a cure in patients with non-small cell lung cancer (NSCLC).^{2,3} A lobectomy remains the standard therapeutic option in most patients with early stage NSCLC. Although the operative mortality and morbidity

following a lobectomy has decreased over the past decade, both factors still contribute significantly to the cost of treating such patients. Careful selection of the operative modalities is therefore required to reduce the overall mortality and morbidity.

Clinicians are often faced with the problem of morbidity, especially pulmonary complications, which tend to be impossible to predict. Pulmonary complications after lung resection can have serious consequences because the remaining lung tissue is severely compromised during the immediate postoperative period. In the present study we have reviewed a series of patients with NSCLC to evaluate the feasibility of preoperatively predicting pulmonary complications.

Methods

PATIENTS

From April 1994 to December 1998, 299 patients with primary lung cancer underwent lung resection at the Department of Surgery II, University of Occupational and Environmental Health, Kitakyushu, Japan. Of these, 89 patients with stage I-IIIa NSCLC in whom lobectomy was performed were assessed. All patients underwent the following diagnostic procedures preoperatively: brain computed tomographic (CT) scan, body CT scan, a pulmonary perfusion scintigraphic scan, and a bone scintigraphic scan. Spirometric tests were performed according to established guidelines.⁴ We calculated the predicted residual pulmonary function combined with xenon ventilation/perfusion scans from the ratio of right versus left fractional perfusion. The formula for the predicted residual pulmonary function is as follows:

Predicted residual pulmonary function = pulmonary function $\times$ (1 - ratio of lung perfusion on the resected side $\times$ number of resected pulmonary segments/number of pulmonary segments on the resected side)

In our institution surgery is indicated in patients with a predicted postoperative forced vital capacity (FVC) of $>800 \text{ ml/m}^2$ or a forced expiratory volume in one second (FEV₁) of $>600 \text{ ml/m}^2$. Routine systematic lymphadenectomy of hilar and mediastinal lymph nodes was performed in all cases. The charts were then retrospectively reviewed. The ineligibility factors are shown in table 1. The 19 variables listed in table 2 were evaluated.

POSTOPERATIVE COMPLICATIONS

The pulmonary complications included atelectasis and sputum retention requiring bron-

Department of
Surgery II, School of
Medicine, University
of Occupational and
Environmental Health,
School of Medicine,
1-1 Iseigaoka
Yahatanishi-ku,
Kitakyushu 807-8555,
Japan

H Uramoto
H Imoto
M Takenoyama
T Yoshimatsu
T Oyama
T Osaki
K Yasumoto

Department of
Thoracic Surgery,
Shin-Kokura Hospital,
Kitakyushu, Japan
R Nakanishi

Clinical Epidemiology,
School of Medicine,
University of
Occupational and
Environmental Health,
School of Medicine,
Kitakyushu, Japan
Y Fujino

Correspondence to:
Dr H Uramoto
hidetaka@med.uoeh-u.ac.jp

Received 1 June 2000
Returned to authors
26 July 2000
Revised version received
15 August 2000
Accepted for publication
18 September 2000

Table 1 Ineligible factors

Personal characteristics
Karnofsky performance status: 70% or less
Diabetes mellitus, cardiac disease or others requiring medication
Preoperative chemotherapy and/or radiotherapy
Interstitial change detected by CT scan
Pulmonary function test
Both predicted postoperative FVC of 800 ml/m ² or less and predicted postoperative FEV ₁ of 600 ml/m ² or less
Tumour characteristics
Small cell carcinoma or low grade malignant tumour
Confirmed histological diagnosis of stage IIIB or IV
Surgical characteristics
Residual tumour at the surgical margin
Incomplete systematic lymphadenectomy such as sampling
Less than lobectomy such as wedge resection or partial resection
Extent resection such as bilobectomy or pneumonectomy
Mechanical stapling devices for bronchial stump closure
Combined resection
Previous thoracic surgery such as thoracoplasty
Blood transfusion at operation

CT = computed tomography; FEV₁ = forced expiratory volume in one second; FVC = forced vital capacity.

chocscopy, pneumonia defined by radiographic infiltrates plus at least two of the following: temperature >37.7°C, white blood cell count >10 500/mm³, initiation of antibiotic therapy, and

Table 2 Variables (category) evaluated for morbidity

Preoperative general factors (12 variables)
Age
Sex
Brinkmann index
Serum total protein
Serum albumin
Serum LDH
%FVC
%FEV ₁
FEV ₁ (%)
RV/TLC
Pao ₂
Paco ₂
Surgical factors (three variables)
Operative procedure
Operated site
Existence of adhesion at operation
Oncological factors (four variables)
Pathological T stage
Pathological N stage
Pathological stage
Histological type

LDH = lactate dehydrogenase; FVC = forced vital capacity; FEV₁ = forced expiratory volume in one second; RV = residual volume; TLC = total lung capacity; Pao₂, Paco₂ = arterial oxygen and carbon dioxide tensions.

Table 3 Postoperative complications in patients undergoing a lobectomy with NSCLC

Complication type	No of patients (%)
Any complication	37 (41.2)
Respiratory	20 (22.5)
Atelectasis	8 (9.0)
Air leak	7 (7.9)
Pneumonia	2 (2.2)
Sputum retention	2 (2.2)
Bronchial asthma	2 (2.2)
Chylothorax	1 (1.1)
Empyema	1 (1.1)

Some patients had more than one complication.

Table 4 Univariate analysis of the perioperative variables contributing to the occurrence of pulmonary complications following a lobectomy

Variable	Category	No of patients	No (%) of patients with pulmonary complications	p value
Serum LDH (U/l)	<230	83	16 (19.3%)	0.007
	≥230	6	4 (66.7%)	
FVC (%)	<80	5	3 (60.0%)	0.04
	≥80	84	17 (20.2%)	
Pao ₂ (kPa)	<10.6	15	8 (53.3%)	0.002
	≥10.6	74	12 (16.2%)	

LDH = lactate dehydrogenase; FVC = forced vital capacity; Pao₂ = arterial oxygen tension.

air leak requiring a thoracostomy tube drainage for more than seven days, bronchial asthma, chylothorax, and empyema. Cardiac complications included atrial fibrillation, bradycardia, and premature ventricular contractions, all of which required treatment. Other complications included liver dysfunction, gastric ulcer defined by endoscopy, depression, wound infection, and colitis, which all required treatment.

STATISTICAL ANALYSIS

The χ^2 test was used to examine the association between the incidence of complications and clinicopathological features. When the expected value was <5 the Fisher's exact test was used. Variables with a p value of <0.3 were entered into the multivariate logistic regression program of the SAS.⁵ The cut off values were chosen to reflect extreme values over the normal range, or the clinical usefulness of these parameters. A p value of <0.05 was considered significant.

Results

STUDY POPULATION

The mean (SD) age of the patients was 67.2 (8.6) years (range 42–85). There were 62 men and 27 women. Twenty four patients had squamous cell carcinoma, 60 had adenocarcinoma, three had large cell carcinoma, one had adenosquamous cell carcinoma, and one had large cell neuroendocrine carcinoma. According to the new international staging system for lung cancer⁶ following a complete systematic lymphadenectomy carried out in all patients, the staging was as follows: 43 patients were stage IA, 26 were stage IB, two were stage IIA, six patients were stage IIB, and 12 patients were stage IIIA. Pulmonary function tests showed a restrictive defect in five (5.6%), an obstructive defect in eight (9.0%), and a combined defect in three (3.4%).

COMPLICATIONS

All patients were extubated on the day of the operation. There were no deaths (operative mortality rate 0%) and re-operation was required in only one patient (1.1%) following empyema. No patient needed either mechanical ventilation or a tracheostomy. Overall, postoperative complications occurred in 37 patients (41.2%). Pulmonary complications occurred in 20 patients (22.5%). The postoperative complications and their incidence are shown in table 3.

PREDICTORS OF POSTOPERATIVE COMPLICATIONS

The univariate predictions of pulmonary complications including age, sex, history of smoking, serum total protein, serum albumin, %FEV₁, FEV₁%, RV/TLC, hypercapnia, operative procedure, operated site, the existence of adhesions at operation, the pathological T and N stage, and the histological type were not predictive of pulmonary complications. FVC of <80%, serum LDH levels of ≥230 U/l, and arterial oxygen tension (Pao₂) of <10.6 kPa (80 mm Hg) were predictive of complications (table 4). In a multivariate analysis these three factors were found to be significant independent predictors of pulmonary complications (table 5).

Table 5 Multivariate analysis of the perioperative variables contributing to the occurrence of pulmonary complications following a lobectomy

Variable	Category	Odds ratio	95% confidence interval	p value
Serum LDH (U/l)	≥230	10.530	1.434 to 77.331	0.0207
	<230	1		
RV/TLC (%)	≥30	6.015	1.074 to 33.688	0.0412
	<30	1		
Pao ₂ (kPa)	<10.6	5.584	1.407 to 22.168	0.0145
	≥10.6	1		

LDH = lactate dehydrogenase; RV = residual volume; TLC = total lung capacity; Pao₂ = arterial oxygen tension.

Discussion

Advances in operative techniques and postoperative care have led to a further decline in the postoperative complication rates following a lobectomy with a mortality rate of 1–9% reported in recent studies^{7–9} and morbidity occurring at an alarming rate of 11–47%.^{8–10} These mortality and morbidity data are generally obtained from eligible patients who are selected on the basis of their pulmonary function and other organ functions. We also selected surgical candidates on the basis of either a predicted postoperative FVC of >800 ml/m² or an FEV₁ of >600 ml/m². The resulting 0% mortality suggests that our criteria appear to be generally acceptable. However, some morbidity still remains even in these selected patients. We therefore evaluated whether it is possible to predict pulmonary complications after a lobectomy because even patients eligible for surgery develop pulmonary complications.

Many investigators have tried to identify the factors that predict the occurrence of postoperative complications following a lobectomy. Previous studies have shown a relationship between perioperative pulmonary complications and preoperative variables including age,^{8–10} sex,⁸ obesity,¹¹ chronic obstructive pulmonary disease,^{12–13} cigarette smoking,¹⁰ the measurement of maximum oxygen consumption during exercise,¹⁴ FEV₁,⁸ %FEV₁,¹⁰ predicted FEV₁,¹⁵ and transfer factor (TLCO).^{16–17} However, these factors are not yet applicable at all institutions because the eligibility for surgery at each institution is different.

Our results suggest that the serum LDH level is strongly associated with postoperative morbidity. To our knowledge, only one report to date has previously evaluated the LDH level for predicting postoperative morbidity.¹⁸ A high LDH level indicates interstitial fibrosis of the lung following alveolar damage. No interstitial changes were detected on the CT scan and no liver dysfunction was seen in patients with a high LDH level in our series, although we did not investigate the isozyme of LDH. A high LDH level may indicate minimal interstitial damage of the lung which is not seen radiographically. We suggest that measurement of the LDH level, which is widely available, inexpensive, and quick and easy to perform, may have the potential to predict the occurrence of pulmonary complications.

In general, respiratory muscle weakness and semi-starvation is observed in patients following surgery. The contributing factors in these patients are postulated to be malnutrition related to biochemical, physiological, and ana-

tomical changes, reduced blood flow to the respiratory muscles, and increased RV. An increase in RV means that the lung is still hyperinflated even after maximal expiratory effort, thereby increasing the lower secretion of sputum resulting from impaired movement of the diaphragm. An increased RV/TLC may thus also have an impact on the gas component in the alveolar space.

A low Pao₂, which reflects diffusion impairment, unevenness of perfusion/ventilation, or existence of shunt, was independently associated with the occurrence of major postoperative pulmonary complications. Careful management may therefore be required for patients who simultaneously have a high level of LDH, a raised RV/TLC, and a low Pao₂ because these three factors seem to be correlated. In these patients preoperative pulmonary rehabilitation should be performed to prevent pulmonary complications, including the immediate cessation of smoking, use of bronchodilators and/or antibiotics, hydration, and chest physiotherapy.

In conclusion, LDH levels of ≥230 U/l, RV/TLC of ≥30%, and Pao₂ of <10.6 kPa (80 mm Hg) are the most predictive criteria for postoperative morbidity in surgical candidates who had both a predicted postoperative FVC of >800 ml/m² and FEV₁ of >600 ml/m². An accurate prediction of postoperative morbidity would improve patient selection for operative procedures and also help to reduce the overall medical costs related to the treatment of such patients. Further studies in a prospective setting with a larger series of patients are required.

- Landis SH, Murry T, Bolden S, et al. Cancer statistics, 1998. *CA Cancer J Clin* 1998;48:6–29.
- Damstrup L, Poulson HS. Review of the curative role of radiotherapy in the treatment of non-small cell lung cancer. *Lung Cancer* 1994;11:153–78.
- Grilli R, Oxman AD, Julian JA. Chemotherapy for advanced non-small cell lung cancer; how much benefit with enough? *J Clin Oncol* 1993;11:1855–72.
- American Thoracic Society. Standardization of spirometry: 1987 update. *Am Rev Respir Dis* 1987;136:1285–98.
- SAS Institute Inc. *SAS/STAT User's Guide*, Release 6.03. Cary, NC: SAS Institute, 1988.
- Mountain CF. Revisions in the international system for staging lung cancer. *Chest* 1997;111:1710–7.
- Ginsberg RJ, Hill LD, Eagan RT, et al. Modern thirty-day operative mortality for surgical resections for lung cancer. *J Thorac Cardiovasc Surg* 1983;86:654–8.
- Keagy BA, LoresME, Starek PJK, et al. Elective pulmonary lobectomy: factors associated with mortality and operative morbidity. *Ann Thorac Surg* 1985;40:349–52.
- Nagasaki F, Flehinger BJ, Martini N. Comparisons of surgery treatment of lung cancer. *Chest* 1982;82:25–9.
- Dales RE, Dionne G, Leech JA, et al. Preoperative prediction of pulmonary complications following thoracic surgery. *Chest* 1993;104:155–9.
- Tris GM. Preoperative evaluation of pulmonary function. Validity, indications, and benefits. *Am Rev Respir Dis* 1979;119:293–310.
- Tarah S, Moffitt EA, Sessler AD, et al. Risk of anesthesia and surgery in patient with chronic bronchiolitis and chronic obstructive pulmonary disease. *Surgery* 1973;74:720–66.
- Williams CD, Brenowitz JB. "Prohibitive" lung function and major surgical procedures. *Am J Surg* 1976;132:763–6.
- Smith TP, Kinasevitz GT, Tucker WY, et al. Exercise capacity as a predictor of post-thoracotomy morbidity. *Am Rev Respir Dis* 1984;129:730–4.
- Kearney DJ, Lee TH, Reilly JJ, et al. Assessment of operative risk in patients undergoing lung resection. *Chest* 1994;105:753–9.
- Markos J, Mullan BP, Hillman DR, et al. Preoperative assessment as a predictor of mortality and morbidity after lung resection. *Am Rev Respir Dis* 1989;139:902–10.
- Fergusom MK, Little L, Rizzo L, et al. Diffusing capacity predicts morbidity and mortality after pulmonary resection. *J Thorac Cardiovasc Surg* 1988;96:894–900.
- Mitsudomi T, Mizoue T, Yoshimatsu T, et al. Postoperative complications after pneumonectomy for treatment of lung cancer: multivariate analysis. *J Surg Oncol* 1996;61:218–22.