REVIEW SERIES

Chronic obstructive pulmonary disease • 1: Susceptibility factors for COPD the genotype–environment interaction

A J Sandford, E K Silverman

Thorax 2002;57:736–741

Genetic factors including α_1 -antitrypsin deficiency are important in COPD. Candidate gene association studies in COPD are reviewed. Efforts to identify interactions between genetic factors and environmental determinants such as smoking may lead to improved understanding of the pathogenesis of the disease.

Chronic obstructive pulmonary disease (COPD) is a complex disease which is influenced by genetic factors, environmental influences, and genotype–environment interactions. Genotype–environment interactions have been of interest to geneticists for decades. In 1946 Haldane noted that “the interaction of nature and nurture is one of the central problems in genetics”, and it remains so today.¹ A genotype–environment interaction can be defined as a non-additive contribution of genetic and environmental factors to the expression of a phenotype.² Thus, the presence of both genetic and environmental influences on a phenotype does not necessarily imply a genotype–environment interaction; non-additivity of these influences must be present, and non-additivity is observed on a certain scale of measurement. The scale of measurement of clinical variables may not correspond to the scale on which genetic and environmental factors act. Transformations of scale (e.g. logarithmic) may remove genotype–environment interactions, which can be useful for analytical approaches that assume that no genotype–environment interaction is present.³

It is not necessary to know the specific genetic and environmental influences on a phenotype to determine if a genotype–environment interaction is present. Many agricultural studies have compared crop strains in different geographical sites, providing evidence for a genotype–environment interaction without identifying the specific genetic or environmental differences involved.⁴ Of course, identification of the specific genes and the specific environmental factors that interact is of great interest in human genetics, and progress in the human genome project is making the identification of such interactions possible.

Cigarette smoking is clearly the major environmental determinant of COPD; the ability to quantify exposure to cigarette smoke with reasonable accuracy using a questionnaire provides unique opportunities for the study of genotype–environment interactions. For many other complex human disorders, environmental factors are probably important but are difficult to measure. In addition to cigarette smoking, a number of other environmental factors may also be impor-

tant in the development of COPD. Recent work has focused on latent adenoviral infections as a potential environmental trigger for COPD⁵; in addition, childhood respiratory infections, air pollution, and occupational exposures have been suggested as environmental determinants.⁶ In this review we will discuss the genetics of α_1 -antitrypsin (AAT) deficiency, familial aggregation of COPD unrelated to AAT deficiency, and association studies of candidate genes in COPD in the context of genotype–environment interactions.

SEVERE AAT DEFICIENCY

Severe AAT deficiency usually results from a PI ZZ or PI Znull genotype, which is typically diagnosed as a PI Z phenotype by isoelectric focusing of serum. PI Z is frequently associated with the development of early onset COPD. Several large series of AAT deficient individuals have clearly shown that PI Z subjects who smoke cigarettes tend to develop more severe pulmonary impairment at an earlier age than non-smoking PI Z individuals.^{7–9} In addition, a follow up study of participants in the Swedish AAT Deficiency Register has shown that the rate of decline in forced expiratory volume in 1 second (FEV₁) is significantly higher in PI Z current smokers than in never smokers or ex-smokers.¹⁰

A significant fraction of the variability in pulmonary function among PI Z individuals is certainly explained by cigarette smoking; however, the development of COPD in PI Z subjects, even among current or ex-smokers, is not absolute. Silverman and colleagues¹¹ studied 52 PI Z subjects and found that index PI Z subjects, who were tested for AAT deficiency because they had COPD and who were the first PI Z identified in their family, all had significantly reduced FEV₁ values. Many non-index PI Z subjects, who were ascertained by various other means including family studies, population screening, and liver disease, had preserved pulmonary function. By comparing the slopes of the relationship between FEV₁ and pack years of smoking in PI Z, PI MZ, and PI M subjects, Silverman and colleagues showed that there was a significant genotype–environment interaction between PI type and pack years of smoking (fig 1).¹² However, the PI $\times$ smoking interaction was influenced by ascertainment; if only index PI Z subjects were included, all of whom had profoundly reduced pulmonary function, the greater sensitivity of PI Z subjects to increased smoking intensity could not be shown.

With subjects identified from the Danish AAT Register, Seersholm and colleagues also found significantly higher FEV₁ values in non-index PI Z

See end of article for authors' affiliations

Correspondence to: Dr A J Sandford, University of British Columbia McDonald Research Laboratories/iCAPTURE Center, St Paul's Hospital, Vancouver, BC, Canada; asandford@mrl.ubc.ca

Figure 1 FEV₁ as a function of pack years of smoking in non-index PI Z subjects (solid line), PI MZ subjects (dotted line), and PI M subjects (dashed line) in the Washington University AAT study. The steeper decline in FEV₁ in response to pack years in PI Z subjects compared with PI M and PI MZ subjects indicates a significant genotype–environment interaction. Adapted from Silverman *et al.*¹²

subjects (not identified because they had COPD) than in index PI Z subjects (identified because they had COPD), despite similar ages and smoking histories.¹³ Moreover, Seersholm *et al* showed that index PI Z subjects had markedly reduced survival compared with non-index PI Z subjects.¹⁴

Few studies have considered whether factors other than smoking influence the development of lung disease. Black and Kueppers¹⁵ studied 54 PI Z individuals and found significant variability in pulmonary function and clinical symptoms, especially among non-smoking PI Z individuals. Piitulainen *et al*¹⁶ studied more than 200 non-smoking PI Z subjects from the Swedish AAT Deficiency Register, so they were able to examine the impact of risk factors for reduced pulmonary function without confounding by smoking. Not surprisingly, increasing age was associated with reduced pulmonary function in these non-smoking PI Z subjects. Male sex, wheezing symptoms, and occupational exposures to gas, fumes, or dust were also associated with reduced pulmonary function. Subsequent analyses in this population suggested that agricultural occupation and use of a kerosene heater were associated with reduced FEV₁ levels; however, the generalisability of these findings may be limited since only a small number of PI Z subjects had these exposures.¹⁷

Mayer and colleagues recently assessed occupational exposures as a potential contributor to variable expression of lung disease in 128 PI Z subjects.¹⁸ They found that high mineral dust exposure was associated with reduced FEV₁ and chronic cough, independent of smoking history. Although genotype–environment interactions have not been formally demonstrated, occupational and other environmental exposures may also be important determinants of the development of lung disease in PI Z subjects.

ROLE OF OTHER GENETIC FACTORS IN COPD

Familial aggregation of spirometric parameters and COPD

Studies of pulmonary function measurements performed in the general population and in twins have suggested that genetic factors influence variation in pulmonary function. Redline and colleagues assessed spirometric parameters in twins who were not selected for respiratory problems.¹⁹ For FEV₁, a higher correlation in monozygotic twins (0.72) than in dizygotic twins (0.27) suggested that genetic factors influence variation in pulmonary function. Although such general population studies suggest that genetic factors influence pulmonary function, they do not address whether genetic factors influence the development of chronic airflow obstruction.

However, studies in relatives of patients with COPD have supported a role for genetic factors in the disease. Several studies in the 1970s and 1980s reported higher rates of airflow obstruction in first degree relatives of patients with COPD

than in control subjects.^{20–22} For example, Kueppers *et al* found that the mean FEV₁ in siblings of patients with COPD (90% of predicted) was significantly lower than in the siblings of matched control subjects (103% of predicted).²³ Additional studies have shown familial aggregation for chronic bronchitis.^{24–25}

More recently, Silverman and colleagues have focused on subjects with severe early onset COPD unrelated to severe AAT deficiency.²⁶ Proband in this study had FEV₁ <40% predicted at age <53 years. Among first degree relatives of early onset COPD probands, highly significant differences in FEV₁ and FEV₁/FVC were found when current or ex-smoking first degree relatives of early onset COPD probands were compared with lifelong non-smoking control subjects. This pattern of smoking related susceptibility, which was confirmed by multivariate analysis adjusting for age and pack years of smoking, would be consistent with genetic risk factors which interact with cigarette smoking to result in COPD. A similar pattern of smoking related susceptibility was also seen for chronic bronchitis and bronchodilator responsiveness.^{26–27}

Segregation analysis and linkage analysis of pulmonary function and COPD

Segregation analysis, a statistical genetic method that determines whether the pattern of transmission of a phenotype within families is consistent with a major gene effect, has been applied to pulmonary function measurements in several studies. However, the results of these studies have been inconsistent. Using segregation analysis, Rybicki and colleagues²⁸ found that a major gene influencing FEV₁ was suggested in 85 COPD families but not in 56 non-COPD families. In segregation analysis of 455 randomly ascertained families from the NHLBI Family Heart Study, Wilk *et al*²⁹ found evidence for a dominant major gene influencing FEV₁. However, Chen *et al*³⁰ performed segregation analysis of FEV₁ in 214 randomly selected families and found no evidence for a major gene influence. Similarly, Givelber and colleagues found no evidence for a major gene influence on FEV₁ in segregation analysis of 1408 families from the Framingham Study.³¹

Linkage studies have only recently been studied in COPD.^{31a–31b} The absence of an accepted genetic model for the transmission of COPD from segregation analysis suggests that non-parametric linkage analysis methods will probably be required for COPD related phenotypes. Progress in linkage analysis may provide useful guidance for the selection of candidate genes for association studies.

ASSOCIATION STUDIES OF CANDIDATE GENES

Proteases

Heterozygotes for PI deficiency alleles

The issue of whether heterozygotes for PI deficiency alleles are at increased risk for the development of COPD is contentious. PI MS and PI MZ heterozygotes have reductions in serum AAT levels to ~80% and 60% of normal, respectively. PI SZ compound heterozygotes are rare but have levels even lower at ~40% of normal. PI SZ individuals (n=50) were shown to have “similar” levels of airflow obstruction to PI Z individuals, but only if they were smokers.³² In contrast, the frequency of the PI SZ genotype in 702 patients with COPD was not significantly different from that in the general population.³³

Several case-control studies have shown an increased prevalence of PI MZ heterozygotes in patients with COPD compared with controls.³⁴ In one of the largest studies, 526 German patients with COPD were compared with two control populations from the same country.³⁵ There was a significant increase in the frequency of PI MZ heterozygotes in the patients (6%) compared with controls (1%), yielding an odds

ratio (OR) of 5. In other studies the cases and controls were often not selected from the same population, raising concerns of spurious associations due to population stratification.

An alternative approach to case-control studies is to phenotype a general population sample for *PI* variants and compare the prevalence of COPD in *PI* MZ individuals with that in *PI* M individuals.³⁴ Many of these studies are based on small numbers of individuals and have insufficient power to detect an effect of the *PI* MZ or *PI* MS phenotype. In a recent large cohort study the prevalence of obstructive pulmonary disease—defined as hospital discharge diagnosis of either asthma, chronic bronchitis or emphysema—was compared in 1551 *PI* MZ individuals and 14 484 controls from the general population of unknown *PI* genotype.³⁶ The risk for obstructive pulmonary disease was significantly increased in the *PI* MZ individuals compared with the controls (relative risk (RR) = 2). However, only first degree relatives of *PI* Z COPD patients had an increased risk, suggesting that other genetic or environmental factors contribute to the increased risk in these patients. Most recently, Dahl *et al*³⁷ have performed a large cross sectional study of 9187 individuals from the general population. Each individual underwent pulmonary function testing and was genotyped for the S and Z *PI* variants. Of the genotypic groups, only the *PI* SZ and *PI* ZZ individuals showed an increased prevalence of airflow obstruction (FEV₁ <80% predicted). Neither the *PI* MS or *PI* MZ genotype was associated with a lower level of lung function in individuals without COPD. However, in patients with COPD FEV₁ was 655 ml less in *PI* MZ individuals than in *PI* M individuals after adjustment for confounding variables. The observation that the *PI* MZ genotype was only associated with reduced lung function in those with COPD again suggests that other predisposing factors must exist.

3' *PI* mutation

Several polymorphisms of the *PI* gene are not associated with AAT deficiency—for example, a G→A transition in the 3' region of the gene. The A allele of this polymorphism was associated with COPD in some populations^{38, 39} but not in others.^{40, 41} The A allele was also associated with decreased transcription factor binding and decreased gene expression in vitro.^{42, 43} The 3' mutation could therefore affect the acute phase response leading to attenuated upregulation of AAT synthesis when inflammation is present. In contrast to the in vitro data, the 3' polymorphism was not associated with a reduced AAT acute phase response in patients undergoing open heart surgery⁴⁴ or in patients who had cystic fibrosis.⁴⁵ The role of the 3' polymorphism in the pathogenesis of COPD therefore remains unclear.

α_1 -Antichymotrypsin

α_1 -Antichymotrypsin (AAT) is a protease inhibitor that is secreted by hepatocytes and alveolar macrophages. The gene contains several polymorphisms, and the Ala variant of an amino acid substitution (Pro²²⁷→Ala) that causes AAT deficiency was associated with COPD in one population⁴⁶ but not others.^{41, 47} This variant is rare and larger cohorts will be required to identify its role definitively in the pathogenesis of COPD. A recent Japanese study of 53 patients with COPD and 65 controls matched for age, sex, and smoking reported that homozygotes for the Ala variant of an amino acid substitution (Ala-15Thr) in the AAT signal peptide were increased in the patients compared with the controls (OR=3).⁴⁸ This association is difficult to interpret since the Ala variant was not associated with AAT deficiency but it may reflect linkage disequilibrium with another allele.

Xenobiotic metabolising enzymes

Figure 2 summarises the function of the xenobiotic metabolising enzymes that have been investigated as risk factors for the development of COPD.

Figure 2 Overview of xenobiotic metabolism focusing on genes investigated as candidates for COPD. X_1 , X_2 , and X_3 are different xenobiotic substrates. GSTs = glutathione S-transferases.

Microsomal epoxide hydrolase

Microsomal epoxide hydrolase (EPHX1) is an enzyme that plays an important role in the metabolism of highly reactive epoxide intermediates formed in cigarette smoke. EPHX1 is expressed in several different cell types including hepatocytes and bronchial epithelial cells. Two common polymorphisms occur in the *EPHX1* gene: Tyr¹¹³→His and His¹³⁹→Arg. These polymorphisms correlated with the level of EPHX1 enzymatic activity in transfected cell lines⁴⁹ but not in liver tissue samples.⁵⁰ The slow metabolising form of EPHX1 was found in a higher proportion of patients with emphysema (22%) and COPD (19%) than in controls (6%), giving an odds ratio of 5.⁵¹ In a smaller Japanese study the slow metabolising form of EPHX1 was associated with more severe COPD.⁵² These results were not confirmed in a Korean population.⁵³ However, *EPHX1* genotypes were associated with rate of decline of lung function in smokers.⁵⁴

Glutathione S-transferases

Glutathione S-transferases (GSTs) are enzymes that play an important role in detoxifying various aromatic hydrocarbons found in cigarette smoke. *GSTM1* is expressed in the liver and the lung. Homozygous deletion of the *GSTM1* gene occurs in approximately 50% of Caucasians and was associated with emphysema in patients who had lung cancer (OR=2)⁵⁵ and with severe chronic bronchitis in heavy smokers (OR=3).⁵⁶ However, in a Korean study there was no association of *GSTM1* and *GSTT1* polymorphisms with COPD.⁵³

GSTP1 is expressed in the same cell types as *GSTM1*, although at a higher level.⁵⁷ There is a polymorphism at position 105 (Ile¹⁰⁵→Val), leading to an increased catalytic activity of the enzyme in vitro.⁵⁸ Homozygotes for the isoleucine allele were significantly increased in Japanese patients with COPD compared with controls (OR=3).⁵⁹

Antioxidants

Heme oxygenase-1

Heme oxygenase-1 (HMOX1) degrades heme to biliverdin and has been found to provide cellular protection against heme and non-heme mediated oxidant injury.^{60, 61} A microsatellite (GT)_n polymorphism within the heme oxygenase gene (*HMOX1*) promoter was associated with emphysema in Japanese smokers.⁶² In this study of 100 patients with emphysema and 101 matched controls, alleles with a high number of GT repeats (≥30) were more prevalent in the cases (21%) than in the controls (10%) yielding an odds ratio of 2. The authors hypothesised that the association was due to the effect of

Table 1 Genes investigated as candidates for COPD

Gene	References
Alpha-1 antitrypsin	7–18, 32–45
Alpha-1 antichymotrypsin	41, 46–48
Microsomal epoxide hydrolase	51–54
Glutathione S-transferase M1	53, 55, 56
Glutathione S-transferase T1	53
Glutathione S-transferase P1	59
Heme oxygenase-1	62
Tumour necrosis factor- α	54, 64–68
Cystic fibrosis transmembrane conductance regulator	69–71
Cytochrome P4501A1	77
ABH secretor, Lewis and ABO blood groups	78–84
HLA	82, 85, 86
Vitamin D binding protein	87, 88
Immunoglobulin deficiency	89–91

purine-pyrimidine repeats in promoting the formation of left handed Z DNA, thus downregulating gene expression.⁶³ A high number of GT repeats in the *HMOX1* promoter could suppress expression of the gene and leave the lung susceptible to oxidant injury. In support of this hypothesis, reporter gene constructs with low numbers of GT repeats upregulated gene expression in response to hydrogen peroxide whereas constructs with a higher number of repeats did not.⁶²

Inflammatory mediators

Tumour necrosis factor α

Tumour necrosis factor α (TNF α) is a proinflammatory cytokine that has many effects relevant to the pathogenesis of COPD including neutrophil release from the bone marrow and neutrophil activation. The *TNF* gene contains a G→A transition in the promoter (*TNF* G-308A) that was associated with the level of TNF production in vitro.⁶⁴ An association of the *TNF* -308A allele with COPD was found in a Taiwanese population (OR=11).⁶⁵ The patients were selected based on the presence of chronic bronchitis and impaired lung function (FEV₁ <80% predicted and FEV₁/FVC <0.69). This association was confirmed recently in a Japanese study in which 106 patients with an FEV₁ of <80% predicted and an FEV₁/FVC of <0.7 were compared with 110 asymptomatic smokers/ex-smokers and 129 adult blood donors.⁶⁶ *TNF* -308A was significantly increased in the cases compared with both the control groups (OR=3). However, no association of the -308A allele with COPD was found in a study of 53 patients with COPD and 65 controls from a Japanese population.⁶⁷ Studies of Caucasian populations have found no association of *TNF* -308A with COPD⁶⁸ or rate of decline of lung function.³⁴

Mucociliary clearance

Cystic fibrosis transmembrane regulator

In 1989 mutations in the cystic fibrosis transmembrane conductance regulator (*CFTR*) gene were identified as the cause of cystic fibrosis (CF). CF carriers may also be predisposed to respiratory disease. IVS8 is a *CFTR* polymorphism consisting of a variable length thymine repeat in intron 8 of the gene. The IVS8-5T allele results in reduced *CFTR* gene expression. Studies of IVS8-5T as a risk factor for COPD have yielded conflicting results.^{69–70} A recent study by Tzetzis *et al.*⁷¹ in a Greek population of 12 patients with obstructive lung diseases and 52 controls screened for variants in the entire *CFTR* coding region found no significant increase in CF-causing mutations in the patients compared with the controls. The frequency of the Met allele of the Met470Val polymorphism was increased in the patients (71%) compared with the controls (36%). However, the Met470 variant is associated with increased *CFTR* chloride channel activity⁷² and therefore the reason for the association with COPD is unclear.

Other genes

Several other genes have been investigated as risk factors for the development of COPD and these are listed in table 1.

FUTURE STUDIES OF GENOTYPE–ENVIRONMENT INTERACTION

In most of the studies discussed in this review there has been little specific consideration of genotype–environment interactions. Of interest, segregation analysis and linkage analysis typically assume that no genotype–environment interaction is present. Approaches which include genotype–environment interactions in segregation analysis have been developed, but they have not been applied in COPD.⁷³ New methods of including genotype–environment interaction in linkage analysis and family based association analysis have also been developed, which may be useful in future studies of the genetics of COPD.^{73–74} In addition to assessing the significance of genotype–environment interactions, inclusion of these interactions in linkage and association studies may actually improve the detection of main genetic effects because the relationship between genetic and environmental factors is modelled more accurately.

The most commonly performed design used in the study of genetics in COPD has been the case-control genetic association study. Future association studies may also be strengthened by the inclusion of environmental risk factors in the study design and analysis. The methodology employed could be a case-control approach using multivariate regression models to examine interactions. An alternative methodology could be to use a case only design⁷⁵ if the genotype and environmental risk factors are independent.⁷⁶

In conclusion, genotype–environment interactions are likely to be essential contributors to the development of COPD. Intensified efforts to identify these interactions may lead to improved understanding of the pathogenesis of the disease.

Authors' affiliations

A J Sandford, University of British Columbia McDonald Research Laboratories/iCAPTURE Center, St Paul's Hospital, Vancouver, BC, Canada

E K Silverman, Channing Laboratory and Pulmonary and Critical Care Division, Department of Medicine, Brigham and Women's Hospital, Boston, MA 02115, USA

This work was supported by NIH grants HL 61575 and HL 68926 (EKS) and a Parker B Francis Fellowship (AJS).

REFERENCES

- 1 Haldane JBS. The interaction of nature and nurture. *Ann Eugen* 1946;**13**:197–205.
- 2 Hartl DL, Clark AG. *Principles of population genetics*. Sunderland, MA: Sinauer Associates, 1997.
- 3 Falconer DS, Mackay TFC. *Introduction to quantitative genetics*. Harlow: Longman, 1996.
- 4 Westcott B. Some methods of analysing genotype-environment interaction. *Heredity* 1986;**56**:243–53.
- 5 Retamales I, Elliott WM, Meshi B, *et al*. Amplification of inflammation in emphysema and its association with latent adenoviral infection. *Am J Respir Crit Care Med* 2001;**164**:469–73.
- 6 Silverman EK, Speizer FE. Risk factors for the development of chronic obstructive pulmonary disease. *Med Clin North Am* 1996;**80**:501–22.
- 7 Tobin MJ, Cook PJ, Hutchison DC. Alpha 1 antitrypsin deficiency: the clinical and physiological features of pulmonary emphysema in subjects homozygous for Pi type Z. A survey by the British Thoracic Association. *Br J Dis Chest* 1983;**77**:14–27.
- 8 Janus ED, Phillips NT, Carrell RW. Smoking, lung function, and α_1 -antitrypsin deficiency. *Lancet* 1985;**i**:152–4.
- 9 Larsson C. Natural history and life expectancy in severe alpha1-antitrypsin deficiency, Pi Z. *Acta Med Scand* 1978;**204**:345–51.
- 10 Piitulainen E, Eriksson S. Decline in FEV₁ related to smoking status in individuals with severe alpha1-antitrypsin deficiency (PiZZ). *Eur Respir J* 1999;**13**:247–51.
- 11 Silverman EK, Pierce JA, Province MA, *et al*. Variability of pulmonary function in alpha1-antitrypsin deficiency: clinical correlates. *Ann Intern Med* 1989;**111**:982–91.
- 12 Silverman EK, Province MA, Campbell EJ, *et al*. Family study of alpha 1-antitrypsin deficiency: effects of cigarette smoking, measured genotype,

- and their interaction on pulmonary function and biochemical traits. *Genet Epidemiol* 1992;9:317-31.
- 13 **Seersholm N**, Kok-Jensen A, Dirksen A. Decline in FEV₁ among patients with severe hereditary alpha 1-antitrypsin deficiency type PiZ. *Am J Respir Crit Care Med* 1995;152:1922-5.
 - 14 **Seersholm N**, Kok-Jensen A. Survival in relation to lung function and smoking cessation in patients with severe hereditary alpha 1-antitrypsin deficiency. *Am J Respir Crit Care Med* 1995;151:369-73.
 - 15 **Black LF**, Kueppers F. α_1 -antitrypsin deficiency in nonsmokers. *Am Rev Respir Dis* 1978;117:421-8.
 - 16 **Piitulainen E**, Tornling G, Eriksson S. Effect of age and occupational exposure to airway irritants on lung function in non-smoking individuals with alpha 1-antitrypsin deficiency (PiZZ). *Thorax* 1997;52:244-8.
 - 17 **Piitulainen E**, Tornling G, Eriksson S. Environmental correlates of impaired lung function in non-smokers with severe alpha 1-antitrypsin deficiency (PiZZ). *Thorax* 1998;53:939-43.
 - 18 **Mayer AS**, Stoller JK, Bucher Bartelson B, et al. Occupational exposure risks in individuals with Pi*Z alpha(1)-antitrypsin deficiency. *Am J Respir Crit Care Med* 2000;162:553-8.
 - 19 **Redline S**, Tishler PV, Rosner B, et al. Genotypic and phenotypic similarities in pulmonary function among family members of adult monozygotic and dizygotic twins. *Am J Epidemiol* 1989;129:827-36.
 - 20 **Larson RK**, Barman ML, Kueppers F, et al. Genetic and environmental determinants of chronic obstructive pulmonary disease. *Ann Intern Med* 1970;72:627-32.
 - 21 **Cohen BH**, Diamond EL, Graves CG, et al. A common familial component in lung cancer and chronic obstructive pulmonary disease. *Lancet* 1977;ii:523-6.
 - 22 **Cohen BH**. Chronic obstructive pulmonary disease: a challenge in genetic epidemiology. *Am J Epidemiol* 1980;112:274-88.
 - 23 **Kueppers F**, Miller RD, Gordon H, et al. Familial prevalence of chronic obstructive pulmonary disease in a matched pair study. *Am J Med* 1977;63:336-42.
 - 24 **Higgins M**, Keller J. Familial occurrence of chronic respiratory disease and familial resemblance in ventilatory capacity. *J Chron Dis* 1975;28:239-51.
 - 25 **Tager I**, Tishler PV, Rosner B, et al. Studies of the familial aggregation of chronic bronchitis and obstructive airways disease. *Int J Epidemiol* 1978;7:55-62.
 - 26 **Silverman EK**, Chapman HA, Drazen JM, et al. Genetic epidemiology of severe, early-onset chronic obstructive pulmonary disease. Risk to relatives for airflow obstruction and chronic bronchitis. *Am J Respir Crit Care Med* 1998;157:1770-8.
 - 27 **Celedon JC**, Speizer FE, Drazen JM, et al. Bronchodilator responsiveness and serum total IgE levels in families of probands with severe early-onset COPD. *Eur Respir J* 1999;14:1009-14.
 - 28 **Rybicki BA**, Beaty TH, Cohen BH. Major genetic mechanisms in pulmonary function. *J Clin Epidemiol* 1990;43:667-75.
 - 29 **Wilk JB**, Djousse L, Arnett DK, et al. Evidence for major genes influencing pulmonary function in the NHLBI Family Heart Study. *Genet Epidemiol* 2000;19:81-94.
 - 30 **Chen Y**, Horne SL, Rennie DC, et al. Segregation analysis of two lung function indices in a random sample of young families: the Humboldt Family Study. *Genet Epidemiol* 1996;13:35-47.
 - 31 **Givelber RJ**, Courpmitree NN, Gottlieb DJ, et al. Segregation analysis of pulmonary function among families in the Framingham Study. *Am J Respir Crit Care Med* 1998;157:1445-51.
 - 31a **Silverman EK**, Palmer LJ, Mosley JD, et al. Genome wide linkage analysis of quantitative spirometric phenotypes in severe early-onset chronic obstructive pulmonary disease. *Am J Hum Genet* 2002;70:1229-39.
 - 31b **Silverman EK**, Mosley JD, Palmer LJ, et al. Genome-wide linkage analysis of severe, early-onset chronic obstructive pulmonary disease: airflow obstruction and chronic bronchitis phenotypes. *Hum Mol Genet* 2002;11:623-32.
 - 32 **Turino GM**, Barker AF, Brantly ML, et al. Clinical features of individuals with Pi*SZ phenotype of α_1 -antitrypsin deficiency. α_1 -Antitrypsin Deficiency Registry Study Group. *Am J Respir Crit Care Med* 1996;154:1718-25.
 - 33 **Alvarez-Granda L**, Cabero-Perez MJ, Bustamante-Ruiz A, et al. Pi SZ phenotype in chronic obstructive pulmonary disease. *Thorax* 1997;52:659-61.
 - 34 **Sandford AJ**, Weir TD, Paré PD. Genetic risk factors for chronic obstructive pulmonary disease. *Eur Respir J* 1997;10:1380-91.
 - 35 **Bartmann K**, Fooke-Achterra M, Koch G, et al. Heterozygosity in the Pi-system as a pathogenetic cofactor in chronic obstructive pulmonary disease (COPD). *Eur J Respir Dis* 1985;66:284-96.
 - 36 **Seersholm N**, Wilcz JT, Kok-Jensen A, et al. Risk of hospital admission for obstructive pulmonary disease in alpha(1)-antitrypsin heterozygotes of phenotype PiMZ. *Am J Respir Crit Care Med* 2000;161:81-4.
 - 37 **Dahl M**, Nordestgaard BG, Lange P, et al. Molecular diagnosis of intermediate and severe alpha(1)-antitrypsin deficiency: MZ individuals with chronic obstructive pulmonary disease may have lower lung function than MM individuals. *Clin Chem* 2001;47:56-62.
 - 38 **Kalsheker NA**, Watkins GL, Hill S, et al. Independent mutations in the flanking sequence of the α_1 -antitrypsin gene are associated with chronic obstructive airways disease. *Dis Markers* 1990;8:151-7.
 - 39 **Poller W**, Meisen C, Olek K. DNA polymorphisms of the α_1 -antitrypsin gene region in patients with chronic obstructive pulmonary disease. *Eur J Clin Invest* 1990;20:1-7.
 - 40 **Sandford AJ**, Spinelli JJ, Weir TD, et al. Mutation in the 3' region of the α_1 -antitrypsin gene and chronic obstructive pulmonary disease. *J Med Genet* 1997;34:874-5.
 - 41 **Bentazzo MG**, Gile LS, Bombieri C, et al. α_1 -antitrypsin Taq I polymorphism and α_1 -antichymotrypsin mutations in patients with obstructive pulmonary disease. *Respir Med* 1999;93:648-54.
 - 42 **Morgan K**, Scobie G, Kalsheker NA. Point mutation in a 3' flanking sequence of the α_1 -antitrypsin gene associated with chronic respiratory disease occurs in a regulatory sequence. *Hum Mol Genet* 1993;2:253-7.
 - 43 **Morgan K**, Scobie G, Marsters P, et al. Mutation in an α_1 -antitrypsin enhancer results in an interleukin-6 deficient acute-phase response due to loss of cooperativity between transcription factors. *Biochim Biophys Acta* 1997;1362:67-76.
 - 44 **Sandford AJ**, Chagani T, Spinelli JJ, et al. α_1 -antitrypsin genotypes and the acute-phase response to open heart surgery. *Am J Respir Crit Care Med* 1999;159:1624-8.
 - 45 **Mahadeva R**, Westerbeek RC, Perry DJ, et al. α_1 -antitrypsin deficiency alleles and the Taq-I G→A allele in cystic fibrosis lung disease. *Eur Respir J* 1998;11:873-9.
 - 46 **Poller W**, Faber JP, Scholz S, et al. Mis-sense mutation of α_1 -antichymotrypsin gene associated with chronic lung disease. *Lancet* 1992;339:1538.
 - 47 **Sandford AJ**, Chagani T, Weir TD, et al. α_1 -antichymotrypsin mutations in patients with chronic obstructive pulmonary disease. *Dis Markers* 1998;13:257-60.
 - 48 **Ishii T**, Matsuse T, Teramoto S, et al. Association between alpha-1-antichymotrypsin polymorphism and susceptibility to chronic obstructive pulmonary disease. *Eur J Clin Invest* 2000;30:543-8.
 - 49 **Hassett C**, Aicher L, Sidhu JS, et al. Human microsomal epoxide hydrolase: genetic polymorphism and functional expression in vitro of amino acid variants. *Hum Mol Genet* 1994;3:421-8.
 - 50 **Kitteringham NR**, Davis C, Howard N, et al. Interindividual and interspecies variation in hepatic microsomal epoxide hydrolase activity: studies with cis-stilbene oxide, carbamazepine 10, 11-epoxide and naphthalene. *J Pharmacol Exp Ther* 1996;278:1018-27.
 - 51 **Smith CA**, Harrison DJ. Association between polymorphism in gene for microsomal epoxide hydrolase and susceptibility to emphysema. *Lancet* 1997;350:630-3.
 - 52 **Yoshikawa M**, Hiyama K, Ishioka S, et al. Microsomal epoxide hydrolase genotypes and chronic obstructive pulmonary disease in Japanese. *Int J Mol Med* 2000;5:49-53.
 - 53 **Yim JJ**, Park GY, Lee CT, et al. Genetic susceptibility to chronic obstructive pulmonary disease in Koreans: combined analysis of polymorphic genotypes for microsomal epoxide hydrolase and glutathione S-transferase M1 and T1. *Thorax* 2000;55:121-5.
 - 54 **Sandford AJ**, Chagani T, Weir TD, et al. Susceptibility genes for rapid decline of lung function in the Lung Health Study. *Am J Respir Crit Care Med* 2001;163:469-73.
 - 55 **Harrison DJ**, Cantlay AM, Rae F, et al. Frequency of glutathione S-transferase M1 deletion in smokers with emphysema and lung cancer. *Hum Exp Toxicol* 1997;16:356-60.
 - 56 **Baranova H**, Perriot J, Albuissou E, et al. Peculiarities of the GSTM1 0/0 genotype in French heavy smokers with various types of chronic bronchitis. *Hum Genet* 1997;99:822-6.
 - 57 **Cantlay AM**, Smith CA, Wallace WA, et al. Heterogeneous expression and polymorphic genotype of glutathione S-transferases in human lung. *Thorax* 1994;49:1010-4.
 - 58 **Sundberg K**, Johansson AS, Stenberg G, et al. Differences in the catalytic efficiencies of allelic variants of glutathione transferase P1-1 towards carcinogenic diol epoxides of polycyclic aromatic hydrocarbons. *Carcinogenesis* 1998;19:433-6.
 - 59 **Ishii T**, Matsuse T, Teramoto S, et al. Glutathione S-transferase P1 (GSTP1) polymorphism in patients with chronic obstructive pulmonary disease. *Thorax* 1999;54:693-6.
 - 60 **Otterbein LE**, Lee PJ, Chin BY, et al. Protective effects of heme oxygenase-1 in acute lung injury. *Chest* 1999;116:61S-63S.
 - 61 **Choi AM**, Alam J. Heme oxygenase-1: function, regulation, and implication of a novel stress-inducible protein in oxidant-induced lung injury. *Am J Respir Cell Mol Biol* 1996;15:9-19.
 - 62 **Yamada N**, Yamaya M, Okinaga S, et al. Microsatellite polymorphism in the heme oxygenase-1 gene promoter is associated with susceptibility to emphysema. *Am J Hum Genet* 2000;66:187-95.
 - 63 **Naylor LH**, Clark EM. d(TG)n.d(CA)n sequences upstream of the rat prolactin gene form Z-DNA and inhibit gene transcription. *Nucl Acids Res* 1990;18:1595-601.
 - 64 **Bouma G**, Crusius JB, Oudkerk Pool M, et al. Secretion of tumour necrosis factor α and lymphotoxin α in relation to polymorphisms in the TNF genes and HLA-DR alleles. Relevance for inflammatory bowel disease. *Scand J Immunol* 1996;43:456-63.
 - 65 **Huang SL**, Su CH, Chang SC. Tumour necrosis factor- α gene polymorphism in chronic bronchitis. *Am J Respir Crit Care Med* 1997;156:1436-9.
 - 66 **Sakao S**, Tatsumi K, Igari H, et al. Association of tumor necrosis factor alpha gene promoter polymorphism with the presence of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2001;163:420-2.
 - 67 **Ishii T**, Matsuse T, Teramoto S, et al. Neither IL-1beta, IL-1 receptor antagonist, nor TNF-alpha polymorphisms are associated with susceptibility to COPD. *Respir Med* 2000;94:847-51.
 - 68 **Higham MA**, Pride NB, Alikhan A, et al. Tumour necrosis factor-alpha gene promoter polymorphism in chronic obstructive pulmonary disease. *Eur Respir J* 2000;15:281-4.
 - 69 **Pignatti PF**, Bombieri C, Benetazzo M, et al. CFTR gene variant IVS8-5T in disseminated bronchiectasis. *Am J Hum Genet* 1996;58:889-92.

- 70 **Bombieri C**, Benetazzo M, Saccomani A, *et al*. Complete mutational screening of the CFTR gene in 120 patients with pulmonary disease. *Hum Genet* 1998;**103**:718–22.
- 71 **Tzetis M**, Efthymiadou A, Strofalis S, *et al*. CFTR gene mutations—including three novel nucleotide substitutions—and haplotype background in patients with asthma, disseminated bronchiectasis and chronic obstructive pulmonary disease. *Hum Genet* 2001;**108**:216–21.
- 72 **Cuppens H**, Lin W, Jaspers M, *et al*. Polyvariant mutant cystic fibrosis transmembrane conductance regulator genes. The polymorphic (Tg)m locus explains the partial penetrance of the T5 polymorphism as a disease mutation. *J Clin Invest* 1998;**101**:487–96.
- 73 **Gauderman WJ**, Siegmund KD. Gene-environment interaction and affected sib pair linkage analysis. *Hum Hered* 2001;**52**:34–46.
- 74 **Umbach DM**. Invited commentary: on studying the joint effects of candidate genes and exposures. *Am J Epidemiol* 2000;**152**:701–3.
- 75 **Khoury MJ**, Flanders WD. Nontraditional epidemiologic approaches in the analysis of gene-environment interaction: case-control studies with no controls! *Am J Epidemiol* 1996;**144**:207–13.
- 76 **Albert PS**, Ratnasinghe D, Tangrea J, *et al*. Limitations of the case-only design for identifying gene-environment interactions. *Am J Epidemiol* 2001;**154**:687–93.
- 77 **Cantlay AM**, Lamb D, Gillooly M, *et al*. Association between the CYP1A1 gene polymorphism and susceptibility to emphysema and lung cancer. *J Clin Pathol Mol Pathol* 1995;**48**:M210–4.
- 78 **Cohen BH**, Ball WC, Brashears S, *et al*. Risk factors in chronic obstructive pulmonary disease (COPD). *Am J Epidemiol* 1977;**105**:223–32.
- 79 **Beatty TH**, Menkes HA, Cohen BH, *et al*. Risk factors associated with longitudinal change in pulmonary function. *Am Rev Respir Dis* 1984;**129**:660–7.
- 80 **Krzyzanowski M**, Jedrychowski W, Wysocki M. Factors associated with the change in ventilatory function and the development of chronic obstructive pulmonary disease in a 13 year follow-up of the Cracow study. Risk of chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1986;**134**:1011–9.
- 81 **Cohen BH**, Bias WB, Chase GA, *et al*. Is ABH nonsecretor status a risk factor for obstructive lung disease? *Am J Epidemiol* 1980;**111**:285–91.
- 82 **Kauffmann F**, Kleisbauer JP, Cambon-De-Mouzon A, *et al*. Genetic markers in chronic air-flow limitation: a genetic epidemiologic study. *Am Rev Respir Dis* 1983;**127**:263–9.
- 83 **Horne SL**, Cockcroft DW, Lovegrove A, *et al*. ABO, Lewis and secretor status and relative incidence of airflow obstruction. *Dis Markers* 1985;**3**:55–62.
- 84 **Kauffmann F**, Frette C, Pham QT, *et al*. Associations of blood group-related antigens to FEV₁, wheezing, and asthma. *Am J Respir Crit Care Med* 1996;**153**:76–82.
- 85 **Sugiyama Y**, Kudoh S, Maeda H, *et al*. Analysis of HLA antigens in patients with diffuse panbronchiolitis. *Am Rev Respir Dis* 1990;**141**:1459–62.
- 86 **Keicho N**, Tokunaga K, Nakata K, *et al*. Contribution of HLA genes to genetic predisposition in diffuse panbronchiolitis. *Am J Respir Crit Care Med* 1998;**158**:846–50.
- 87 **Horne SL**, Cockcroft DW, Dosman JA. Possible protective effect against chronic obstructive airways disease by the GC 2 allele. *Hum Heredity* 1990;**40**:173–6.
- 88 **Schellenberg D**, Paré PD, Weir TD, *et al*. Vitamin D binding protein variants and the risk of COPD. *Am J Respir Crit Care Med* 1998;**157**:957–61.
- 89 **Bjorkander J**, Bake B, Oxelius VA, *et al*. Impaired lung function in patients with IgA deficiency and low levels of IgG2 or IgG3. *N Engl J Med* 1985;**313**:720–4.
- 90 **O'Keeffe S**, Gzel A, Drury R, *et al*. Immunoglobulin G subclasses and spirometry in patients with chronic obstructive pulmonary disease. *Eur Respir J* 1991;**4**:932–6.
- 91 **Webb DR**, Condemi JJ. Selective immunoglobulin A deficiency and chronic obstructive lung disease. A family study. *Ann Intern Med* 1974;**80**:618–21.