- 18 Jones SE, Kelly DJ, Cox AJ, *et al.* Mast cell infiltration and chemokine expression in progressive renal disease. *Kidney Int* 2003;**64**:906–13.
- 19 Gimenez-Gallego G, Navia MA, Reuben JP, *et al.* Purification, sequence, and model structure of charybdotoxin, a potent selective inhibitor of calcium-activated potassium channels. *Proc Natl Acad Sci USA* 1998;**95**:3329–33.
- 20 Wulff H, Gutman GA, Cahalan MD, *et al.* Delineation of the clotrimazole/TRAM-34 binding site on the intermediate conductance calcium-activated potassium channel, IKCa1. *J Cell Chem* 2001;**276**:32040–5.
- 21 Logsdon NJ, Kang J, Togo JA, *et al.* A novel gene, hKCa4, encodes the calcium-activated potassium channel in human T lymphocytes. *J Biol Chem* 1997;**272**:32723–6.
- 22 Bradding P, Okayama Y, Kambe N, *et al.* Ion channel gene expression in human lung, skin, and cord blood-derived mast cells. *J Leukoc Biol* 2003;**73**:614–20.
- 23 Wulff H, Miller MJ, Hansel W, *et al.* Design of a potent and selective inhibitor of the intermediate-conductance Ca^{2+} -activated K^{+} channel, IKCa1: a potential immunosuppressant. *Proc Natl Acad Sci USA* 2000;**97**:8151–6.
- 24 Schwab A, Schuricht B, Seeger P, *et al.* Migration of transformed renal epithelial cells is regulated by K^{+} channel modulation of actin cytoskeleton and cell volume. *Pflügers Arch* 1999;**438**:330–7.
- 25 Schneider SW, Pagel P, Rotsch C, *et al.* Volume dynamics in migrating epithelial cells measured with atomic force microscopy. *Pflügers Arch* 2000;**439**:297–303.
- 26 Schwab A, Wulf A, Schulz C, *et al.* Subcellular distribution of calcium-sensitive potassium channels (IK1) in migrating cells. *J Cell Physiol* 2006;**206**:86–94.
- 27 Ghanshani S, Wulff H, Miller MJ, *et al.* Up-regulation of the IKCa1 potassium channel during T-cell activation. Molecular mechanism and functional consequences. *J Biol Chem* 2000;**275**:37137–49.
- 28 Wulff H, Knaus HG, Pennington M, *et al.* K^{+} channel expression during B cell differentiation: implications for immunomodulation and autoimmunity. *J Immunol* 2004;**173**:776–86.
- 29 Grgic I, Eichler I, Heinau P, *et al.* Selective blockade of the intermediate-conductance Ca^{2+} -activated K^{+} channel suppresses proliferation of microvascular and macrovascular endothelial cells and angiogenesis in vivo. *Arterioscler Thromb Vasc Biol* 2005;**25**:704–9.
- 30 Aktas H, Fluckiger R, Acosta JA, *et al.* Depletion of intracellular Ca^{2+} stores, phosphorylation of eIF2 alpha, and sustained inhibition of translation initiation mediate the anticancer effects of clotrimazole. *Proc Natl Acad Sci USA* 1998;**95**:8280–5.
- 31 Zhang W, Ramamoorthy Y, Kilcarslan T, *et al.* Inhibition of cytochromes P450 by antifungal imidazole derivatives. *Drug Metab Dispos* 2002;**30**:314–8.
- 32 Penso J, Beitner R. Clotrimazole decreases glycolysis and the viability of lung carcinoma and colon adenocarcinoma cells. *Eur J Pharmacol* 2002;**45**:227–35.
- 33 Wojtulewski JA, Gow PJ, Walter J, *et al.* Clotrimazole in rheumatoid arthritis. *Ann Rheum Dis* 1980;**39**:469–72.
- 34 Kohler R, Wulff H, Eichler I, *et al.* Blockade of the intermediate-conductance calcium-activated potassium channel as a new therapeutic strategy for restenosis. *Circulation* 2003;**108**:1119–25.
- 35 Urbahns K, Horvath E, Stasch JP, *et al.* 4-Phenyl-4H-pyrans as IK(Ca) channel blockers. *Bioorg Med Chem Lett* 2005;**13**:2637–9.

LUNG ALERT

Lung cancer is more common but less often fatal in women

▲ International Early Lung Cancer Action Program Investigators. Women's susceptibility to tobacco carcinogens and survival after diagnosis of lung cancer. *JAMA* 2006;**296**:180–4

Lung cancer is on the rise, especially in women. Previous studies have suggested that women have a higher relative risk of developing lung cancer. This large follow up study from the International Early Lung Cancer Action Programme (ELCAP) investigators was designed to assess the risk of lung cancer related to smoking in women compared with men. Survival after diagnosis was also evaluated, using the data from the present study combined with those from the original ELCAP study.

An additional 6296 women and 8139 men (all asymptomatic) were screened with baseline CT scans. All patients with lung nodules underwent immediate biopsy or surgical resection. A panel of five expert lung pathologists reported the histological samples.

Lung cancer was diagnosed in 111/6296 women (1.7%) and 93/8139 men (1.1%). Combining the two series together, lung cancer was diagnosed in 156/7498 women (2.1%) and 113/9427 men (1.2%). This gave an odds ratio of cancer in women of 1.9 (95% CI 1.5 to 2.5) corrected for age and pack-year smoking history. It was also noted that survival was better in women, regardless of the stage of lung cancer and after matching for cell type and treatment. The hazard ratio for a fatal outcome in women was 0.48 (95% CI 0.25 to 0.89). This is an interesting result, but the authors could not find any specific reason for the difference in survival. It has been suggested that lung cancer in women may be less aggressive or relatively more curable than in men, and this needs to be evaluated further.

M S Anwar

Respiratory Registrar, Whipps Cross University Hospital, London, UK; Muhammad.anwar@whippsx.nhs.uk