- 26 **Griffiths TL**, Henson LC, Whipp BJ. Influence of inspired oxygen concentration on the dynamics of the exercise hyperpnoea in man. *J Physiol* 1986;**380**:387–403.
- 27 **Hughes RL**, Clode M, Edwards RHT, *et al.* Effect of inspired O₂ on cardiopulmonary and metabolic responses to exercise in man. *J Appl Physiol* 1968;**24**:336–47.
- 28 **Davies CTM**, Sargeant AJ. Physiological responses to one- and two-leg exercise breathing air and 45% oxygen. *J Appl Physiol* 1974;**36**:142–8.
- 29 **Casaburi R**, Stremel RW, Whipp BJ, *et al.* Alteration by hyperoxia of ventilatory dynamics during sinusoidal work. *J Appl Physiol* 1980;**48**:1083–91.
- 30 **Eklblom B**, Huot R, Stein EM, *et al.* Effect of changes in arterial oxygen content on circulation and physical performance. *J Appl Physiol* 1975;**39**:71–5.
- 31 **Bye PTP**, Esau SA, Walley KR, *et al.* Ventilatory muscles during exercise in air and oxygen in normal men. *J Appl Physiol* 1984;**56**:464–71.
- 32 **Welch HG**, Bonde-Petersen F, Graham T, *et al.* Effects of hyperoxia on leg blood flow and metabolism during exercise. *J Appl Physiol* 1977;**42**:385–90.
- 33 **Palange P**, Galassetti P, Mannix ET, *et al.* Oxygen effect on O₂ deficit and VO₂ kinetics during exercise in obstructive pulmonary disease. *J Appl Physiol* 1995;**78**:2228–34.
- 34 **O'Donnell DE**, Chau LKL, Bertley JC, *et al.* Qualitative aspects of exertional breathlessness in chronic airflow limitation: pathophysiologic mechanisms. *Am J Respir Crit Care Med* 1997;**155**:109–15.
- 35 **Hogan MC**, Richardson RS, Haseler LJ. Human muscle performance and PCr hydrolysis with varied inspired oxygen fractions: a ³¹P-MRS study. *J Appl Physiol* 1999;**99**:1367–73.
- 36 **Gosselin N**, Durand F, Poulain M, *et al.* Effect of acute hyperoxia during exercise on quadriceps electrical activity in active COPD patients. *Acta Physiol Scand* 2004;**181**:333–43.
- 37 **Gandevia SC**. Neural mechanisms underlying the sensation of breathlessness: kinesthetic parallels between respiratory and limb muscles. *Aust NZ J Med* 1988;**18**:83–91.

LUNG ALERT

Elastin fragments are pro-inflammatory in the progression of emphysema

▲ Houghton AM, Quintero PA, Perkins DL, *et al.* Elastin fragments drive disease progression in a murine model of emphysema. *J Clin Invest* 2006;**116**:753–9

Mice deficient in the macrophage elastase matrix metalloproteinase-12 (MMP-12) do not develop cigarette smoke induced emphysema or show high levels of macrophage accumulation in the lung on exposure to cigarette smoke. However, monocyte migration to the lung in MMP-12 deficient mice can occur in the presence of an appropriate stimulus. This study used an in vivo model to investigate that stimulus.

Bronchoalveolar lavage (BAL) fluid from wild-type smoke exposed mice (and not MMP-12 deficient controls) showed monocyte chemotactic activity. MMP-12 itself was not chemotactic. Fractionation of BAL fluid from wild-type mice found that elastin fragments were the only matrix protein fragments present in the chemotactic fraction of the fluid. The use of an elastin fragment antibody resulted in monocyte chemotaxis inhibition, even in the presence of cigarette smoke.

Porcine pancreatic elastase generates a model of emphysema that closely resembles human emphysema. Elastin fragment inhibition by elastin fragment antibody reduced lung macrophage accumulation and airspace disease in this model. The authors hypothesise that alveolar macrophage activation occurs in response to chronic cigarette smoke exposure, resulting in macrophage release of MMP-12. MMP-12 degrades lung extracellular matrix components and generates the production of elastin fragments that are chemotactic to monocytes and which drive disease progression.

Y Aggarwal

Senior House Officer, St George's Hospital, London, UK; dr_y_aggarwal@hotmail.com