- 20 Albain K, Scott CB, Rusch VR, *et al*. Phase III study of concurrent chemotherapy and full course radiotherapy (CT/RT) versus CT/RT induction followed by resection for stage IIIA (pN2) NSCLC. *Lung Cancer* 2003;**41**(Suppl 2):S4.
- 21 Izbicki J, Passlick B, Karg O, *et al*. Impact of radical systematic mediastinal lymphadenectomy on tumour staging in lung cancer. *Ann Thorac Surg* 1995;**59**:209–14.
- 22 Ginsberg R, Rubinstein LV, for the Lung Cancer Study Group. Randomised trial of lobectomy versus limited resection for T1N0 non-small cell lung cancer. *Ann Thorac Surg* 1995;**60**:615–23.
- 23 Rubinstein L, Ginsberg RJ. Lobectomy versus limited resection in T1N0 lung cancer. *Ann Thorac Surg* 1996;**62**:1249–50.
- 24 Sugi K, Kaneda Y, Esato K. Video-assisted thoracoscopic lobectomy achieves a satisfactory long term prognosis in patients with clinical stage 1A lung cancer. *World J Surg* 2000;**24**:27–31.
- 25 Morrison R, Deeley TJ, Cleland WP. The treatment of carcinoma of the bronchus: a clinical trial to compare surgery and supervoltage radiotherapy. *Lancet* 1963;**1**:683–4.
- 26 Shepherd F, Johnston MR, Payne D, *et al*. Randomised study of chemotherapy and surgery versus radiotherapy for stage IIIA non-small-cell lung cancer: a National Cancer Institute of Canada Clinical Trials Group Study. *Br J Cancer* 1998;**78**:683–5.
- 27 Warram J, on behalf of collaborating investigators. Preoperative irradiation of cancer of the lung: final report of a therapeutic trial (a collaborative study). *Cancer* 1975;**36**:914–25.
- 28 Warram J, on behalf of collaborating investigators. Preoperative irradiation of cancer of the lung: preliminary report of a therapeutic trial (a collaborative study). *Cancer* 1969;**23**:419–29.
- 29 Splinter T, van Schil PE, Kramer GWPM, *et al*. Randomised trial of surgery versus radiotherapy in patients with stage IIIA (N2) non-small cell lung cancer after response to induction chemotherapy (EORTC 08941). *Clin Lung Cancer* 2000;**2**:69–72.
- 30 Keller S. Complete mediastinal lymph node dissection – does it make a difference? *Lung Cancer* 2002;**36**:7–8.
- 31 Allen M, for the American College of Surgeons Oncology Group (Thoracic Organ Site Committee). Randomised trial of mediastinal lymph node sampling versus complete lymphadenectomy during pulmonary resection in patients with N0 or N1 (less than hilar) non-small cell lung cancer. Available at http://www.acosog.org/studies/synopses/Z0030_Synopsis.pdf (accessed October 2004).
- 32 Naruke T, Suemasu K, Ishikawa S. Surgical treatment for lung cancer with metastasis to mediastinal lymph nodes. *J Thorac Cardiovasc Surg* 1976;**71**:279–85.
- 33 Martini N, Flehinger BJ. The role of surgery in N2 lung cancer. *Surg Clin North Am* 1987;**67**:1037.
- 34 Feinstein A, Sosin DM, Wells CK. The Will Rogers phenomenon. Stage migration and new diagnostic techniques as a source of misleading statistics for survival in cancer. *N Engl J Med* 1985;**312**:1604–8.
- 35 Bach P, Cramer LD, Schrag D, *et al*. The influence of hospital volume on survival after resection for lung cancer. *N Engl J Med* 2001;**345**:181–8.
- 36 Kiser A, Detterbeck FC. General aspects of surgical treatment. In: Detterbeck F, Rivera MP, Socinski MA, Rosenman JG, eds. *Diagnosis and treatment of lung cancer: an evidence based guide*. Philadelphia: WB Saunders, 2001:133–47.

LUNG ALERT

High prevalence of PE in patients with unexplained exacerbation of COPD

▲ Tillie-Leblond I, Marquette CH, Perez T, *et al*. Pulmonary embolism in patients with unexplained exacerbation of chronic obstructive pulmonary disease: prevalence and risk factors. *Ann Intern Med* 2006;**144**:390–6

This single centre study enrolled 197 patients with COPD requiring admission to hospital for an unexplained acute exacerbation. Patients with any evidence of lower respiratory tract infection, pneumothorax, and those requiring invasive ventilation were excluded. Patients with clinical and radiological findings deemed out of keeping with the degree of hypoxaemia were included. All patients were investigated with computed tomogram pulmonary angiography (CTPA) and venous lower limb ultrasonography.

Evidence of pulmonary embolus (PE) was found in 49 of the 197 patients (25%, 95% CI 19 to 32). This is consistent with previously published data. The diagnosis of PE was determined by positive CTPA and deep vein thrombosis (DVT) on ultrasound in 19 patients; 24 patients had a positive CTPA alone and six with a negative CTPA had DVT seen on ultrasound. The patients were retrospectively categorised into low, intermediate, and high probability according to the Geneva score. Eleven of the 119 patients (9%) in the low probability group had PE, 35 of 75 patients (46%) in the intermediate probability category had PE, and all three in the high probability group had PE.

The diagnosis of PE could not therefore be excluded on the basis of a low probability Geneva score. The only reliable risk factors identified in this study group were previous thromboembolism, malignancy, and a decreased arterial carbon dioxide tension compared with baseline.

A Patel

Senior Clinical Fellow, Royal Free Hospital, London, UK; anantpatel478@yahoo.co.uk