

- 21 **Troosters T**, Vilaro J, Rabinovich R, *et al*. Physiological responses to the 6-min walk test in patients with chronic obstructive pulmonary disease. *Eur Respir J* 2002;**20**:564–9.
- 22 **Casas A**, Vilaro J, Rabinovich R, *et al*. Encouraged 6-min walking test indicates maximum sustainable exercise in COPD patients. *Chest* 2005;**128**:55–61.
- 23 **Connolly CK**, Chan NS. Salbutamol and ipratropium in partially reversible airway obstruction. *Br J Dis Chest* 1987;**81**:55–61.
- 24 **Hay JG**, Stone P, Carter J, *et al*. Bronchodilator reversibility, exercise performance and breathlessness in stable chronic obstructive pulmonary disease. *Eur Respir J* 1992;**5**:659–64.
- 25 **Spence DP**, Hay JG, Carter J, *et al*. Oxygen desaturation and breathlessness during corridor walking in chronic obstructive pulmonary disease: effect of oxitropium bromide. *Thorax* 1993;**48**:1145–50.
- 26 **Eiser N**, Willsher D, Dore CJ. Reliability, repeatability and sensitivity to change of externally and self-paced walking tests in COPD patients. *Respir Med* 2003;**97**:407–14.
- 27 **Redelmeier DA**, Bayoumi AM, Goldstein RS, *et al*. Interpreting small differences in functional status: the six-minute walk test in chronic lung disease patients. *Am J Respir Crit Care Med* 1997;**155**:1278–82.
- 28 **McDowell I**, Newell C. *Measuring health: a guide to rating scales and questionnaires*, 2nd ed. New York: Oxford University Press, 1996.
- 29 **Man WD**, Soliman MG, Gearing J, *et al*. Symptoms and quadriceps fatigability after walking and cycling in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2003;**168**:562–7.
- 30 **O'Donnell DE**, Lam M, Webb KA. Spirometric correlates of improvement in exercise performance after anticholinergic therapy in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1999;**160**:542–9.
- 31 **Sciurba F**, Criner GJ, Lee SM, *et al*. Six-minute walk distance in chronic obstructive pulmonary disease: reproducibility and effect of walking course layout and length. *Am J Respir Crit Care Med* 2003;**167**:1522–7.
- 32 **Lacasse Y**, Wong E, Guyatt GH, *et al*. Meta-analysis of respiratory rehabilitation in chronic obstructive pulmonary disease. *Lancet* 1996;**348**:1115–9.
- 33 **Eaton T**, Young P, Nicol K, *et al*. The endurance shuttle walking test: a responsive measure in pulmonary rehabilitation for COPD patients. *Chronic Respir Dis* 2006;**3**:3–9.
- 34 **Laude EA**, Duffy NC, Baveystock C, *et al*. The effect of helium and oxygen on exercise performance in chronic obstructive pulmonary disease: a randomized crossover trial. *Am J Respir Crit Care Med* 2006;**173**:865–70.

LUNG ALERT

CT screening helps to detect resectable lung cancers and may improve survival

▲ Henschke CI, Yankelevitz DF, Libby DM, *et al*. Survival of patients with stage I lung cancer detected on ct screening the international early lung cancer action programme investigators. *N Engl J Med* 2006;**355**:1763–71.

This international uncontrolled prospective study screened 31 567 asymptomatic individuals using low dose CT scan between 1993 and 2005. Positive tests were defined on baseline screening as the identification of a least one solid or partly solid non-calcified pulmonary nodule >5 mm in diameter, at least one non-solid non-calcified pulmonary nodule >8 mm in diameter or a solid endobronchial node; and on annual screening as any new non-calcified nodule. Patients with suspicious lesions underwent investigation with positron emission tomography (PET), biopsy or repeat CT scan. The study identified 484 participants with lung cancer, 412 (85%) of whom had clinical stage I disease with an estimated 10 year survival rate of 88%. Survival rates increased to 92% in the 302 individuals with stage I cancer who underwent surgical resection within 1 month of diagnosis.

The study is limited by lack of a control group and disease specific mortality as its endpoint, resulting in lead time, length time and over diagnosis bias. The majority of lung cancers detected were found on baseline screening (405) compared with annual CT scan (75) and survival rates are statistically projected rather than true values. CT scan as a primary screening tool is non-invasive with a relative low morbidity, however it yields no information regarding the biological behaviour of the nodule and confirmatory diagnosis requires further invasive procedures. CT scan-based screening programmes may prove difficult to implement within the current National Health Service system in the UK, although this American study demonstrates that it can detect curable cancers.

Helen Liddicoat

SHO in Respiratory Medicine, Homerton University Hospital, London, UK; drhliddicoat@hotmail.com