Ventilation heterogeneity is a major determinant of airway hyperresponsiveness in asthma, independent of airway inflammation

Sue R Downie, Cheryl M Salome, Sylvia Verbanck, Bruce Thompson, Norbert Berend, Gregory G King

Thorax 2007;62:684–689. doi: 10.1136/thx.2006.069682

See end of article for authors' affiliations

Correspondence to: Ms Sue Downie, Woolcock Institute of Medical Research, P O Box M77, Camperdown, NSW 2050, Australia; sued@woolcock.org.au

Received 3 August 2006
Accepted 11 January 2007
Published Online First
20 February 2007

Airway hyperresponsiveness (AHR) is a common feature of asthma that is defined as the ability of airways to narrow too easily and by too much in response to provoking stimuli. In subjects with asthma with severe AHR, excessive airway narrowing represents the potential for severe and life-threatening asthma attacks.¹ AHR is a clinically useful tool for treatment monitoring,² and is associated with impaired development of lung function in childhood.³ Although AHR has been shown to be associated with airway inflammation⁴ and ventilation heterogeneity,⁵ the mechanisms that cause AHR are poorly understood.

Although airway inflammation is regarded as the underlying cause of AHR,⁶ it is not known exactly how inflammation may cause AHR. Significant associations between AHR and markers of eosinophilic inflammation in induced sputum⁷ and the level of nitric oxide in exhaled breath⁴ have been observed, but other studies have failed to show any association between airway inflammation and AHR.^{8–9} Evidence suggests that other non-eosinophilic pathways such as neutrophilic inflammation are important contributors to AHR and asthma.¹⁰ Exhaled nitric oxide is an indirect marker of eosinophilic airway inflammation that is easy to perform, is reduced with anti-inflammatory inhaled corticosteroid treatment (ICS),² and is more closely associated with AHR and active asthma than other markers of airway inflammation such as sputum eosinophils⁴ and serum markers.¹¹ For these reasons, exhaled nitric oxide may be a suitable marker of eosinophilic inflammation to explore the relationship between AHR, airway inflammation and ventilation heterogeneity, and to determine the effect of ICS treatment on this relationship.

Background: Airway hyperresponsiveness is the ability of airways to narrow excessively in response to inhaled stimuli and is a key feature of asthma. Airway inflammation and ventilation heterogeneity have been separately shown to be associated with airway hyperresponsiveness. A study was undertaken to establish whether ventilation heterogeneity is associated with airway hyperresponsiveness independently of airway inflammation in subjects with asthma and to determine the effect of inhaled corticosteroids on this relationship.

Methods: Airway inflammation was measured in 40 subjects with asthma by exhaled nitric oxide, ventilation heterogeneity by multiple breath nitrogen washout and airway hyperresponsiveness by methacholine challenge. In 18 of these subjects with uncontrolled symptoms, measurements were repeated after 3 months of treatment with inhaled beclomethasone dipropionate.

Results: At baseline, airway hyperresponsiveness was independently predicted by airway inflammation (partial $r^2=0.20$, $p<0.001$) and ventilation heterogeneity (partial $r^2=0.39$, $p<0.001$). Inhaled corticosteroid treatment decreased airway inflammation ($p=0.002$), ventilation heterogeneity ($p=0.009$) and airway hyperresponsiveness ($p<0.001$). After treatment, ventilation heterogeneity was the sole predictor of airway hyperresponsiveness ($r^2=0.64$, $p<0.001$).

Conclusions: Baseline ventilation heterogeneity is a strong predictor of airway hyperresponsiveness, independent of airway inflammation in subjects with asthma. Its persistent relationship with airway hyperresponsiveness following anti-inflammatory treatment suggests that it is an important independent determinant of airway hyperresponsiveness. Normalisation of ventilation heterogeneity is therefore a potential goal of treatment that may lead to improved long-term outcomes.

Ventilation heterogeneity is another potential cause of AHR.⁵ Supportive evidence for a causal association arises from computer simulations whereby a more uneven distribution of ventilation and airway narrowing throughout the lung results in a greater overall increase in airway resistance following bronchoconstriction (ie, AHR).^{12–14} Ventilation heterogeneity can be assessed non-invasively by multiple breath nitrogen washout (MBNW).¹⁵ The greater the degree of ventilation heterogeneity within the lung during continuous tidal breathing of 100% oxygen, the greater the nitrogen phase III slope in each subsequent MBNW expiration. More specifically, the phase III slope is determined by convective gas transport, predominantly in the conductive airways, and by diffusive transport in the more peripheral acinar airways. From multiple breath washout tests using inert gases with different diffusivities,¹⁶ SF₆ and helium, it has been possible to partition the convection from the diffusion-dependent portion of the phase III slope, and derive corresponding indices of conductive airway heterogeneity (S_{cond}) and acinar heterogeneity (S_{acin}). The value of these indices increases as ventilation heterogeneity in the corresponding lung zone increases. The mathematical

Abbreviations: AHR, airway hyperresponsiveness; CEV, cumulative expired volume; CFC-BDP, chlorofluorocarbon beclomethasone dipropionate; DRR, dose response ratio; FEV₁, forced expiratory volume in 1 s; FRC, functional residual capacity; FE_{NO}, fraction of nitric oxide in exhaled breath; FVC, forced vital capacity; HFA-BDP, hydrofluoroalkane beclomethasone dipropionate; ICS, inhaled corticosteroid; LCI, lung clearance index; MBNW, multiple breath nitrogen washout; ROC, receiver operator characteristics; S_{acin} , ventilation heterogeneity in acinar lung zone; S_{cond} , ventilation heterogeneity in conducting airways

derivation of these indices is explained in the online supplement available at <http://thorax.bmj.com/supplemental>. Using this MBNW analysis, ventilation heterogeneity in conductive and acinar lung zones has been shown to be abnormal in asthma¹⁷ and to be more sensitive than spirometry for detecting early changes to the peripheral airways associated with cigarette smoking.¹⁵

We hypothesised that there would be an association between ventilation heterogeneity and AHR and have tested whether this association is independent of airway inflammation in a cross-sectional study of subjects with asthma of a wide range of severity. In a subgroup of these subjects with clinically uncontrolled asthma, we further tested the relationship between airway inflammation, ventilation heterogeneity and AHR following 3 months of treatment with ICS.

METHODS

Study design

Subjects in whom asthma had been diagnosed by a respiratory physician according to the NIH guidelines¹⁸ were enrolled in the study. At the baseline visit all subjects underwent the following tests in the order shown: (1) atopic status was determined by skin prick testing with mean weal diameters ≥ 4 mm regarded as positive; (2) Juniper asthma control questionnaire; (3) the fraction of nitric oxide in exhaled breath (FE_{NO}) was used as a marker of airway inflammation; (4) MBNW was performed to measure ventilation heterogeneity; and (5) methacholine challenge was undertaken to determine AHR. A subgroup of patients with at least mild persistent asthma, as determined by GINA guidelines,¹⁹ were given ICS for 3 months with either chlorofluorocarbon beclomethasone dipropionate (CFC-BDP) 750 μ g twice daily via Autohaler or hydrofluoroalkane beclomethasone dipropionate (HFA-BDP) 400 μ g twice daily via Autohaler (clinically equivalent doses). Since our aim was to investigate overall treatment effects, all treatment data were pooled for analyses.

Written informed consent was obtained from all subjects and the study was approved by the human ethics review committee of the South-Western Area Health Service. This study is registered on the Australian Clinical Trials Registry (#ACTRN012605000317695).

Subjects

Subjects with asthma were recruited by advertising throughout the University of Sydney. Inclusion criteria for all subjects with asthma were: (1) no smoking within the last 6 months and <10 pack-years smoking history; (2) no current lung disease other than asthma; (3) no oral prednisone use in the last 4 weeks; and (4) no respiratory tract infection in the last 4 weeks. Additional inclusion criteria for entry into the 3 month treatment period were: (1) asthma symptoms on at least three occasions a week over the previous month; (2) AHR defined as a provoking dose of methacholine causing a 20% fall in FEV_1 of ≤ 6.1 μ mol; and (3) taking no more than 800 μ g/day CFC-BDP equivalent of ICS.

Two cohorts of healthy non-asthmatic subjects were recruited from our research staff and the University of Sydney. The first cohort was recruited for the sensitivity and specificity analysis of a parameter from the MBNW test in detecting AHR, and the second cohort was recruited for testing the repeatability of the MBNW indices. Healthy non-asthmatic subjects were eligible to participate if they had (1) no history of or medication use relating to chronic respiratory disease; (2) no smoking within the last 6 months and <10 pack-years smoking history; and (3) no respiratory tract infection in the last 4 weeks.

Symptoms

All the subjects with asthma completed the Juniper asthma control questionnaire²⁰ at the baseline visit and after 3 months of treatment (in the treatment subgroup) to assess asthma symptoms within the previous week.

Exhaled nitric oxide

FE_{NO} was measured using an offline technique²¹ according to American Thoracic Society guidelines. The subject exhaled over 5–15 s into a nitric oxide impermeable polyethylene bag (Scholle Industries Pty Ltd, Elizabeth West, Australia) at a constant flow rate of 0.2 l/s, monitored by rotameter (Dwyer Flowmeter Model VFSS-25, AMBIT Instruments Pty Ltd, Parramatta, Australia). The exhaled gas was analysed using a chemiluminescence analyser (Thermo Environmental Instruments Model 42C, Franklin, Massachusetts, USA). The upper limit of normal for FE_{NO} has been established as 13 ppb.²²

Multiple breath nitrogen washout (MBNW)

The MBNW was performed using a closed circuit bag-in-box breathing system to deliver 100% oxygen during inspiration with separate capture of exhaled breath. The nitrogen concentration was measured at the mouth with a Model 721 KaeTech Nitrogen Analyser (KaeTech Instruments Inc, Green Bay, Wisconsin, USA), while flow and volume were recorded from the box by a pneumotachograph. Subjects breathed 100% oxygen at a tidal volume of 1–1.3 litres until the mean expired nitrogen concentration fell to $<2\%$. Three tests were performed with successive tests starting after alveolar nitrogen had returned to baseline. Ventilation heterogeneity indices in the conductive (S_{cond}) and acinar (S_{acin}) lung zones were derived as previously described¹⁷ (see calculations in online supplement available at <http://thorax.bmj.com/supplemental>). The upper limit of normal for S_{cond} was 0.037/l and for S_{acin} was 0.130/l, derived from the mean + 1.96 $\times$ SD of the value obtained previously in normal subjects.²³ The lung clearance index (LCI) was calculated by dividing the cumulative expired volume (CEV) during the washout by the patient's functional residual capacity (FRC) determined from the washout. Details of a repeatability study for S_{cond} , S_{acin} and LCI obtained from an independent group of patients with asthma and healthy non-asthmatic patients are given in the online supplement available at <http://thorax.bmj.com/supplemental>.

Spirometry and bronchial challenge

Spirometry was performed using a Sensormedics Vmax spirometer (Sensormedics Corporation, Yorba Linda, California, USA) and methacholine challenge tests were performed using the rapid method²⁴ via hand-held DeVilbiss No 45 nebulisers in which doubling doses ranging from 0.05 μ mol to 6.1 μ mol were administered until the final dose was reached or the forced expiratory volume in 1 s (FEV_1) fell by $\geq 20\%$. Short acting β -agonists were withheld for 6 h and long-acting β -agonists for 24 h before testing. Response to challenge was measured by the dose response ratio (DRR), which is a continuous variable based on the two-point slope.²⁵ The DRR is calculated from the final step in the challenge test as % fall in FEV_1 /methacholine dose (μ mol), and a constant of 3 is added to allow for log transformation of zero or negative values. Greater values of DRR indicate greater AHR, and a DRR value of $>6.28\%$ fall in FEV_1 / μ mol methacholine + 3 indicates the presence of AHR. This allows inclusion of data from subjects who had a fall in FEV_1 of $<20\%$ either at baseline or following treatment.

Table 1 Characteristics of study subjects

	Entire asthma group (n = 40)	ICS treatment asthma subgroup (n = 18)		Absolute difference† (95% CI)	p Value†
		Before treatment	After treatment		
M/F	20/20	9/9	–	–	–
Mean (range) age (years)	32.1 (18–66)	33.9 (18–63)	–	–	–
Atopic (Y/N)	40/0	–	–	–	–
FE _{NO} (ppb)*	15.4 (12.2 to 19.4)	14.1	8.0	6.5 (1.5 to 11.6)	0.002
S _{cond} (/l)	0.055 (0.03)	0.067	0.052	0.02 (0.01 to 0.03)	0.009
S _{acin} (/l)	0.146 (0.02)	0.159	0.142	0.02 (0.00 to 0.03)	0.035
LCI (CEV/FRC)	8.16 (1.1)	8.54	7.78	0.8 (0.4 to 1.1)	0.001
DRR (% fall FEV ₁ /μmol methacholine+3)*	18.9 (12.4 to 28.8)	22.0	10.1	1.1 (0.6 to 1.6)‡	<0.001
FEV ₁ (% predicted)	82.4 (11.7)	80.1	85.9	5.8 (3.2 to 8.4)	<0.001
FEV ₁ /FVC ratio	0.76 (0.08)	0.73	0.75	0.02 (0.00 to 0.04)	0.04
β agonist use (times/day)	1.4 (2.4)	1.95	0.54	0.8 (–0.0 to 1.6)	0.07
BDP equivalent dose (μg/day)	320 (449.6)	270	1500	–	–
Juniper symptom score	1.28 (0.84)	1.48	0.81	0.7 (0.4 to 0.9)	<0.001

Values are mean (SD) unless otherwise stated.
*Geometric mean and 95% CI
†Comparison between values before and after treatment.
‡Doubling dose difference in DRR.
BDP, beclomethasone dipropionate; CEV, cumulative expired volume; DRR, dose response ratio; FEV₁, forced expiratory volume in 1 s; FE_{NO}, fraction of nitric oxide in exhaled breath; FRC, functional residual capacity; FVC, forced vital capacity; ICS, inhaled corticosteroid; LCI, lung clearance index; MBNW, multiple breath nitrogen washout; S_{cond}, ventilation heterogeneity in the conducting airways; S_{acin}, ventilation heterogeneity in acinar lung zone; ppb, parts per billion.

Statistical analysis

Data were analysed using the Analyse-it for Microsoft Excel software (Analyse-It Software Ltd, Leeds, UK). DRR and FE_{NO} were log-normally distributed and were log₁₀ transformed for all analyses. Change in DRR after treatment was expressed as change in doubling dose of methacholine (change in log DRR/0.3). The relationship between AHR (expressed as DRR) and potential predictive factors was examined using univariate analysis (with Pearson’s correlation coefficient) and multiple linear regression analyses. The sensitivity and specificity of S_{cond} in predicting AHR was assessed by receiver operator characteristics (ROC) analysis. The paired Student’s *t* test was used for parametric data and the Wilcoxon signed rank test for non-parametric data to compare outcomes after treatment. Data are presented as means with 95% confidence intervals (CI) unless otherwise specified.

RESULTS

Subjects

Forty subjects with asthma participated in the baseline study; 28 had suboptimally controlled asthma and were eligible for the 3 month treatment study and 24 of these agreed to participate. Four subjects from the treatment subgroup withdrew from the study during treatment and the results of two subjects were excluded from the analysis owing to an upper respiratory tract infection at their post-treatment visit. Eighteen subjects were therefore included in the treatment subgroup analysis.

Baseline results of entire asthma group

The characteristics of all 40 patients with asthma at baseline are shown in table 1. AHR was present in 31 of the 40 subjects, and log DRR correlated positively with airway inflammation (log FE_{NO}), S_{cond} (fig 1A) and LCI and negatively with % predicted FEV₁ and FEV₁/FVC ratio (table 2). Log DRR was not correlated with S_{acin} (p = 0.67). Log FE_{NO} was not correlated with S_{cond} (r = 0.29, p = 0.07), S_{acin} (r = 0.20, p = 0.22) or LCI (r = 0.14, p = 0.40) at baseline.

In a multiple linear regression model which included the significant factors in table 2, log DRR was significantly predicted by both log FE_{NO} (partial r² = 0.20, p<0.001, β coefficient 0.9, 95% CI 0.5 to 1.3) and S_{cond} (partial r² = 0.39, p<0.001, β coefficient 8.5, 95% CI 4.8 to 12.2). LCI, % predicted

FEV₁ and FEV₁/FVC ratio were not significant predictors of log DRR (p = 0.90, p = 0.25 and p = 0.97, respectively) when included separately or together in this model.

Inhaled corticosteroid treatment subgroup

The characteristics of the 18 subjects with asthma in the treatment subgroup are shown in table 1. Following treatment there were significant improvements in DRR, FE_{NO}, S_{cond}, S_{acin}, LCI, FEV₁ (% predicted) and the Juniper symptom score (table 1), but AHR was still present in 10 of the 18 subjects. The univariate correlations at baseline in this subgroup were similar to those in the group as a whole (table 2, fig 1B). However, LCI, % predicted FEV₁ and the FEV₁/FVC ratio were not correlated with log DRR either before or after treatment in the treatment subgroup (table 2).

In a multiple linear regression model, log DRR in the treatment subgroup was predicted solely by S_{cond} both before (r² = 0.38, p = 0.004, β coefficient 8.3, 95% CI 3.1 to 13.6) and after treatment (r² = 0.64, p<0.001, β coefficient 10.9, 95% CI 6.8 to 15). Log FE_{NO}, LCI, % predicted FEV₁ and the FEV₁/FVC ratio were not significant predictors of log DRR either before (p = 0.18, p = 0.90, p = 0.24, p = 0.99, respectively) or after treatment (p = 0.38, p = 0.30, p = 0.25, p = 0.42, respectively) in the multivariate regression model.

The magnitude of improvement in AHR correlated with both the reductions in FE_{NO} and S_{cond} (F = 3.94, p = 0.04 for the multivariate model), but there was no correlation between the reductions in S_{cond} and FE_{NO}0.2 following treatment (r = 0.07, p = 0.77). There were no differences in the magnitude of improvements in FE_{NO} (p = 0.26), S_{cond} (p = 0.81) or DRR (p = 0.90) between subjects treated with HFA-BDP or CFC-BDP (corresponding post-treatment data are represented by squares and triangles in fig 1B).

S_{cond} as a predictor of AHR

Based on the correlation between AHR and S_{cond}, we combined the patients with asthma from this study (n = 40) with a cohort of 17 healthy non-asthmatic subjects (their characteristics are summarised in table E1 in the supplementary data available online at <http://thorax.bmj.com/supplemental>) to test whether S_{cond} was a significant predictor of the presence of AHR using ROC analysis (fig 2). The area under the ROC curve was 0.88

Figure 1 (A) Conductive airway ventilation heterogeneity (S_{cond}) correlates with airway hyperresponsiveness (AHR) (log DRR) at baseline ($r=0.63$, $p<0.001$). Closed circles represent those who participated in the inhaled corticosteroid treatment study and open circles represent those who did not. (B) S_{cond} continues to correlate with AHR following treatment ($r=0.82$, $p<0.001$). Triangles represent subjects treated with chlorofluorocarbon beclomethasone dipropionate (CFC-BDP) and squares represent subjects treated with hydrofluoroalkane beclomethasone dipropionate (HFA-BDP).

($p<0.0001$), and the cut-off value of $S_{cond} = 0.037/l$ corresponding to the upper limit of normal range²⁶ gave a good combination of sensitivity and specificity (71.9% and 90.0%, respectively) for detecting the presence of AHR. Using the same cut-off value (0.037/l), S_{cond} remained a significant predictor of

Figure 2 Sensitivity and specificity of ventilation heterogeneity in conducting airways (S_{cond}) in predicting the presence or absence of airway hyperresponsiveness in a combination of 40 patients with asthma and 17 healthy non-asthmatic subjects at baseline.

the presence of AHR after treatment in our 18 subjects from the subgroup (area under the ROC curve = 0.96, $p<0.0001$) with a high sensitivity and specificity (90.0% and 87.5%, respectively, see fig E2 in the supplementary data available online at <http://thorax.bmj.com/supplemental>).

Repeatability of S_{cond} , S_{acin} and LCI

The characteristics of the cohort of 10 subjects with asthma and 11 healthy non-asthmatic subjects who participated in the repeatability study of the MBNW indices are summarised in table E2 in the supplementary data (available online at <http://thorax.bmj.com/supplemental>). In subjects with asthma, S_{cond} , S_{acin} and LCI had a good repeatability with an intraclass correlation coefficient of 0.84 (95% limits of agreement ± 0.026) for S_{cond} , 0.95 (95% limits of agreement ± 0.027) for S_{acin} , and 0.91 (95% limits of agreement ± 0.735) for LCI. The repeatability results for the healthy non-asthmatic subjects are shown in table E3 in the supplementary data (available online at <http://thorax.bmj.com/supplemental>).

Table 2 Pearson correlation coefficients (r) of baseline variables with airway hyperresponsiveness (dose response ratio, DRR) for the asthmatic group as a whole and for the treatment subgroup before and after treatment

	Baseline asthma group (n = 40)		Baseline treatment subgroup (n = 18)		Post-treatment treatment subgroup (n = 18)	
	r	p Value	r	p Value	r	p Value
Airway inflammation (FE_{NO})	0.62	<0.001	0.51	0.03	0.61	0.008
Ventilation heterogeneity (S_{cond})	0.63	<0.001	0.64	0.004	0.82	<0.001
Ventilation heterogeneity (S_{acin})	0.07	0.67	0.04	0.86	-0.02	0.93
Ventilation heterogeneity (LCI)	0.38	0.02	0.31	0.21	0.47	0.05
FEV_1 % predicted	-0.44	0.005	-0.40	0.10	-0.37	0.13
FEV_1/FVC ratio	-0.47	0.002	-0.34	0.17	-0.21	0.41

FEV_1 , forced expiratory volume in 1 s; FE_{NO} , fraction of nitric oxide in exhaled breath; FVC, forced vital capacity; LCI, lung clearance index; S_{cond} , ventilation heterogeneity in conducting airways; S_{acin} , ventilation heterogeneity in acinar lung zone

DISCUSSION

In this study we have shown that baseline ventilation heterogeneity correlates strongly with the severity of AHR in subjects with asthma with a range of disease severity. In particular, S_{cond} predicted AHR to methacholine independently of airway inflammation and airway calibre. Importantly, the strong correlation between S_{cond} and AHR persisted after airway inflammation was reduced with ICS treatment. Following treatment, S_{cond} was, in fact, the sole independent predictor of AHR. The improvement in S_{cond} with ICS treatment suggests that it is determined, at least in part, by steroid-responsive inflammatory processes. However, the residual ventilation heterogeneity which persisted after ICS treatment suggests that non-steroid responsive or non-inflammatory factors also make an important contribution to ventilation heterogeneity. The conductive airways, where the residual ventilation heterogeneity responsible for AHR after anti-inflammatory treatment originates, clearly constitute an important therapeutic target. These findings are novel and have clinical implications in patients with asthma.

Ventilation heterogeneity in the lungs is a well recognised feature of asthma,¹² but its clinical significance—apart from the deleterious effects on gas exchange²⁷—has been uncertain. In our study, the main contribution to ventilation heterogeneity that was associated with AHR originated in the conductive airways. Gustafsson *et al*⁵ reported a correlation between airway responsiveness to cold air challenge and ventilation heterogeneity measured by single breath washout using helium and SF₆, which indicated an involvement of peripheral airways but did not assess the contribution of airway inflammation. It is impossible independently to determine ventilation heterogeneity of conductive airways from the single breath washout, making comparison with the present study difficult. However, despite methodological differences in the determination of ventilation heterogeneity and AHR, the observed correlations between ventilation heterogeneity and AHR in both studies suggest a robust association. In the present study, AHR was not correlated with baseline acinar (S_{acin}) ventilation heterogeneity, signalling that the important lung region with regard to the relationship between ventilation heterogeneity and AHR is specifically in the conducting airways. The weak correlation between the LCI and AHR indicates that this measure of specific ventilation distribution between relatively large lung units, which also partly contributes to S_{cond} , is neither sensitive nor specific as a measure of the conductive airway heterogeneity underlying AHR.

The significant reduction in ventilation heterogeneity in our subjects following anti-inflammatory ICS treatment suggests that ventilation heterogeneity is partly due to steroid-responsive airway inflammation. After treatment, residual ventilation heterogeneity persisted in the presence of normalised mean exhaled nitric oxide levels, suggesting that other inflammatory processes not related to exhaled nitric oxide, or non-inflammatory processes, could also contribute to ventilation heterogeneity. Exhaled nitric oxide correlates with sputum eosinophils,²⁸ but it does not reflect the full spectrum of inflammatory and non-inflammatory processes associated with asthma. This may explain the lack of association between improvement in exhaled nitric oxide and ventilation heterogeneity during treatment. Ventilation heterogeneity could also continue to improve with further ICS treatment, as has been shown to occur with AHR which can continue to improve up to and after 12 months of treatment.²⁹ It would be valuable in future studies also to assess any additional benefits of systemic treatments on ventilation heterogeneity.

The residual ventilation heterogeneity present at the end of treatment may also be caused by non-inflammatory structural

changes such as airway remodelling or increased smooth muscle tone. Structural changes induced by airway remodelling are likely to be heterogeneously distributed among parallel pathways of the bronchial tree. These structural changes manifest as altered composition and organisation of the soft tissues of the airway walls and are believed to be a result of chronic or repeated episodes of acute inflammation.³⁰ The changes include thickening of the airway wall, smooth muscle hyperplasia and hypertrophy, subepithelial fibrosis and mucous metaplasia. It is possible that many, if not all, of these abnormalities may contribute to ventilation heterogeneity by exaggerating the inherent heterogeneity of the airway tree and by being unevenly distributed themselves.

Increased baseline ventilation heterogeneity may cause a heterogeneous distribution of methacholine aerosol resulting in high concentrations of methacholine delivered to a small surface area of the airways. In a previous study, central-to-peripheral airway differences in the pattern of methacholine deposition in subjects with asthma did not alter AHR.³¹ However, a recent imaging study in sheep³² showed that the regions of lung with the highest baseline ventilation were the same regions which became constricted after methacholine, suggesting a link between parallel ventilation heterogeneity and methacholine deposition. This could partly explain the association between S_{cond} (representative of parallel ventilation heterogeneity) and AHR. The exact mechanism by which ventilation heterogeneity may cause AHR remains unclear. However, a recent modelling study¹² showed that only minor alterations in structure, function or smooth muscle activity in a virtually uniform airway tree are required to trigger extreme ventilation heterogeneity during uniformly applied bronchoconstriction.

One of the strengths of our study was the wide range of asthma severity in the group of subjects tested at baseline. Indices of ventilation heterogeneity from the MBNW were robust, repeatable and comparable with values obtained in other subjects with asthma.^{17–33} The strong predictive capacity of S_{cond} for the presence of AHR, as inferred from our ROC analysis (fig 2), is reinforced because the upper limit of the normal range for S_{cond} (0.037/l) corresponded to a high combination of sensitivity and specificity for detecting the presence of AHR in the present study. This raises the possibility that S_{cond} could at least partly be used to predict AHR as part of the clinical management of asthma. MBNW tests can be performed in patients whose lung function is too low to permit bronchial challenge and has been successfully administered in non-sedated children with a number of airway diseases.³⁴

In summary, we have shown for the first time that ventilation heterogeneity in the conducting airways is a significant predictor of AHR in subjects with asthma, independent of airway inflammation as measured by exhaled nitric oxide, both before and after ICS treatment. Inflammatory and non-inflammatory processes which could alter the structural features within the airway tree may contribute to ventilation heterogeneity. Based on the strength of the correlations in this study and on the results of computational modelling,^{12–14} it is possible that ventilation heterogeneity is an important contributor to AHR in asthma. The index S_{cond} derived from the MBNW is a measure of conducting airway function that is likely to be useful in the clinical management of asthma and for the assessment of new treatment strategies and drugs in this disease. These findings suggest that greater understanding of the determinants of ventilation heterogeneity may lead to more efficacious asthma treatment and can provide an additional tool for monitoring treatment, resulting in improved long-term outcomes for patients with asthma.

ACKNOWLEDGEMENTS

The authors thank the volunteers who participated in the study; Nathan Brown, Chantale Diba, Phillip Munoz, Dr Ricardo Starling-Schwanz, Dr Stephen Vincent and Dr Ciça Santos who assisted with patient visits; Gunnar Unger and Tom Li who gave engineering support for the multiple breath nitrogen washout; and Wei Xuan for his advice on statistical analysis.

Additional data are available in the online supplement at <http://thorax.bmj.com/supplemental>.

Authors' affiliations

Sue R Downie, Cheryl M Salome, Norbert Berend, Gregory G King, Woolcock Institute of Medical Research, Camperdown, Sydney, Australia
Sylvia Verbanck, Respiratory Division, Academic Hospital, Vrije Universiteit Brussels, 1090 Brussels, Belgium
Bruce Thompson, Department of Allergy, Immunology, and Respiratory Medicine, The Alfred Hospital and Monash University, Melbourne, Australia
Norbert Berend, Gregory G King, Department of Respiratory Medicine, Royal North Shore Hospital, Sydney, Australia

This project was funded by the Cooperative Research Centre for Asthma.

Competing interests: The authors have the following potential conflicts of interest to declare: Cheryl M Salome states that her research group has received support from Boehringer Ingelheim, GlaxoSmithKline and AstraZeneca but these do not pose any conflict of interest for this paper. Norbert Berend serves on advisory boards for GlaxoSmithKline, Boehringer Ingelheim and AstraZeneca, and The Woolcock Institute of Medical Research of which he is the Director receives research grants from GlaxoSmithKline. Gregory King has received travel sponsorship from GlaxoSmithKline to attend the ATS ASM 2003 (approximately \$A9000) and a GlaxoSmithKline meeting in 2003 (approximately \$A10 000), travel sponsorship from AstraZeneca to attend ATS ASM 2004 (approximately \$A10 000) and two AstraZeneca scientific meetings (approximate combined value \$A20 000), an honorarium paid to his research institute from GlaxoSmithKline to speak at a sponsored conference in South East Asia (\$A3000) in 2005, a travel grant from AstraZeneca to attend the ATS 2005 (\$A6000) and a travel grant from GlaxoSmithKline to attend the ATS 2006 (\$A5000). A proportion of Dr King's research work is conducted at the Woolcock Institute of Medical Research which receives unrestricted grants from AstraZeneca, GlaxoSmithKline, Boehringer Ingelheim and Vita Medical for such work. The Woolcock Institute of Medical Research also has a consultancy agreement with Pfizer, Boehringer Ingelheim, AstraZeneca and GlaxoSmithKline for which Dr King provides consultancy services related to asthma and COPD. His research group receives a proportion of the grants and monies that arise from those companies as part of specific and general allocations of those funds for research purposes across all research groups of the Woolcock Institute of Medical Research. Dr King does not own any stocks, equity or patents that pose a conflict of interest.

REFERENCES

- Macklem P. Bronchial hyporesponsiveness. *Chest* 1987;**91**:189-91S.
- Leuppi JD, Salome CM, Jenkins CR, *et al*. Predictive markers of asthma exacerbation during stepwise dose reduction of inhaled corticosteroids (see comment). *Am J Respir Crit Care Med* 2001;**163**:406-12.
- Xuan W, Peak J, Toelle B, *et al*. Lung function growth and its relation to airway hyperresponsiveness and recent wheeze. Results from a longitudinal population study. *Am J Respir Crit Care Med* 2000;**161**:1820-4.
- Reid DW, Johns DP, Feltis B, *et al*. Exhaled nitric oxide continues to reflect airway hyperresponsiveness and disease activity in inhaled corticosteroid-treated adult asthmatic patients. *Respirology* 2003;**8**:479-86.
- Gustafsson PM, Ljungberg HK, Kjellman B. Peripheral airway involvement in asthma assessed by single-breath SF₆ and He washout. *Eur Respir J* 2003;**21**:1033-9.
- Report National Heart, Lung and Blood Institute/World Health Organization. *Pocket guide for asthma management and prevention - Global Initiative for Asthma* 2005:1-27.
- Obase Y, Shimoda T, Mitsuta K, *et al*. Correlation between airway hyperresponsiveness and airway inflammation in a young adult population: eosinophil, ECP, and cytokine levels in induced sputum. *Ann Allergy Asthma Immunol* 2001;**86**:304-10.
- Crimi E, Spanevello A, Neri M, *et al*. Dissociation between airway inflammation and airway hyperresponsiveness in allergic asthma. *Am J Respir Crit Care Med* 1998;**157**:4-9.
- Moller GM, Overbeek SE, van Helden-Meeuwsen CG, *et al*. Eosinophils in the bronchial mucosa in relation to methacholine dose-response curves in atopic asthma. *J Appl Physiol* 1999;**86**:1352-6.
- Simpson JL, Scott RJ, Boyle MJ, *et al*. Differential proteolytic enzyme activity in eosinophilic and neutrophilic asthma. *Am J Respir Crit Care Med* 2005;**172**:559-65.
- Lanz MJ, Leung DY, McCormick DR, *et al*. Comparison of exhaled nitric oxide, serum eosinophilic cationic protein, and soluble interleukin-2 receptor in exacerbations of pediatric asthma. *Pediatr Pulmonol* 1997;**24**:305-11.
- Venegas JG, Winkler T, Musch G, *et al*. Self-organized patchiness in asthma as a prelude to catastrophic shifts. *Nature* 2005;**434**:777-82.
- Thorpe CW, Bates JHT. Effect of stochastic heterogeneity on lung impedance during acute bronchoconstriction: a model analysis. *J Appl Physiol* 1997;**82**:1616-25.
- Lutchen KR, Gillis H. Relationship between heterogeneous changes in airway morphometry and lung resistance and elastance. *J Appl Physiol* 1997;**83**:1192-201.
- Verbanck S, Schuermans D, Meysman M, *et al*. Noninvasive assessment of airway alterations in smokers. The small airways revisited. *Am J Respir Crit Care Med* 2004;**170**:414-9.
- Crawford AB, Makowska M, Paiva M, *et al*. Convection- and diffusion-dependent ventilation maldistribution in normal subjects. *J Appl Physiol* 1985;**59**:838-46.
- Verbanck S, Schuermans D, Noppen M, *et al*. Evidence of acinar airway involvement in asthma. *Am J Respir Crit Care Med* 1999;**159**:1545-50.
- National Heart Lung and Blood Institute. *Guidelines for the diagnosis and management of asthma*. Washington, DC: NIH, 1997.
- Global Initiative for Asthma. *Pocket guide for asthma management and prevention*. Medical Communications Resources, 2004.
- Juniper EF, O'Byrne PM, Guyatt GH, *et al*. Development and validation of a questionnaire to measure asthma control. *Eur Respir J* 1999;**14**:902-7.
- Salome CM, Roberts AM, Brown NJ, *et al*. Exhaled nitric oxide measurements in a population sample of young adults. *Am J Respir Crit Care Med* 1999;**159**:911-6.
- Deykin A, Massaro AF, Drazen JM, *et al*. Exhaled nitric oxide as a diagnostic test for asthma: online versus offline techniques and effect of flow rate (see comment). *Am J Respir Crit Care Med* 2002;**165**:1597-601.
- King GG, Downie SR, Verbanck S, *et al*. Effects of methacholine on small airway function measured by forced oscillation technique and multiple breath nitrogen washout in normal subjects. *Respir Physiol Neurobiol* 2005;**148**:165-77.
- Yan K, Salome CM, Woolcock AJ. Rapid method for the measurement of bronchial responsiveness. *Thorax* 1983;**38**:760-5.
- O'Connor G, Sparrow D, Taylor D, *et al*. Analysis of dose response curves to methacholine. An approach suitable for population studies. *Am Rev Respir Dis* 1987;**136**:1412-7.
- King GG, Carroll JD, Muller NL, *et al*. Heterogeneity of narrowing in normal and asthmatic airways measured by HRCT. *Eur Respir J* 2004;**24**:211-8.
- Rodriguez-Roisin R. Gas exchange abnormalities in asthma. *Lung* 1990;**168**(Suppl):599-605.
- Beier J, Beeh KM, Kornmann O, *et al*. Sputum induction leads to a decrease of exhaled nitric oxide unrelated to airflow. *Eur Respir J* 2003;**22**:354-7.
- Ward C, Pais M, Bish R, *et al*. Airway inflammation, basement membrane thickening and bronchial hyperresponsiveness in asthma (see comment). *Thorax* 2002;**57**:309-16.
- Vignola AM, Mirabella F, Costanzo G, *et al*. Airway remodeling in asthma. *Chest* 2003;**123**(3 Suppl):417-22S.
- O'Riordan TG, Walser TGL, Smaldone GC. Changing patterns of aerosol deposition during methacholine bronchoprovocation. *Chest* 1993;**103**:1385-9.
- Venegas JG, Schroeder T, Harris S, *et al*. The distribution of ventilation during bronchoconstriction is patchy and bimodal: a PET imaging study. *Respir Physiol Neurobiol* 2005;**148**:57-64.
- Verbanck S, Schuermans D, Paiva M, *et al*. Nonreversible conductive airway ventilation heterogeneity in mild asthma. *Am J Respir Crit Care Med* 2003;**167**:1380-6.
- Aurora P, Kozłowska W, Stocks J. Gas mixing efficiency from birth to adulthood measured by multiple-breath washout. *Respir Physiol Neurobiol* 2005;**148**:125-39.