

- 93 **Gabbay E**, Dark JH, Ashcroft T, *et al*. Recurrence of Langerhans' cell granulomatosis following lung transplantation. *Thorax* 1998;**53**:326–7.
- 94 **Habib SB**, Congleton J, Carr D, *et al*. Recurrence of recipient Langerhans' cell histiocytosis following bilateral lung transplantation. *Thorax* 1998;**53**:323–5.
- 95 **Sullivan EJ**. Lymphangioleiomyomatosis: a review. *Chest* 1998;**114**:1689–703.
- 96 **Kristof AS**, Moss J. Lymphangioleiomyomatosis. In: Schwarz MI, King TE, eds. *Interstitial lung disease*. Hamilton, Ontario: B C Decker, 2003:851–64.
- 97 **Aubry MC**, Myers JL, Ryu JH, *et al*. Pulmonary lymphangioleiomyomatosis in a man. *Am J Respir Crit Care Med* 2000;**162**(2 Pt 1):749–52.
- 98 **Taylor JR**, Ryu J, Colby TV, *et al*. Lymphangioleiomyomatosis. Clinical course in 32 patients. *N Engl J Med* 1990;**323**:1254–60.
- 99 **Chu SC**, Horiba K, Usuki J, *et al*. Comprehensive evaluation of 35 patients with lymphangioleiomyomatosis. *Chest* 1999;**115**:1041–52.
- 100 **Pallisa E**, Sanz P, Roman A, *et al*. Lymphangioleiomyomatosis: pulmonary and abdominal findings with pathologic correlation. *Radiographics* 2002;**22**(Special No):S185–98.
- 101 **Avila NA**, Kelly JA, Chu SC, *et al*. Lymphangioleiomyomatosis: abdominopelvic CT and US findings. *Radiology* 2000;**216**:147–153.
- 102 **Matsui K**, Tatsuguchi A, Valencia J, *et al*. Extrapulmonary lymphangioleiomyomatosis (LAM): clinicopathologic features in 22 cases. *Hum Pathol* 2000;**31**:1242–8.
- 103 **Neumann HP**, Schwarzkopf G, Henske EP. Renal angiomyolipomas, cysts, and cancer in tuberous sclerosis complex. *Semin Pediatr Neurol* 1998;**5**:269–75.
- 104 **Moss J**, Avila NA, Barnes PM, *et al*. Prevalence and clinical characteristics of lymphangioleiomyomatosis (LAM) in patients with tuberous sclerosis complex. *Am J Respir Crit Care Med* 2001;**164**:669–71.
- 105 **Costello LC**, Hartman TE, Ryu JH. High frequency of pulmonary lymphangioleiomyomatosis in women with tuberous sclerosis complex. *Mayo Clin Proc* 2000;**75**:591–4.
- 106 **Carsillo T**, Astrinidis A, Henske EP. Mutations in the tuberous sclerosis complex gene TSC2 are a cause of sporadic pulmonary lymphangioleiomyomatosis. *Proc Natl Acad Sci USA* 2000;**97**:6085–90.
- 107 **Aicher LD**, Campbell JS, Yeung RS. Tuberin phosphorylation regulates its interaction with hamartin. Two proteins involved in tuberous sclerosis. *J Biol Chem* 2001;**276**:21017–21.
- 108 **Nellist M**, van Slegtenhorst MA, Goedbloed M, *et al*. Characterization of the cytosolic tuberin-hamartin complex. Tuberin is a cytosolic chaperone for hamartin. *J Biol Chem* 1999;**274**:35647–52.
- 109 **Ferrans VJ**, Yu ZX, Nelson WK, *et al*. Lymphangioleiomyomatosis (LAM): a review of clinical and morphological features. *J Nippon Med Sch* 2000;**67**:311–29.
- 110 **Maruyama H**, Seyama K, Sobajima J, *et al*. Multifocal micronodular pneumocyte hyperplasia and lymphangioleiomyomatosis in tuberous sclerosis with a TSC2 gene. *Mod Pathol* 2001;**14**:609–14.
- 111 **Kalassian KG**, Doyle R, Kao P, *et al*. Lymphangioleiomyomatosis: new insights. *Am J Respir Crit Care Med* 1997;**155**:1183–6.
- 112 **Aberle DR**, Hansell DM, Brown K, *et al*. Lymphangiomyomatosis: CT, chest radiographic, and functional correlations. *Radiology* 1990;**176**:381–7.
- 113 **Lynch DA**, Brown KK, Lee J-S, *et al*. Imaging of diffuse infiltrative lung disease. In: Lynch DA, Newell JD, Lee J-S, eds. *Imaging of diffuse lung disease*. Hamilton, Ontario and Lewiston, NY: B C Decker, 2000:57–140.
- 114 **Taveira-DaSilva AM**, Stylianou MP, Hedin CJ, *et al*. Decline in lung function in patients with lymphangioleiomyomatosis treated with or without progesterone. *Chest* 2004;**126**:1867–74.
- 115 **Urban T**, Lazor R, Lacronique J, *et al*. Pulmonary lymphangioleiomyomatosis. A study of 69 patients. Groupe d'Etudes et de Recherche sur les Maladies "Orphelines" Pulmonaires (GERM"O" P). *Medicine (Baltimore)* 1999;**78**:321–37.
- 116 **Johnson SR**, Tattersfield AE. Decline in lung function in lymphangioleiomyomatosis: relation to menopause and progesterone treatment. *Am J Respir Crit Care Med* 1999;**160**:628–33.
- 117 **Kenerson HL**, Aicher LD, True LD, *et al*. Activated mammalian target of rapamycin pathway in the pathogenesis of tuberous sclerosis complex renal tumors. *Cancer Res* 2002;**62**:5645–50.
- 118 **Tyrrell VJ**, Asher MI, Chan Y. Subpleural lung cysts in Down's syndrome. *Pediatr Pulmonol* 1999;**28**:145–8.
- 119 **Joshi VV**, Kasznica J, Ali Khan MA, *et al*. Cystic lung disease in Down's syndrome: a report of two cases. *Pediatr Pathol* 1986;**5**:79–86.
- 120 **Nickerson ML**, Warren MB, Toro JR, *et al*. Mutations in a novel gene lead to kidney tumors, lung wall defects, and benign tumors of the hair follicle in patients with the Birt-Hogg-Dube syndrome. *Cancer Cell* 2002;**2**:157–64.
- 121 **Schmidt LS**, Warren MB, Nickerson ML, *et al*. Birt-Hogg-Dube syndrome, a genodermatosis associated with spontaneous pneumothorax and kidney neoplasia, maps to chromosome 17p11.2. *Am J Hum Genet* 2001;**69**:876–82.
- 122 **Kupres KA**, Krivda SJ, Turiansky GW. Numerous asymptomatic facial papules and multiple pulmonary cysts: a case of Birt-Hogg-Dube syndrome. *Cutis* 2003;**72**:127–31.

## LUNG ALERT

### Invariant natural killer T cells in asthma and COPD: back to square one!

▲ Vijayanand P, Seumois G, Pickard C, *et al*. Invariant natural killer T cells in asthma and chronic obstructive pulmonary disease. *N Engl J Med* 2007;**356**:1410–22.

Recent studies have suggested the possibility of invariant natural killer T cells (iNKT) playing an important role in the pathogenesis of asthma. To explore this hypothesis, the authors measured the numbers of iNKT in the airways of patients with stable mild/moderate asthma, patients with stable or exacerbated chronic obstructive pulmonary disease (COPD) and controls.

Cells from induced sputum, bronchoalveolar lavage (BAL) and/or bronchial biopsies were labelled with fluorescent monoclonal antibodies for CD3, CD4 and iNKT specific domains: V $\alpha$ 24, V $\beta$ 11, V $\alpha$ 24-J $\alpha$ 18 and CD1d tetramers loaded with  $\alpha$ galactosylceramide. Flow cytometry with serial gating was used for phenotyping and excluded cells other than iNKT—the failure to do so may have been a drawback of a previous study. Quantitative polymerase chain reaction to detect gene expression for V $\alpha$ 24 and V $\beta$ 11 domains was also used.

iNKT constituted <2% of CD3+ cells and <1.5% of CD3+/CD4+ cells in BAL from patients with asthma. Positive controls confirmed the accuracy of antibodies/gating strategy to detect iNKT. Similarly, very few iNKT were identified in induced sputum of controls, patients with asthma and patients with COPD with no significant differences among groups. iNKT constituted <1.7% of CD3+ cells on biopsy analysis. BAL and sputum cells from patients with asthma and COPD did not express mRNA for V $\alpha$ 24 and V $\beta$ 11, despite the presence of T cell receptors. This ruled out the presence of numerous iNKT among T cells and confirmed flow cytometry findings.

The authors concluded that iNKT are a minority cell population in airways of patients with asthma and COPD. The authors questioned the role of iNKT in the pathogenesis of asthma. These findings, in contrast with a recent report, suggested that future therapeutic strategies should continue to focus on class II major histocompatibility complex restricted cells.

**S A Nachman**

Assistant Professor of Clinical Medicine, Columbia University Medical Center at Harlem Hospital, New York, NY, USA; san14@columbia.edu