14. Chinn S, Rona RJ. Height and age adjustment for cross sectional studies of lung function in children aged 6–11 years. *Thorax* 1992;**47**:707–14.
15. Yan K, Salome C, Woolcock AJ. Rapid method for measurement of bronchial responsiveness. *Thorax* 1983;**38**:760–5.
16. Rabe-Hesketh S, Skrondal A. Classical latent variable models for medical research. *Stat Methods Med Res* 2008;**17**:5–32.
17. Stein RT, Holberg CJ, Morgan WJ, et al. Peak flow variability, methacholine responsiveness and atopy as markers for detecting different wheezing phenotypes in childhood. *Thorax* 1997;**52**:946–52.
18. Turner SW, Palmer LJ, Rye PJ, et al. Infants with flow limitation at 4 weeks: outcome at 6 and 11 years. *Am J Respir Crit Care Med* 2002;**165**:1294–8.
19. Oddy WH, Holt PG, Sly PD, et al. Association between breast feeding and asthma in 6 year old children: findings of a prospective birth cohort study. *BMJ* 1999;**319**:815–9.
20. Young AF, Powers JR, Bell SL. Attrition in longitudinal studies: who do you lose? *Aust NZ J Public Health* 2006;**30**:353–61.
21. Baker D, Henderson J. Differences between infants and adults in the social aetiology of wheeze. The ALSPAC Study Team. Avon Longitudinal Study of Pregnancy and Childhood. *J Epidemiol Community Health* 1999;**53**:636–42.
22. Asher MI, Keil U, Anderson HR, et al. International Study of Asthma and Allergies in Childhood (ISAAC): rationale and methods. *Eur Respir J* 1995;**8**:483–91.
23. Elphick HE, Sherlock P, Foxall G, et al. Survey of respiratory sounds in infants. *Arch Dis Child* 2001;**84**:35–9.
24. Elphick HE, Ritson S, Rodgers H, et al. When a “wheeze” is not a wheeze: acoustic analysis of breath sounds in infants. *Eur Respir J* 2000;**16**:593–7.
25. Lowe LA, Simpson A, Woodcock A, et al. Wheeze phenotypes and lung function in preschool children. *Am J Respir Crit Care Med* 2005;**171**:231–7.
26. Guilbert TW, Morgan WJ, Zeiger RS, et al. Long-term inhaled corticosteroids in preschool children at high risk for asthma. *N Engl J Med* 2006;**354**:1985–97.
27. Murray CS, Woodcock A, Langley SJ, et al. Secondary prevention of asthma by the use of Inhaled Fluticasone propionate in Wheezy Infants (IFWIN): double-blind, randomised, controlled study. *Lancet* 2006;**368**:754–62.
28. van der Velden V, Laan MP, Baert MR, et al. Selective development of a strong Th2 cytokine profile in high-risk children who develop atopy: risk factors and regulatory role of IFN-gamma, IL-4 and IL-10. *Clin Exp Allergy* 2001;**31**:997–1006.
29. Sporik R, Holgate ST, Platts-Mills TA, et al. Exposure to house-dust mite allergen (Der p I) and the development of asthma in childhood. A prospective study. *N Engl J Med* 1990;**323**:502–7.
30. Brussee JE, Smit HA, van Strien RT, et al. Allergen exposure in infancy and the development of sensitization, wheeze, and asthma at 4 years. *J Allergy Clin Immunol* 2005;**115**:946–52.
31. Singh AM, Moore PE, Gern JE, et al. Bronchiolitis to asthma: a review and call for studies of gene-virus interactions in asthma causation. *Am J Respir Crit Care Med* 2007;**175**:108–19.
32. Hoffjan S, Nicolae D, Ostrovskaya I, et al. Gene-environment interaction effects on the development of immune responses in the 1st year of life. *Am J Hum Genet* 2005;**76**:696–704.
33. Lau S, Illi S, Sommerfeld C, et al. Transient early wheeze is not associated with impaired lung function in 7-yr-old children. *Eur Respir J* 2003;**21**:834–41.
34. Haland G, Carlsen KC, Sandvik L, et al. Reduced lung function at birth and the risk of asthma at 10 years of age. *N Engl J Med* 2006;**355**:1682–9.
35. Turner SW, Palmer LJ, Rye PJ, et al. The relationship between infant airway function, childhood airway responsiveness, and asthma. *Am J Respir Crit Care Med* 2004;**169**:921–7.
36. International Study of Asthma and Allergies in Childhood (ISAAC) Steering Committee. Worldwide variation in prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and atopic eczema: ISAAC. *Lancet* 1998;**351**:1225–32.

Pulmonary puzzle

An unusual cystic lung lesion

CLINICAL PRESENTATION

An 80-year-old man was admitted because he had experienced haemoptysis for 4 days. He had smoked 40 cigarettes daily for 50 years. He had no medical history except for chronic obstructive pulmonary disease. Physical examination and laboratory data were unremarkable. A chest radiograph revealed a mass lesion in the right upper lobe of the lung. A CT scan of the patient's chest (fig 1A) showed a well defined cystic mass with mural nodules in the medial aspect of the right upper lung. Magnetic resonance imaging (MRI) of the chest showed a well defined cystic mass with multiple mural nodules in the right upper lung zone, 5×7.5×7.5 cm in size (fig 1B). A CT-guided biopsy was performed and the pathology of the specimen indicated chronic inflammation.

Figure 1 (A) CT scan of the chest showing a well defined cystic mass with mural nodules 10×7.5×7.0 cm in the medial aspect of the right upper lung. There were no enlarged mediastinal or hilar nodes. (B) MRI of the thorax showing a well defined cystic mass with mural nodules 8.5×7.5×7.5 cm in the right upper lung.

QUESTION

What is your diagnosis?

See page 1005

This case was submitted by:

T-W Huang, Y-L Cheng, S-C Lee

Division of Thoracic Surgery, Tri-Service General Hospital, National Defense Medical Center, Taipei, Taiwan, ROC

Correspondence to: Professor S-C Lee, Division of Thoracic Surgery, Tri-Service General Hospital, 325, Section 2, Cheng Gong Road, Taipei 114, Taiwan, ROC; leesc001@yahoo.com.tw

Competing interests: None.

Patient consent: Obtained.

Thorax 2008;**63**:980. doi:10.1136/thx.2007.091751