32. **Hybertson BM**, Lee YM, Cho HG, *et al.* Alveolar type II cell abnormalities and peroxide formation in lungs of rats given IL-1 intratracheally. *Inflammation* 2000;**24**:289–303.
33. **Goggel R**, Winoto-Morbach S, Vielhaber G, *et al.* PAF-mediated pulmonary edema: a new role for acid sphingomyelinase and ceramide. *Nat Med* 2004;**10**:155–60.
34. **Roux J**, Kawakatsu H, Gartland B, *et al.* Interleukin-1beta decreases expression of the epithelial sodium channel alpha-subunit in alveolar epithelial cells via a p38 MAPK-dependent signaling pathway. *J Biol Chem* 2005;**280**:18579–89.
35. **Altemeier WA**, Matute-Bello G, Frevert CW, *et al.* Mechanical ventilation with moderate tidal volumes synergistically increases lung cytokine response to systemic endotoxin. *Am J Physiol Lung Cell Mol Physiol* 2004;**287**:L533–42.
36. **Hollingsworth JW**, Li Z, Brass DM, *et al.* CD44 regulates macrophage recruitment to the lung in lipopolysaccharide-induced airway disease. *Am J Respir Cell Mol Biol* 2007;**37**:248–53.
37. **Hollingsworth JW**, Whitehead GS, Lin KL, *et al.* TLR4 signaling attenuates ongoing allergic inflammation. *J Immunol* 2006;**176**:5856–62.
38. **Jiang D**, Liang J, Fan J, *et al.* Regulation of lung injury and repair by Toll-like receptors and hyaluronan. *Nat Med* 2005;**11**:1173–9.
39. **Okutani D**, Han B, Mura M, *et al.* High-volume ventilation induces pentraxin 3 expression in multiple acute lung injury models in rats. *Am J Physiol Lung Cell Mol Physiol* 2007;**292**:L144–53.
40. **Santos CC**, Zhang H, Liu M, *et al.* Bench-to-bedside review: Biotrauma and modulation of the innate immune response. *Crit Care* 2005;**9**:280–6.
41. **Fisher CJ Jr**, Dhainaut JF, Opal SM, *et al.* Recombinant human interleukin 1 receptor antagonist in the treatment of patients with sepsis syndrome. Results from a randomized, double-blind, placebo-controlled trial. Phase III rhIL-1ra Sepsis Syndrome Study Group. *JAMA* 1994;**271**:1836–43.
42. **Opal SM**, Fisher CJ Jr, Dhainaut JF, *et al.* Confirmatory interleukin-1 receptor antagonist trial in severe sepsis: a phase III, randomized, double-blind, placebo-controlled, multicenter trial. The Interleukin-1 Receptor Antagonist Sepsis Investigator Group. *Crit Care Med* 1997;**25**:1115–24.
43. **Matthay MA**, Wiener-Kronish JP. Intact epithelial barrier function is critical for the resolution of alveolar edema in humans. *Am Rev Respir Disease* 1990;**142**:1250–7.
44. **Ware LB**, Matthay MA. Alveolar fluid clearance is impaired in the majority of patients with acute lung injury and the acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2001;**163**:1376–83.

Pulmonary puzzle

ANSWER

From question on page 128

The bronchial biopsy showed normal columnar epithelium along with metaplastic squamous epithelium. Below the mucosa there were nodules of hyaline cartilage showing focal calcification and cancellous bone with thin trabeculae (fig 1D, arrow) and areas of fatty marrow (fig 1D, arrowhead). These features are diagnostic of “tracheobronchopathia osteochondroplastica”.

Tracheobronchopathia osteochondroplastica (TO) is a rare benign condition of unknown aetiology involving the tracheobronchial tree and was first described in 1857. It usually affects adults with equal gender distribution. TO is characterised by the development of multiple osseous and cartilaginous nodules in the submucosa of the trachea and the main bronchi. Typically, these nodules develop in the cartilaginous anterior and lateral parts of the trachea with sparing of the posterior part. The precise incidence and prevalence of this disorder is unknown but autopsy reporting suggests 3/1000. TO may be an incidental diagnosis or patients may present with chronic respiratory symptoms. The abnormal nodules can lead to altering degrees of airway obstruction and can impair clearance of bronchial secretions. CT may show dense submucosal nodules in the trachea and major bronchi with or without calcification with sparing of the posterior wall. Bronchoscopy shows multiple nodular lesions in the above location and the appearances are characteristic. However, these can be misinterpreted. A typical “gritty” feeling on contact of these nodules with the bronchoscope is described. Histology shows variable amount of submucosal cartilaginous and osseous nodules. Bone tissue may be calcified and/or contain haematopoietic bone marrow. This condition usually follows a benign course and in most cases disease progression is undetectable.^{1–4}

Thorax 2008;**63**:153. doi:10.1136/thx.2007.088237a

REFERENCES

1. **Leske V**, Lazor R, Coetmeur D, *et al.* Tracheobronchopathia osteochondroplastica: a study of 41 patients. *Medicine (Baltimore)* 2001;**80**:378–90.
2. **Lundgren R**, Stjernberg NL. Tracheobronchopathia osteochondroplastica. A clinical bronchoscopic and spirometric study. *Chest* 1981;**80**:706–9.
3. **Prince JS**, Duhamel DR, Levin DL, *et al.* Nonneoplastic lesions of the tracheobronchial wall: radiologic findings with bronchoscopic correlation. *Radiographics* 2002;**22**:S215–30.
4. **Karlikaya C**, Yuksel M, Kilicli S, *et al.* Tracheobronchopathia osteochondroplastica. *Respirology* 2000;**5**:377–80.