

time factors[MeSH] OR waiting lists[MeSH] OR delay[text word] OR Timeliness[text word] OR Time[text word] OR prognosis[MESH]
AND 1995/01/01[PDat]: 2007/06/01[PDat] AND Humans[Mesh] AND English[lang]

Studies of timeliness and outcome in lung cancer care

lung neoplasms[MESH] OR lung cancer*[text word]
AND
time factors[MeSH] OR waiting lists[MeSH] OR delay[text word] OR timeliness[text word] AND mortality[MeSH Terms] OR survival rate[MESH] OR survival analysis[MESH] OR survival[MeSH Terms]
AND 1995/01/01[PDat]: 2007/06/01[PDat] AND Humans[Mesh] AND English[lang]

Studies aimed at improving timeliness of lung cancer care

lung neoplasms[MeSH] OR lung cancer*[text word]
AND
delivery of health care[MESH] OR program development[MESH] OR management information systems[MESH] OR outcome and process assessment (health care)[MESH] OR referral and consultation[MESH] OR practice management[MESH]
AND
time factors[MeSH] OR waiting lists[MeSH] OR delay[text word] OR timeliness[text word] OR time[text word]
AND 1995/01/01[PDat]: 2007/06/01[PDat] AND Humans[Mesh] AND English[lang]

Pulmonary puzzle

A 63-year-old male with marked eosinophilia and dyspnoea on exertion

CLINICAL PRESENTATION

A 63-year-old male farmer presented marked eosinophilia. He had no symptoms of fever, night sweats or weight loss and no signs of anaemia, jaundice or lymphadenopathy. The leucocyte count was $17.27 \times 10^9/l$, with 53.1% eosinophils, but no eggs or parasites were found in his faeces. Blood chemistry results were within normal limits except for an elevated alanine aminotransferase level of 77 U/l. Serum total immunoglobulin E (IgE) was high at 340 kU/l. Specific IgG antibodies to *Taenia solium*, *Paragonimus westermani*, *Sparganum mansoni* and *Clonorchis sinensis* were negative. The patient's first absolute eosinophil count during his hospitalisation was $27.25 \times 10^9/l$. Both a chest CT scan and an echocardiograph were normal. A contrast-enhanced CT scan of the liver showed multiple, small, ill-defined, round, low-attenuating nodules with hepatomegaly. There was no evidence of lymphadenopathy on both a chest and abdominal CT scan. Bone marrow specimens revealed normocellular marrow with marked eosinophilia and no evidence of eosinophilic leukaemia. His eosinophils had increased to $46.32 \times 10^9/l$, so he was treated with prednisolone at 1 mg/kg/day. The pronounced eosinophilia improved and he was discharged. We tapered the dose of prednisolone to 0.8 mg/kg/day over 3 months and his eosinophil counts returned to normal at $0.02 \times 10^9/l$.

Four months later, he was admitted again with dyspnoea on exertion and hypoxaemia. The lower lungs exhibited fine bilateral crackles. The leucocyte count was $13.39 \times 10^9/l$ with 0.7% eosinophils ($0.10 \times 10^9/l$). A chest radiograph showed peripheral reticulonodular opacities in both lungs. A high-resolution CT showed fine reticulation and irregular linear opacity with predominant subpleural distribution (fig 1). Pulmonary function tests showed a reduction in diffusing capacity (55% predicted).



Figure 1 High-resolution CT through the lower lung zone on 11 February 2005. The scan shows bilateral irregular linear opacity with predominant subpleural distribution.

QUESTION

What questions might you ask the farmer?

See page 777 for the answer.

S-H Kim,¹ Y Nawa,² H Y Kim,¹ W Kwon,³ S J Yong,¹ S H Jung⁴

¹ Department of Internal Medicine, Yonsei University Wonju College of Medicine, Wonju, Korea; ² Division of Parasitology, Department of Infectious Diseases, Faculty of Medicine, University of Miyazaki, Kiyotake, Miyazaki, Japan; ³ Department of Diagnostic Radiology, Yonsei University Wonju College of Medicine, Wonju, Korea; ⁴ Department of Pathology, Yonsei University Wonju College of Medicine, Wonju, Korea

Correspondence to: Dr S H Jung, Department of Pathology, Yonsei University Wonju College of Medicine, Wonju, Korea; soonheej@yonsei.ac.kr

Competing interests: None.

Patient consent: Obtained.

Provenance and peer review: Not commissioned; externally peer reviewed.

Thorax 2009;64:756. doi:10.1136/thx.2008.101352