

26. **Bergeron A**, Soler P, Kambouchner M, *et al*. Cytokine profiles in idiopathic pulmonary fibrosis suggest an important role for TGF-beta and IL-10. *Eur Respir J* 2003;**22**:69–76.
27. **Carre P**, Leophonte P. [Cytokines and pulmonary fibrosis]. *Rev Mal Respir* 1993;**10**:193–207.
28. **Hodge G**, Hodge S, Chambers D, *et al*. Acute lung transplant rejection is associated with localized increase in T-cell IFN-gamma and TNF-alpha proinflammatory cytokines in the airways. *Transplantation* 2007;**84**:1452–8.
29. **Bates RC**, Mercurio AM. Tumor necrosis factor-alpha stimulates the epithelial-to-mesenchymal transition of human colonic organoids. *Mol Biol Cell* 2003;**14**:1790–800.
30. **Armendariz-Borunda J**, Katayama K, Seyer JM. Transcriptional mechanisms of type I collagen gene expression are differentially regulated by interleukin-1 beta, tumor necrosis factor alpha, and transforming growth factor beta in Ito cells. *J Biol Chem* 1992;**267**:14316–21.
31. **Regan MC**, Kirk SJ, Hurson M, *et al*. Tumor necrosis factor-alpha inhibits in vivo collagen synthesis. *Surgery* 1993;**113**:173–7.
32. **Solis-Herruzo JA**, Brenner DA, Chojkier M. Tumor necrosis factor alpha inhibits collagen gene transcription and collagen synthesis in cultured human fibroblasts. *J Biol Chem* 1988;**263**:5841–5.


## Pulmonary puzzle

### ANSWER

From the question on page 756

The history of the farmer's eating habits and diet had to be taken into consideration and questioned. It was found that he had a history of eating raw meat, with frequent consumption of uncooked dog and cow liver. The level of eosinophil cationic protein was 12.50 µg/l and the bronchoalveolar lavage fluid contained 20.8% monocytes, 4.6% neutrophils, 69.4% lymphocytes and 5.2% eosinophils. An open lung biopsy revealed extensive eosinophilic infiltration and a larval nematode near the pleural surface (fig 1). An ELISA with the larval excretory–secretory antigen of *Toxocara canis* was positive. He was treated with albendazole at 13 mg/kg/day (800 mg/day) for 6 weeks. After 6 months, he had improved chest radiographs and no respiratory symptoms.

Human infection by the nematode *T canis* occurs by ingesting embryonated eggs from the environment or encapsulated larvae from uncooked animal liver.<sup>1</sup> In 1952, Beaver *et al* identified larvae of *T canis* in a liver biopsy specimen and proposed the term visceral larva migrans (VLM).<sup>2</sup> The larvae penetrate the intestinal wall and begin a migration through the tissues. VLM involves several organ systems, such as the liver, lung, skin and, rarely, the central nervous system.<sup>3</sup> Since *T canis* larvae are rarely detected by pathological examination, the most reliable diagnostic method is ELISA with excretory–secretory antigens.<sup>4</sup> We believe that this is the first reported case of toxocariasis by direct detection of a *Toxocara* larva in an adult lung biopsy specimen. Toxocariasis should be considered as a differential diagnosis for adult patients with hypereosinophilic syndrome and a history of eating raw foods.



**Figure 1** The subpleural granuloma associated with chronic organising pneumonia and a multifocal eosinophilic microabscess reveals *Toxocara* larva (inset) with a smooth and thin cuticle (scale bar: 50 µm).

*Thorax* 2009;**64**:777. doi:10.1136/thx.2008.101352a

### REFERENCES

1. **Aragane K**, Akao N, Matsuyama T, *et al*. Fever, cough, and nodules on ankles. *Lancet* 1999;**354**:1872.
2. **Beaver PC**, Snyder CH, Carrera GM, *et al*. Chronic eosinophilia due to visceral larva migrans; report of three cases. *Pediatrics* 1952;**9**:7–19.
3. **Despommier D**. Toxocariasis: clinical aspects, epidemiology, medical ecology, and molecular aspects. *Clin Microbiol Rev* 2003;**16**:265–72.
4. **Jacquier P**, Gottstein B, Stingelin Y, *et al*. Immunodiagnosis of toxocarosis in humans: evaluation of a new enzyme-linked immunosorbent assay kit. *J Clin Microbiol* 1991;**29**:1831–5.