

# Proinflammatory exosomes in bronchoalveolar lavage fluid of patients with sarcoidosis

Khaleda R Qazi,<sup>1</sup> Patricia Torregrosa Paredes,<sup>1</sup> Benita Dahlberg,<sup>2</sup> Johan Grunewald,<sup>2</sup> Anders Eklund,<sup>2</sup> Susanne Gabrielsson<sup>1</sup>

<sup>1</sup>Department of Medicine, Clinical Allergy Research Unit, Karolinska Institutet and University Hospital, Stockholm, Sweden

<sup>2</sup>Department of Medicine, Division of Respiratory Medicine, Lung Research Laboratory, Karolinska Hospital, Stockholm, Sweden

## Correspondence to

Associate Professor Susanne Gabrielsson, Department of Medicine, Clinical Allergy Research Unit, Karolinska Institutet and University Hospital, KS L2:04, SE-171 76 Stockholm, Sweden; [susanne.gabrielsson@ki.se](mailto:susanne.gabrielsson@ki.se)

KRQ and PTP share first authorship of the paper.

Received 25 November 2009

Accepted 29 July 2010

Published Online First

29 September 2010

## ABSTRACT

**Background** Sarcoidosis is a systemic disease of unknown aetiology characterised by granuloma formation and the presence of interferon  $\gamma$  (IFN $\gamma$ )-producing T cells that cause inflammation and tissue damage in multiple organs, especially the lung. Exosomes are nano-sized immunomodulatory vesicles of endosomal origin released from a diverse range of cells and are also found in physiological fluids including bronchoalveolar lavage fluid (BALF) from healthy individuals.

**Objective** To investigate whether exosomes are enriched in the lungs of patients with sarcoidosis compared with healthy individuals and whether they could contribute to pathogenesis.

**Design** BALF exosomes from patients with sarcoidosis (n=36) and healthy controls (n=14) were compared by electron microscopy, flow cytometry, western blot analysis and mass spectrometry. BALF exosomes were incubated with autologous peripheral blood mononuclear cells (PBMCs) or the human bronchial epithelial cell line 16HBE14o-. Cytokines were measured by ELISPOT and ELISA.

**Results** BALF from patients with sarcoidosis showed increased levels of exosomes compared with healthy individuals. Exosomes from patients showed significantly higher expression of MHC class I and II, tetraspanins CD9, CD63 and CD81 as well as neuregulin-1, known to be associated with cancer progression. Furthermore, BALF exosomes from patients induced significantly higher IFN $\gamma$  and interleukin (IL)-13 production in autologous PBMCs compared with healthy individuals and could also stimulate IL-8 production from epithelial cells.

**Conclusion** The results indicate for the first time a role for exosomes in human lung disease with possible contributions to the initiation and progression of inflammation in sarcoidosis. This suggests that exosomes may be a new potential target for the clinical treatment of lung diseases.

## INTRODUCTION

Sarcoidosis is a systemic granulomatous disease with diverse organ system manifestations that predominantly affects the lung. Common symptoms include dyspnoea, cough and fatigue. Spontaneous remission occurs in most patients but some develop chronic disease leading to death in 1–4% of patients. Although the aetiology of sarcoidosis is not known, many observations including the presence of oligoclonal Th1-like CD4+ T cells and macrophages in bronchoalveolar lavage fluid (BALF) and blood, as well as granuloma formation in the lungs, suggest an antigen-driven autoimmune disease.<sup>1–2</sup>

Autoantigens such as vimentin and ATP-synthase have been identified as targets for expanded T cell clones in the lung.<sup>3</sup> Data suggest contributing roles of *Mycobacterium tuberculosis* and *Propionibacterium* spp. in the pathology of sarcoidosis.<sup>4</sup>

Exosomes are nano-sized vesicles (30–100 nm in diameter) produced by inward budding of multivesicular bodies. Fusion of multivesicular bodies with the plasma membrane leads to the extracellular release of intraluminal vesicles now called exosomes.<sup>5</sup> The term exosome was first coined in 1987 by Johnstone who isolated exosome vesicles from culture supernatants<sup>6</sup> and were discovered as a waste product in erythrocyte development. However, a wide variety of other cell types have since been shown to release exosomes, including B cells, dendritic cells (DC), macrophages, mast cells, T cells, epithelial cells, platelets and tumour cells.<sup>5–7</sup> The in vivo role of exosomes still remains elusive, but their presence in several body fluids such as breast milk<sup>8</sup> and plasma<sup>9</sup> suggest a role in vivo. The analysis of exosomes is complicated because of their small size and the lack of exosome-specific markers, hence several methods including electron microscopy are needed to identify them. Exosomes have received great attention lately owing to their potential to induce immune responses or tolerance,<sup>10</sup> depending on their cellular origin. Exosomes loaded with tumour antigens are considered as promising vaccine candidates in cancer<sup>11–12</sup> and infectious diseases such as toxoplasmosis.<sup>13</sup> In 2003 our group isolated exosomes from BALF (BALF exosomes)<sup>14</sup> which had phenotypic similarities to DC-derived exosomes. Studies by Prado *et al* showed that BALF exosomes from tolerised mice can be given intranasally to prevent allergic sensitisation.<sup>15</sup> Furthermore, it was recently shown that BALF exosomes from *Mycobacterium bovis* BCG-infected mice have the capacity to stimulate tumour necrosis factor  $\alpha$  (TNF $\alpha$ ) production in naïve macrophages.<sup>16</sup> Even though the role of exosomes in the lungs or other compartments in humans is not known, they could either participate in immune surveillance<sup>16</sup> or in pathogenesis—for example, by potentiating inflammation. In this study we found major differences between BALF exosomes from patients with sarcoidosis and healthy individuals. Patients showed increased numbers of exosomes in their lung compared with healthy individuals and had higher levels of MHC class I and II, tetraspanins and heat shock protein 70 (HSP70). The novel finding that BALF exosomes contain neuregulin-1 (NRG1), previously connected to cancer progression,<sup>17</sup> suggests a pathological role for exosomes in sarcoidosis. Furthermore, exosomes

from patients could induce IFN $\gamma$  and IL-13 production in autologous peripheral blood mononuclear cells (PBMCs) and IL-8 production in an epithelial cell line. Taken together, these data suggest that exosomes may contribute to the inflammatory state of sarcoidosis.

## METHODS

### Study subjects and collection of BALF

All patients were recruited from Karolinska University Hospital, Sweden and subjected to bronchoscopy with BAL as part of the routine investigation for sarcoidosis. A diagnosis of sarcoidosis was established by several findings including clinical manifestations, radiological findings, BAL cell differential counts with elevated CD4/CD8 ratio and histological findings of granulomas. Disease activity was determined according to established criteria.<sup>18</sup> BALF was obtained from 36 patients with sarcoidosis of median age 42 years (range 23–63); 24 were non-smokers (>2 years) and 12 were current smokers. An overview of the clinical characteristics of the participating subjects is given in table 1. BALF was collected from 14 healthy individuals of median age 25 years (range 20–38), all of whom were non-smokers. The healthy subjects had normal chest x-rays and had no signs of respiratory infection for at least 1 month before BAL. All healthy subjects underwent BAL voluntarily and provided informed consent under protocols approved by the local ethics committee.

Bronchoscopy with BAL (five aliquots of 50 ml phosphate buffered saline (PBS)) was performed as described previously.<sup>19</sup> The BALF was strained through a double layer of Dacron nets (Millipore, Bedford, MA, USA) and centrifuged at 400g for 10 min at 4°C. Cell viability was determined by Trypan blue exclusion and was always >90%. For differential cell counts, cytopspins were prepared at 20g for 3 min and stained in May–Grünwald Giemsa.

### Exosome preparation

Exosomes were isolated by differential ultracentrifugations as described previously<sup>14</sup> with some modifications. BALF was centrifuged at 300g followed by 3000g and 10 000g and filtered through 0.2  $\mu$ m filters (Advantec MFS, Dublin, CA, USA) to remove cell debris and large protein aggregates. Exosomes were pelleted at 140 000g for 2 h 10 min at 4°C. The pellet was dissolved in PBS and the exosomal protein content was measured by D<sub>C</sub> protein assay (Bio-Rad Laboratories, Hercules, CA, USA) according to the manufacturer's instructions.

### Sucrose gradient

BALF exosomes were layered on a continuous sucrose gradient (0.25–2 mM sucrose and 20 mM HEPES/NaOH, pH 7.4) (Sigma-Aldrich, St Louis, MO, USA) prepared manually and centrifuged for 20 h at 79 000g at 4°C. Fractions were collected and the density was determined by refraction index measurements. Each fraction was either coated directly on anti-MHC II beads for analysis by flow cytometry or ultracentrifuged at 200 000g for 35 min for western blot analysis.

### Flow cytometry

Pelleted exosomes were adsorbed onto 4.5  $\mu$ m  $\varnothing$  Dynabeads as described previously.<sup>14</sup> 1  $\mu$ l beads were coated with an amount of BALF exosomes corresponding to 6 ml of the original volume of BALF. Exosomes were stained with antibodies to HLA-DR-FITC, MHC class I-FITC, CD40-FITC, CD54-PE, CD63-PE, CD80-FITC, CD81-PE, CD86-FITC, CD9-FITC (BD Bioscience, San Diego, CA, USA) and MUC-1-FITC (Abcam Ltd, Cambridge,

**Table 1** Clinical characteristics of patients with sarcoidosis: age, gender, smoking habits, mode of onset, chest radiographic disease stage, smoking habits and treatment with inhaled steroids

Patient no.	Age (years)/sex	Smoker	Stage (0–III)*	Treatment	Onset
1	35/M	No	I	No	Acute
2	34/F	No	II	No	Insidious
3	63/M	Yes	II	No	Insidious
4	29/M	Yes	II	No	Acute
5	33/M	Yes	II	No	Insidious
6	58/M	No	I	No	Acute
7	50/M	No	I	No	Acute
8	26/F	Yes	I	No	Insidious
9	24/F	No	II	No	Insidious
10	39/M	No	II	No	Acute
11	39/M	No	I	No	Acute
12	23/M	Yes	I	No	Acute
13	25/F	Yes	I	No	Acute
14	43/M	No	III	No	Insidious
15	44/F	No	II	No	Insidious
16	32/F	No	I	No	Acute
17	41/M	No	III	No	Acute
18	54/M	No	II	Inhaled	Insidious
19	38/M	Yes	I	No	Acute
20	62/F	No	I	No	Insidious
21	44/M	No	III	Inhaled	Insidious
22	63/M	No	0	No	Insidious
23	37/F	No	I	No	Insidious
24	48/M	No	II	No	Acute
25	49/M	Yes	I	No	Insidious
26	44/F	No	I	No	Acute
27	48/F	No	III	No	Insidious
28	52/F	No	I	No	Acute
29	43/M	No	III	No	Insidious
30	36/F	Yes	II	No	Acute
31	34/M	No	I	No	Acute
32	39/M	Yes	I	No	Insidious
33	58/F	No	III	No	Insidious
34	33/M	Yes	I	No	Acute
35	29/F	Yes	I	No	Acute
36	47/F	No	III	Inhaled	Insidious

\*Stages of disease depicted by chest x-ray: stage 0, normal; stage I, bilateral hilar lymphoma (BHL); stage II, BHL and parenchymal infiltrates; stage III, parenchymal infiltrates.

MA, USA) and compared with isotype-matched controls (Becton Dickinson, Erembodegen, Belgium). Samples were analysed by a FACS Calibur flow cytometer (Becton Dickinson) and data were analysed using CellQuest Pro software (Becton Dickinson).

### Transmission electron microscopy

BALF exosomes corresponding to 3 ml of the original volume of BALF was coated to 1  $\mu$ l anti-MHC class II Dynabeads. The beads were dehydrated in ethanol followed by acetone and embedded in LX-112 (Ladd, Burlington, VT, USA) and sections were contrasted as previously described.<sup>8</sup> Digital images were captured by a Mega View III digital camera (Soft Imaging System GmbH, Munster, Germany).

### SDS-PAGE, silver staining and western blot analysis

The pellet from 140 000g centrifugations was denatured in Laemmli sample buffer (Bio-Rad Laboratories), separated on 8–16% SDS polyacrylamide gels (Bio-Rad Laboratories, Hercules, CA, USA) and each lane was loaded with 10  $\mu$ g of exosomes. Gels were either stained with silver staining kit (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's

recommendations or used for immunoblotting. After transfer to PVDF membrane (Millipore, Bedford, MA, USA) and blocking overnight with ECL blocking agent (Amersham Biosciences, Pittsburgh, PA, USA), primary antibodies anti-HLA-DR, anti-HSP70 and  $\beta$ -actin (Santa Cruz Biotechnology, Santa Cruz, CA, USA) or anti-NRG1 (Abcam and Santa Cruz) were added and developed as described previously.<sup>14</sup>

### Mass spectrometry analysis

Protein bands were excised manually from the silver-stained gels and subsequently processed and digested by trypsin using a robotic protein handling system (MassPREP, Waters, Milford, MA, USA). Gel pieces were washed with ammonium bicarbonate/acetonitrile at 40°C followed by reducing proteins using DTT/ammonium bicarbonate. The gels were shrunk in acetonitrile and the proteins were alkylated in iodoacetamide. Trypsin was added and the peptides were extracted with 5% formic acid/2% acetonitrile followed by 2.5% formic acid/50% acetonitrile. The tryptic peptide extracts from in-gel digestion were mixed at 1:1 with a saturated solution of  $\alpha$ -cyano-4-hydroxy cinnamic acid and dried on a standard steel target plate followed by MALDI MS (Voyager DE Pro, Applied Biosystems, Foster City, CA, USA) in positive ion reflectron mode. For database searches and protein identification, the web site of ProteinProspector (<http://prospector.ucsf.edu>) was used.

### IFN $\gamma$ and IL-13 ELISPOT

PBMCs were isolated from freshly drawn whole blood by density centrifugation on Ficoll Paque (Amersham Pharmacia Biotech AB, Uppsala, Sweden). Pelleted exosomes corresponding to 6 ml of the original volume of BALF/ml culture medium were then added to autologous PBMCs (300 000 cells/well) and incubated for 48 h in the anti-IFN $\gamma$  or anti-IL-13 precoated ELISPOT plate (Mabtech, Uppsala, Sweden). Experiments were set up in triplicate and developed according to the manufacturers' instructions. Spots were counted by an ELISPOT reader using AID ELISPOT software (AID, Strassberg, Germany).

### 16HBE14o- cell culture with BALF exosomes

16HBE14o- cells were kindly provided by Dr D Greunert, University of California, San Francisco, USA.<sup>20</sup> Cells were cultured in fibronectin/collagen (BD Biosciences, bovine) coated flasks and grown in minimal essential medium supplemented with 10% fetal calf serum (exosome-depleted), glutamine (200 mM) and penicillin (50 mg/ml) and kept at 37°C with 5% CO<sub>2</sub>. After growth to confluence, cells were trypsinised using Trypsin-EDTA (R&D Systems, Minneapolis, MN, USA). Viability was controlled by trypan blue exclusion.

16HBE14o- cells (50 000 cells/well) were co-cultured in duplicates with BALF exosomes (1  $\mu$ g/well or exosomes corresponding to 6 ml of the original volume of BALF/ml culture medium as indicated) from patients with sarcoidosis or healthy controls in a 24-well plate for 48 h. Supernatants were then collected and stored at -20°C until used for IL-8 ELISA according to the manufacturer's instructions (Biolegend, San Diego, CA, USA).

### Endotoxin test

The endotoxin test was performed on BALF exosomes (9 patients with sarcoidosis, 9 healthy controls) using the Limulus Amebocyte Lysate endotoxin test (Charles River, Wilmington, MA, USA) according to the manufacturer's instructions. No difference in endotoxin levels was detected in exosome preparations between the two groups and no correlation was seen

between the lipopolysaccharide content and cytokine production (data not shown). Lipopolysaccharide levels ranged between 8.1 and 223.2 pg/ml (median 53.4).

### Statistical analyses

The results were expressed as individual data for each group. Data were tested for normality by the D'Agostino and Pearson omnibus normality test. Mean values are shown for normally distributed data, otherwise median values are shown. Unpaired *t* tests were performed on normally distributed data to identify significant differences between experimental groups. Where data were not normally distributed, the Mann-Whitney *U* test was used to compare groups. *p* Values were considered significant at <0.05. Correlations were assessed using the Pearson correlation coefficient for normally distributed data and the Spearman ranked correlation coefficient for non-normally distributed data, and associated *p* values were calculated. Statistical analyses were made with software Graph Pad Prism 5.

## RESULTS

### Characterisation of isolated exosomes


Exosomes were obtained from the BALF of patients with sarcoidosis (*n*=36) and healthy individuals (*n*=14) by differential ultracentrifugation. Patients with sarcoidosis were both smokers and non-smokers, but there was no significant difference in their protein content (figure 1A). The level of BALF exosomes of non-smokers with sarcoidosis was, however, significantly higher than that of healthy individuals (*p*<0.0001). Both electron microscopy (figure 1B, C) and sucrose gradient purification (figure 1D, E) of BAL exosomes revealed a distinct exosome population with typical exosome size (30–100 nm) and density (1.09–1.19 g/ml). They also confirmed that BALF of patients with sarcoidosis contained more exosomes (figure 1B) than BALF from healthy individuals (figure 1C). Immunolabelling showed increased expression of HLA-DR on BALF exosomes from patients with sarcoidosis compared with those from healthy individuals (data not shown). Furthermore, we found a strong correlation between total cell concentration and total exosome protein in BALF (figure 1F).

BALF exosomes were coated to anti-MHC class II beads and analysed by FACS. Exosomes from both smoking and non-smoking patients had significantly upregulated levels of HLA-DR, HLA-ABC, CD9, CD54, CD63 and CD81 compared with healthy individuals (figure 2B–I). There was no difference in CD86 and MUC1 expression (figure 2H,I). CD40 and CD80 were not detected (data not shown). Levels of surface molecules on BALF exosomes did not correlate with stage of disease or disease onset (acute/insidious) (data not shown). A positive correlation was seen between HLA-DR expression and CD9 (*r*=0.62, *p*<0.0001), CD63 (*r*=0.59, *p*<0.0001) (data not shown) and CD81 (*r*=0.79, *p*<0.0001) (figure 3A) on BALF exosomes from both patients and healthy individuals. Interestingly, we found a negative correlation between HLA-DR and MUC1 on BALF exosomes from patients (figure 3B).

### Protein composition and identification

SDS-PAGE was performed with exosomes from 14 patients and 7 healthy controls. Silver staining showed several protein bands ranging from 250 kDa to 10 kDa (figure 4A). Three bands around 50, 20 and 10 kDa (figure 4A, marked with arrows) were stronger for patients with sarcoidosis than for healthy controls in most exosome preparations and were therefore excised and analysed by mass spectrometry (identified proteins are shown





**Figure 1** Enhanced levels of total exosomal protein in bronchoalveolar lavage fluid (BALF) from patients with sarcoidosis. (A) Total exosomal protein content in BALF from patients with sarcoidosis (smokers  $n=12$ , non-smokers  $n=24$ ) and healthy individuals ( $n=14$ ) was determined using the D<sub>C</sub> protein assay. Results are shown as  $\mu\text{g}$  BALF exosomes/ml BALF. Significant differences between groups were assessed using an unpaired t test, \*\*\* $p<0.001$ . (B,C) BALF exosomes from one patient with sarcoidosis (B) and one healthy individual (C) were coated to anti-MHC class II beads. Samples were fixed and analysed by electron microscopy. Scale bar 100 nm. Pictures are from one representative individual out of two. (D) FACS analysis of CD63 expression on anti-MHC II bound exosomes from exosome-containing sucrose gradient fractions. (E) HLA-DR expression by western blotting on the different sucrose density fractions. (F) Correlation analysis between cell concentration in BALF and exosomal protein concentration from patients with sarcoidosis and healthy individuals ( $n=47$ ) assessed using Spearman correlation coefficient ( $r$ ) and associated  $p$  values (\*\*\* $p<0.001$ ).


in table 2). Among the identified proteins we verified the presence of NRG1 on BALF exosomes by sucrose gradient analysis (figure 4B). Western blotting was performed on NRG1 in randomly selected individuals. Three different antibodies directed against full-length NRG1 and different isoforms ((ab2994 (Abcam), SC-348 and SC-80226 (Santa Cruz)) were used. Multiple bands (19–180 kDa) were detected on exosomes when using the SC-348 antibody (9 patients with sarcoidosis, 7 healthy controls, data not shown). However, several high and low molecular weight bands indicative of intracellular cleaved products<sup>21</sup> were prominently stronger in exosomes from patients with sarcoidosis but were hardly detectable in those from healthy controls. Based on the molecular weight previously reported for NRG1, we speculate that the 80–140 kDa immunoreactive bands represent full-length NRG1<sup>21–22</sup> while smaller bands correspond to cleaved NRG1 intracellular fragments or different isoforms.<sup>23</sup> The aberrant upregulation of NRG-1 on exosomes from patients was again proved using the SC-80226 antibody, specifically recognising the NRG1 isoform with a molecular weight of 40 kDa (figure 4C). The intensity of the bands when normalised to loading control showed significantly higher levels of the 40 kDa NRG1 in patients with sarcoidosis than in healthy controls (figure 4D). No significant difference, but the same tendency, was seen for the 44 kDa NRG1  $\beta$  isoform, with a higher level in BALF exosomes from patients compared with healthy controls (figure 4C,F). BALF exosomes from patients with sarcoidosis also showed strong signals for HLA-DR and HSP70 (figure 4C), with significantly higher levels on BALF exosomes from patients with sarcoidosis compared with healthy controls (figure 4E,G).

### Induction of cytokine production from PBMC by BALF exosomes


Sarcoidosis is known as a Th1-mediated disease with IFN $\gamma$  production in the lungs<sup>24</sup> and increased levels of IL-13 have also been reported.<sup>25</sup> We therefore investigated whether BALF exosomes could stimulate autologous PBMC to secrete IFN $\gamma$  as well as IL-13 in vitro measured by ELISPOT. BALF exosomes from patients with sarcoidosis stimulated significantly higher levels of IFN $\gamma$  and IL-13 production in PBMC compared with BALF exosomes from healthy individuals (figure 5A, B). In addition, IFN $\gamma$  spots from exosome stimulation in patients with sarcoidosis were much larger than those obtained with BALF exosomes from healthy individuals. Neither stage of disease (figure 5A–D) nor disease onset (data not shown) correlated with cytokine production.

### BAL exosomes induce the production of proinflammatory IL-8 in lung epithelial cells (16HBE14o-) and autologous PBMC

We further measured levels of IL-8 production by the lung epithelial cell line 16HBE14o- and autologous PBMCs after co-incubation with exosomes from healthy individuals or patients with sarcoidosis. Preliminary results showed that BALF exosomes could trigger significant production of IL-8 in autologous PBMCs compared with unstimulated PBMCs, with no significant difference between the groups (sarcoidosis patients,  $n=7$ ; healthy individuals,  $n=4$ ; data not shown). Epithelial cells also produced IL-8 in response to BALF exosomes. Interestingly, exosomes from patients with sarcoidosis were more potent inducers of IL-8 release than those from healthy individuals, both when adding exosomes from the same original BALF volume (figure 5C) and the same exosomal protein (figure 5D).



**Figure 2** Increased levels of certain exosome-associated proteins from bronchoalveolar lavage fluid (BALF) of patients with sarcoidosis compared with healthy controls. (A) Representative FACS graphs on BALF exosomes from one healthy individual. BALF exosomes from patients with sarcoidosis (smokers n=12, non-smokers n=24) and healthy individuals (n=14) coated to anti-MHC class II beads and stained with antibodies against (B) HLA-DR, (C) HLA-ABC, (D) CD9, (E) CD54, (F) CD63 (G) CD81, (H) CD86 (p=0.390) and (I) MUC-1 (p=0.840). Values are presented as Mean fluorescence intensity (MFI) values normalised to isotype controls. Significant differences between groups were assessed using unpaired t test (B, D, H) or Mann–Whitney test (C, E, G, I). \*p<0.05, \*\*p<0.01, \*\*\*p<0.0001.



**Figure 3** Significant correlation between HLA-DR expression and tetraspanins on bronchoalveolar lavage fluid (BALF) exosomes. (A) Correlation analysis of the expression of HLA-DR and CD81 ( $n=50$ ) on BALF exosomes from healthy individuals and patients with sarcoidosis. (B) HLA-DR and MUC1 ( $n=35$ ) on BALF exosomes from patients with sarcoidosis. Correlation analysis was assessed by Spearman ranked correlation coefficient ( $r$ ) and associated  $p$  values (\*\* $p<0.01$ , \*\*\* $p<0.0001$ ). Exosomes were bound to MHC class II beads. Values are presented as MFI values normalised to isotype controls.


## DISCUSSION

This study shows for the first time that exosomes might be of clinical relevance in a human respiratory disease by potentiating

inflammation. We show that BALF exosomes from patients with sarcoidosis are quantitatively, phenotypically and functionally different from exosomes from healthy individuals. Electron microscopy data showed increased numbers of exosomes in the BALF from patients with sarcoidosis as well as an enrichment of MHC, tetraspanins, HSP70 and NRG1.

Exosomes enriched in HSP70 have previously been shown to be immunostimulatory *in vivo*<sup>26</sup> and might potentiate inflammation. The strong correlation seen between HLA-DR and the tetraspanins (CD9, CD63 and CD81) is explained by the fact that MHC class II is an interaction partner to several members of the tetraspanin family.<sup>27</sup> The negative correlation between HLA-DR and MUC1 in patients with sarcoidosis could be explained by the increase in exosomes from infiltrating cells and thereby a dilution of MUC1-expressing exosomes, likely to derive from epithelial cells.

One major question is whether or not the production of exosomes is beneficial for the patient with sarcoidosis. Exosomes could play a role in the clearance of an underlying infection prior to the onset of sarcoidosis by potentiating the anti-pathogen immune response. Alternatively, exosomes could contribute to the onset of sarcoidosis by exposing self-antigens on their MHC



**Figure 4** Bronchoalveolar lavage fluid (BALF) exosomes from patients with sarcoidosis contain neuregulin-1 (NRG1). (A) Silver staining analysis of BALF exosomes (healthy controls  $n=7$ , sarcoidosis  $n=14$ ). (B) Western blot analysis of NRG1 44 kDa isoform and HLA-DR expression on exosomes purified on sucrose gradient. (C) Western blot analysis of the expression of NRG1 40 kDa isoform, HLA-DR (healthy,  $n=7$ , sarcoidosis,  $n=7$ ), NRG1 44 kDa isoform, HSP70 and  $\beta$ -actin (healthy,  $n=5$ , sarcoidosis,  $n=7$ ) on BALF exosomes. Densitometric quantification of (D) 40 kDa NRG1, (E) HLA-DR, (F) 44 kDa NRG1 ( $p=0.268$ ) and (G) HSP70 on western blot analysis was measured and normalised to loading control ( $\beta$ -actin). Significant differences between groups were assessed using the Mann–Whitney test: \* $p<0.05$ , \*\*\* $p<0.0001$ . AU, arbitrary units.

**Table 2** Protein identifications in bronchoalveolar lavage fluid exosomes by mass spectrometry of bands (10, 20 and 50 kDa) appearing stronger in patients with sarcoidosis by silver staining

Name	Accession code	Corresponding cut band	Mass (kDa)	Function
Ig $\gamma$ -1 chain C region	P01857	50 kDa, 20 kDa, 10 kDa	36106	Interact with protein or protein complex
PHD finger protein 12	Q96QT6	50 kDa	109699	Chromatin-mediated gene regulation
Zink finger protein 749	Q43361	50 kDa	80607	Regulate expression of genes and nucleic acid recognition, reverse transcription and virus assembly
Pro-neuregulin-1, membrane-bound isoform precursor	Q02297	20 kDa	70392	Growth and differentiation of epithelial, glial, neuronal and skeletal muscle cells; inducing expression of acetylcholine receptor; stimulating lobuloalveolar budding and milk production and inducing differentiation of mammary tumour cells
U4/U2 small nuclear ribonucleoprotein Prp31	Q8WWY3	20 kDa	55457	Pre-mRNA splicing

molecules, thus activating autoimmune T cells. We have previously shown in the mouse model that B cell activation is needed for an exosome-mediated T cell response,<sup>28</sup> and that exosomes can carry B cell epitopes. B cell aggregates have been found in sarcoidosis pulmonary lesions.<sup>29</sup> Thus, a role for exosomes in the activation of B cell responses in sarcoidosis is a possibility and could in turn lower the threshold for T cell activation. We propose that the increase in exosome production seen in patients is primarily a result of increased infiltration of cells in the lung. This is supported by the strong correlation seen between the concentration of cells and exosomal protein (figure 1F). However, other factors such as the state of cellular activation or a primary defect in patients could also contribute to the exosomal profile in sarcoidosis. Comparison with an infectious disease with similar levels of cell infiltration such as tuberculosis would resolve some of these questions. Either way, due to the capacity of these exosomes to stimulate cytokine responses, it is likely that they contribute to the inflammatory response in sarcoidosis.


Interestingly, we found NRG1 to be highly upregulated on exosomes from patients with sarcoidosis. NRG1 has not previously been associated with sarcoidosis but has frequently been

implicated in cancer.<sup>17–30</sup> It also serves as a pro-survival and pro-differentiation signal for epithelial cells and induces branching of the developing lung.<sup>31</sup> To our knowledge, this is the first time NRG1 has been detected in sarcoidosis, and an overexpression of NRG1 could have a role in the pathogenesis of sarcoidosis—for example, by promoting cell survival and proliferation. It is noteworthy that sarcoidosis has been associated with an increased incidence of cancer in affected organs.<sup>32</sup> However, a true connection between sarcoidosis and cancer development remains uncertain.

The T cell population in patients with sarcoidosis is skewed<sup>1–2</sup> and only certain clones are activated, suggesting an antigen-specific reaction. Since exosomes from patients induce Th1-like cytokines, such as IFN $\gamma$ , we suggest that exosomes also contribute to inflammation. However, they could also induce IL-13, previously shown to be elevated in BALF cells and PBMCs from patients with sarcoidosis.<sup>25</sup> Whether IL-13 has pro- or anti-inflammatory properties in the lung in patients with sarcoidosis remains an unsettled issue.

We cannot rule out the possibility that exosomes have an immune regulatory role in the lung in sarcoidosis. The modest increase in CD86 compared with MHC class II in exosomes from

**Figure 5** Bronchoalveolar lavage fluid (BALF) exosomes from patients induce interferon- $\gamma$  (IFN $\gamma$ ) and interleukin (IL)-13 in autologous peripheral blood mononuclear cells (PBMCs) and IL-8 in epithelial cells. Incubation of exosomes for 48 h corresponding to 6 ml of original volume of BALF per 500 000 autologous PBMCs. Co-incubations were performed on (A) anti-IFN $\gamma$  (sarcoidosis n=15, healthy n=9) or (B) anti-IL-13 (sarcoidosis n=13, healthy n=9) coated ELISPOT plates. Assays were performed in triplicates and results are expressed as mean spots per  $3 \times 10^5$  PBMCs. (C) IL-8 production by 16HBE14o- cells was measured by ELISA (pg/ml) after 48 h of co-incubation with exosomes from 6 ml original BALF volume or (D) 4  $\mu$ g/ml of exosomal protein from patients with sarcoidosis (n=9) or healthy individuals (n=9). Results are expressed as the mean of duplicates. Production of IL-8 by unstimulated epithelial cells is indicated by the dotted line. Different symbols represent different stages of disease as indicated. Significant differences between groups were assessed using unpaired t test (B–D) or the Mann–Whitney test (A). \*\*p < 0.001, \*\*\*p < 0.0001.





patients with sarcoidosis suggests a reduced co-stimulatory ability in patients. Even though we could stimulate PBMCs from patients in vitro by these exosomes, it is possible that they act in a different way in vivo. The composition of leucocyte subsets in the lung differs from PBMC preparations, and BALF exosomes might be targeting other cells in vivo such as other DC populations. This interaction might therefore result in a different immunological effect such as induction of Th or T regulatory cells. In fact, sarcoidosis has been linked to clinical anergy to common environmental antigens such as *Candida*, possibly due to a reduced antigen presentation function of myeloid DC (mDC).<sup>33</sup>

The mechanism by which BALF exosomes induce IL-8 production in epithelial cells is probably different from that in PBMCs. Mycobacteria have been suggested as causative agents of sarcoidosis and have also been shown to be involved in the activation of epithelial cells.<sup>34</sup> One might speculate that exosomes carrying pathogen-derived molecules such as proteins, RNA or DNA might induce activation through pattern recognition receptors. Alternatively, exosomes from patients might carry pro-inflammatory cytokines. Recent data show that exovesicles from human alveolar epithelial cells can carry TNF $\alpha$  with the ability to induce IL-8 production in epithelial cells.<sup>35</sup>

In view of the number of statistical tests carried out, results which were significant at the 5% level should be interpreted cautiously. However, we feel it is unlikely that these results are due to chance alone because, in the case of HLA-DR expression, we detected significant differences between the groups both by FACS and western blotting analysis (figure 2B and 4E). Furthermore, because HLA-DR is known to be associated with members of the tetraspanin family which is supported by our correlation data (figure 3A), it is likely that the difference in tetraspanin expression seen between groups (figure 2D, G) is truly significant.

In conclusion, this study shows that patients with sarcoidosis have increased production of exosomes in their lungs with the

ability to stimulate cytokine production from autologous PBMCs and epithelial cells, suggesting both an adaptive and an innate mode of activation. These results suggest that exosomes from the BALF of patients with sarcoidosis might have a role in the pathogenesis of the disease and could be of relevance for the development of novel treatment strategies for sarcoidosis and other lung diseases.

**Acknowledgements** We thank the Proteomics and Mass Spectrometry Unit at the Protein Analysis Center at the Karolinska Institutet for helping with mass spectrometry analysis. We are also grateful to Kjell Hultenby for helping with the electron microscopy.

**Funding** This study was supported by grants from the Karolinska Institutet, the Swedish Medical Research Council (no 15242-05-2), the Torsten and Ragnar Söderbergs Foundation, the Stockholm County Council, the Mats Kleberg Foundation, the King Oscar II Jubilee Foundation, the Swedish Cancer Foundation (no. 08 0299), the Swedish Heart-Lung Foundation, the Swedish Society of Medicine, the Centre for Allergy Research Karolinska Institutet, the Swedish Cancer and Allergy Foundation and the Swedish Asthma and Allergy Association's Research Foundation.

**Competing interests** None.

**Ethics approval** This study was conducted with the approval of the local ethics committee of Northern Stockholm.

**Provenance and peer review** Not commissioned; externally peer reviewed.

## REFERENCES

1. Forman JD, Klein JT, Silver RF, *et al.* Selective activation and accumulation of oligoclonal V beta-specific T cells in active pulmonary sarcoidosis. *J Clin Invest* 1994;**94**:1533–42.
2. Mempel M, Flageul B, Suarez F, *et al.* Comparison of the T cell patterns in leprosy and cutaneous sarcoid granulomas. Presence of Valpha24-invariant natural killer T cells in T-cell-reactive leprosy together with a highly biased T cell receptor Valpha repertoire. *Am J Pathol* 2000;**157**:509–23.
3. Wahlstrom J, Dengjel J, Persson B, *et al.* Identification of HLA-DR-bound peptides presented by human bronchoalveolar lavage cells in sarcoidosis. *J Clin Invest* 2007;**117**:3576–82.
4. Drake WP, Newman LS. Mycobacterial antigens may be important in sarcoidosis pathogenesis. *Curr Opin Pulm Med* 2006;**12**:359–63.
5. Denzer K, Kleijmeer MJ, Heijnen HF, *et al.* Exosome: from internal vesicle of the multivesicular body to intercellular signaling device. *J Cell Sci* 2000;**113** (Pt 19):3365–74.
6. Johnstone RM, Adam M, Hammond JR, *et al.* Vesicle formation during reticulocyte maturation. Association of plasma membrane activities with released vesicles (exosomes). *J Biol Chem* 1987;**262**:9412–20.
7. Andre F, Schartz NE, Chaput N, *et al.* Tumor-derived exosomes: a new source of tumor rejection antigens. *Vaccine* 2002;**20**(Suppl 4):A28–31.
8. Admyre C, Johansson SM, Qazi KR, *et al.* Exosomes with immune modulatory features are present in human breast milk. *J Immunol* 2007;**179**:1969–78.
9. Caby MP, Lankar D, Vincendeau-Scherrer C, *et al.* Exosomal-like vesicles are present in human blood plasma. *Int Immunol* 2005;**17**:879–87.
10. Karlsson M, Lundin S, Dahlgren U, *et al.* "Tolerosomes" are produced by intestinal epithelial cells. *Eur J Immunol* 2001;**31**:2892–900.
11. Zitvogel L, Regnault A, Lozier A, *et al.* Eradication of established murine tumors using a novel cell-free vaccine: dendritic cell-derived exosomes. *Nat Med* 1998;**4**:594–600.
12. Escudier B, Dorval T, Chaput N, *et al.* Vaccination of metastatic melanoma patients with autologous dendritic cell (DC) derived-exosomes: results of the first phase I clinical trial. *J Transl Med* 2005;**3**:10.
13. Aline F, Bout D, Amigorena S, *et al.* Toxoplasma gondii antigen-pulsed-dendritic cell-derived exosomes induce a protective immune response against T gondii infection. *Infect Immun* 2004;**72**:4127–37.
14. Admyre C, Grunewald J, Thyberg J, *et al.* Exosomes with major histocompatibility complex class II and co-stimulatory molecules are present in human BAL fluid. *Eur Respir J* 2003;**22**:578–83.
15. Prado N, Marazuela EG, Segura E, *et al.* Exosomes from bronchoalveolar fluid of tolerized mice prevent allergic reaction. *J Immunol* 2008;**181**:1519–25.
16. Bhatnagar S, Shinagawa K, Castellino FJ, *et al.* Exosomes released from macrophages infected with intracellular pathogens stimulate a proinflammatory response in vitro and in vivo. *Blood* 2007;**110**:3234–44.
17. Hayes NV, Gullick WJ. The neuregulin family of genes and their multiple splice variants in breast cancer. *J Mammary Gland Biol Neoplasia* 2008;**13**:205–14.
18. Anon. Statement on sarcoidosis. Joint Statement of the American Thoracic Society (ATS), the European Respiratory Society (ERS) and the World Association of Sarcoidosis and Other Granulomatous Disorders (WASOG) adopted by the ATS Board of Directors and by the ERS Executive Committee, February 1999. *Am J Respir Crit Care Med* 1999;**160**:736–55.

## What is already known on this topic

Exosomes are nano-sized immunomodulatory vesicles of endosomal origin previously found in the BALF of healthy individuals. These exosomes contain both major histocompatibility complex and co-stimulatory molecules and might have a regulatory role in the immune defence in the lung. Sarcoidosis is a Th1-driven inflammatory disease mainly affecting the lungs.

## What this study adds

The results have implications for the use of exosomes as disease markers or targets for clinical treatment of lung diseases. We show that the BALF of patients with sarcoidosis is enriched in exosomes and that these exosomes have increased levels of immunostimulatory molecules such as MHC I and II, CD54 and tetraspanins, as well as NRG1 previously implicated in cancer progression. Exosomes from patients can induce IFN $\gamma$ , IL-13 and IL-8 production, suggesting exosome-mediated activation of both the innate and adaptive immune system. These findings show the clinical relevance of exosomes as inducers of inflammation and suggest their importance in the pathogenesis of sarcoidosis.



19. **Eklund A**, Blaschke E. Relationship between changed alveolar-capillary permeability and angiotensin converting enzyme activity in serum in sarcoidosis. *Thorax* 1986;**41**:629–34.
20. **Cozens AL**, Yezzi MJ, Kunzelmann K, *et al*. CFTR expression and chloride secretion in polarized immortal human bronchial epithelial cells. *Am J Respir Cell Mol Biol* 1994;**10**:38–47.
21. **Frenzel KE**, Falls DL. Neuregulin-1 proteins in rat brain and transfected cells are localized to lipid rafts. *J Neurochem* 2001;**77**:1–12.
22. **Law AJ**, Shannon Weickert C, Hyde TM, *et al*. Neuregulin-1 (NRG-1) mRNA and protein in the adult human brain. *Neuroscience* 2004;**127**:125–36.
23. **Bao J**, Lin H, Ouyang Y, *et al*. Activity-dependent transcription regulation of PSD-95 by neuregulin-1 and Eos. *Nat Neurosci* 2004;**7**:1250–8.
24. **Moller DR**, Forman JD, Liu MC, *et al*. Enhanced expression of IL-12 associated with Th1 cytokine profiles in active pulmonary sarcoidosis. *J Immunol* 1996;**156**:4952–60.
25. **Hauber HP**, Gholami D, Meyer A, *et al*. Increased interleukin-13 expression in patients with sarcoidosis. *Thorax* 2003;**58**:519–24.
26. **Cho JA**, Lee YS, Kim SH, *et al*. MHC independent anti-tumor immune responses induced by Hsp70-enriched exosomes generate tumor regression in murine models. *Cancer Lett* 2009;**275**:256–65.
27. **Engering A**, Pieters J. Association of distinct tetraspanins with MHC class II molecules at different subcellular locations in human immature dendritic cells. *Int Immunol* 2001;**13**:127–34.
28. **Qazi KR**, Gehrman U, Domange Jordo E, *et al*. Antigen-loaded exosomes alone induce Th1-type memory through a B-cell-dependent mechanism. *Blood* 2009;**113**:2673–83.
29. **Fazel SB**, Howie SE, Krajewski AS, *et al*. B lymphocyte accumulations in human pulmonary sarcoidosis. *Thorax* 1992;**47**:964–7.
30. **Laurendeau I**, Ferrer M, Garrido D, *et al*. Gene expression profiling of ErbB receptors and ligands in human meningiomas. *Cancer Invest* 2009;**27**:691–8.
31. **Patel NV**, Acarregui MJ, Snyder JM, *et al*. Neuregulin-1 and human epidermal growth factor receptors 2 and 3 play a role in human lung development in vitro. *Am J Respir Cell Mol Biol* 2000;**22**:432–40.
32. **Askling J**, Grunewald J, Eklund A, *et al*. Increased risk for cancer following sarcoidosis. *Am J Respir Crit Care Med* 1999;**160**:1668–72.
33. **Mathew S**, Bauer KL, Fischroeder A, *et al*. The anergic state in sarcoidosis is associated with diminished dendritic cell function. *J Immunol* 2008;**181**:746–55.
34. **Wickremasinghe MI**, Thomas LH, Friedland JS. Pulmonary epithelial cells are a source of IL-8 in the response to Mycobacterium tuberculosis: essential role of IL-1 from infected monocytes in a NF-kappa B-dependent network. *J Immunol* 1999;**163**:3936–47.
35. **Obregon C**, Rothen-Rutishauser B, Gerber P, *et al*. Active uptake of dendritic cell-derived exosomes by epithelial cells induces the release of inflammatory mediators through a TNF-alpha-mediated pathway. *Am J Pathol* 2009;**175**:696–705.

## Pulmonary puzzle

### ANSWER

From the question on page 1009

The combination of blood eosinophilia persisting for more than 6 months, end-organ damage as demonstrated by lung consolidation and an identified clonal chromosomal abnormality (ie, the FIP1L1/PDGFR $\alpha$  gene arrangement) fulfils the diagnostic criteria for chronic eosinophilic leukaemia (CEL).<sup>1</sup> In the absence of clonal chromosomal abnormalities, our patient would fulfil the criteria for an idiopathic hypereosinophilic syndrome (HES). The lack of relevant travel history along with normal IgE excluded tropical pulmonary eosinophilia as the diagnosis.<sup>2</sup>

The FIP1L1-PDGFR $\alpha$  fusion protein is a constitutively activated tyrosine kinase<sup>3</sup> which occurs due to a deletion on chromosome 4q12. Expression of FIP1L1-PDGFR $\alpha$  in haematopoietic progenitors induces cytokine-independent differentiation of several myeloid lineages, not just eosinophils. Murine studies show that overexpression of interleukin 5 (IL-5) in the presence of FIP1L1-PDGFR $\alpha$  is required to induce eosinophilic tissue infiltration,<sup>4</sup> with studies of human CD34 cells further suggesting that IL-5 helps induce the phenotype but alone is insufficient to induce HES/CEL-like disease.<sup>5</sup>

Treatment was with the tyrosine kinase inhibitor imatinib. Fifteen-year survival in HES prior to imatinib was around 42%. However, imatinib has been shown to induce clinical, haematological and molecular remission in the majority of patients and it is anticipated that long-term survival will significantly improve.<sup>3</sup>

The mechanism for cough in CEL is unknown. We did not demonstrate eosinophilic lung infiltration (no bronchial biopsies

or bronchoalveolar lavage were performed), but suggest that the pulmonary infiltrates seen on the CT scan represent eosinophilic infiltration. This would be consistent with a case of FIP1L1/PDGFR $\alpha$ -related disease who presented with cough,<sup>6</sup> in whom CT scanning showed thickened airways and endobronchial biopsy showed eosinophilic inflammation. The patient also had normal spirometry and a negative histamine challenge test but a sensitive cough response to inhaled capsaicin. We hypothesise that eosinophils or their degranulation products are the cause of the cough, but further studies are required to elucidate exactly how eosinophils induce cough.

*Thorax* 2010;**65**:1024. doi:10.1136/thx.2010.135251

### REFERENCES

1. **Bain B**, Pierre R, Imbert M, *et al*. Chronic eosinophilic leukaemia and the hypereosinophilic syndrome. In: Jaffe ES, Harris NL, Stein H, *et al*, eds. *World Health Organization classification of tumours: pathology and genetics of tumours of haematopoietic and lymphoid tissues*. Lyon, France: IARC Press, 2001:29–31.
2. **Ottesen EA**, Nutman TB. Tropical pulmonary eosinophilia. *Annu Rev Med* 1992;**43**:417–24.
3. **Cools J**, DeAngelo DJ, Gotlib J, *et al*. A tyrosine kinase created by fusion of positive PDGFR $\alpha$  and FIP1L1 genes as a therapeutic target of imatinib in idiopathic hypereosinophilic syndrome. *N Engl J Med* 2003;**348**:1201–14.
4. **Yamada Y**, Rothenberg ME, Lee AW, *et al*. The FIP1L1-PDGFR $\alpha$  fusion gene cooperates with IL-5 to induce murine hypereosinophilic syndrome (HES)/chronic eosinophilic leukaemia(CEL)-like disease. *Blood* 2006;**107**:4071–9.
5. **Buitenhuis M**, Verhagen LP, Cools J, *et al*. Molecular mechanisms underlying FIP1L1-PDGFR $\alpha$ -mediated myeloproliferation. *Cancer Res* 2007;**67**:3759–66.
6. **Chung KF**, Hew M, Score J, *et al*. Cough and hypereosinophilia due to FIP1L1-PDGFR $\alpha$  fusion gene with tyrosine kinase activity. *Eur Respir J* 2006;**27**:230–2.