21. **Mercadier JJ**, Schwartz K, Schiaffino S, *et al*. Myosin heavy chain gene expression changes in the diaphragm of patients with chronic lung hyperinflation. *Am J Physiol* 1998;**274**:L527–34.
22. **Doucet M**, Debigaré R, Joannisse DR, *et al*. Adaptation of the diaphragm and the vastus lateralis muscle in mild to moderate COPD. *Eur Respir J* 2004;**24**:971–9.
23. **American Thoracic Society**. Standards for the diagnosis and care of patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1995;**152**:S77–120.
24. **Quanjer PH**, Tammeling GJ, Cotes JE, *et al*. Lung volumes and forced ventilatory flows. Report working party standardization of lung function tests, European community for steel and coal. Official statement of the European Respiratory Society. *Eur Respir J Suppl* 1993;**16**:5–40.
25. **Heymsfield SB**, Williams PJ, Shils ME, *et al*. *Nutritional assessment by clinical and biochemical methods. Modern nutrition in health and disease*. Philadelphia: Lea & Febiger, 1994:812–41.
26. **Maltais F**, Leblanc P, Whittom F, *et al*. Oxidative enzyme activities of the vastus lateralis muscle and the functional status in patients with COPD. *Thorax* 2000;**55**:848–53.
27. **Blomstrand E**, Celsing F, Fridén J, *et al*. How to calculate human muscle fibre areas in biopsy samples-methodological considerations. *Acta Physiol Scand* 1984;**122**:545–51.
28. **Rabe KF**, Hurd S, Anzueto A, *et al*. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: Gold Executive Summary. *Am J Respir Crit Care Med* 2007;**176**:532–55.
29. **Watz H**, Waschki B, Meyer T, *et al*. Physical activity in patients with COPD. *Eur Respir J* 2009;**33**:262–72.
30. **Orozco-Levi M**. Structure and function of the respiratory muscles in patients with COPD: impairment or adaptation? *Eur Respir J Suppl* 2003;**46**:41s–51s.
31. **Taillandier D**, Aurousseau E, Meynial-Denis D, *et al*. Coordinate activation of lysosomal, Ca^{2+} -activated and ATP-ubiquitin-dependent proteinases in the unweighted rat soleus muscle. *Biochem J* 1996;**316**:65–72.
32. **Degens H**, Alway SE. Control of muscle size during disuse, disease, and aging. *Int J Sports Med* 2006;**27**:94–9.
33. **Caron MA**, Debigaré R, Dekhuijzen PN, *et al*. Comparative assessment of the quadriceps and the diaphragm in patients with copd. *J Appl Physiol* 2009;**107**:952–61.
34. **Degens H**, Swisher AK, Heijdra YF, *et al*. Apoptosis and Id2 expression in diaphragm and soleus muscle from the emphysematous hamster. *Am J Physiol Regul Integr Comp Physiol* 2007;**293**:R135–44.
35. **Bodine SC**, Latres E, Baumhueter S, *et al*. Identification of ubiquitin ligases required for skeletal muscle atrophy. *Science* 2001;**294**:1704–8.
36. **Cho JE**, Fournier M, Da X, *et al*. Time course expression of foxo transcription factors in skeletal muscle following corticosteroid administration. *J Appl Physiol* 2010;**108**:137–45.
37. **Kamei Y**, Miura S, Suzuki M, *et al*. Skeletal muscle FOXO1 (FKHR) transgenic mice have less skeletal muscle mass, down-regulated type I (slow twitch/red muscle) fiber genes, and impaired glycemic control. *J Biol Chem* 2004;**279**:41114–23.
38. **Sandri M**, Sandri C, Gilbert A, *et al*. Foxo transcription factors induce the atrophy-related ubiquitin ligase atrogin-1 and cause skeletal muscle atrophy. *Cell* 2004;**117**:399–412.
39. **Bossola M**, Muscaritoli M, Costelli P, *et al*. Increased muscle proteasome activity correlates with disease severity in gastric cancer patients. *Ann Surg* 2003;**237**:384–9.
40. **Gosker HR**, Engelen MP, van Mameren H, *et al*. Muscle fiber type IIx atrophy is involved in the loss of fat-free mass in chronic obstructive pulmonary disease. *Am J Clin Nutr* 2002;**76**:113–19.

Lung alert

Effect of mild emphysema on left heart function

Severe chronic obstructive pulmonary disease is sometimes associated with cor pulmonale, impaired left ventricular filling and reduced left ventricular stroke volume and cardiac output with preserved left ventricular ejection fraction. However, whether similar changes occur in less severe cases of emphysema and airflow obstruction is unknown.

In this study the authors hypothesised that mild emphysema and airflow obstruction were associated with a decrease in left ventricular end-diastolic volume, left ventricular stroke volume and cardiac output. A total of 2816 individuals (13% current smokers, 38% former smokers and 49% never smokers) with normal lung function as well as airflow obstruction and CT evidence of emphysema were included. Patients with restrictive spirometry and clinical cardiovascular disease were excluded. Emphysema was measured quantitatively on cardiac CT scans, airflow obstruction according to ATS and ERS guidelines and left ventricular volume was determined by cardiac MRI.

After multivariate adjustments, significant associations were noted between left ventricular measurements and both percentage emphysema and the ratio of forced expiratory volume in 1 s to forced vital capacity (FEV₁/FVC). An increase in 10 percentage points in percentage emphysema was associated with a 4.1 ml decrease in left ventricular end-diastolic volume, a 2.7 ml decrease in stroke volume and a 0.19 l/min decrease in cardiac output. A decrease of 10 percentage points in FEV₁/FVC ratio was associated with decreases of 1.7 ml in left ventricular end-diastolic volume, 1.5 ml in stroke volume and 0.10 l/min in cardiac output. These associations were even stronger for current smokers. There were no significant associations between left ventricular ejection fraction and percentage emphysema and FEV₁/FVC ratio.

This study suggests that mild emphysema may be an independent and additional risk factor for cardiac disease. This may have implications for the way we screen and manage patients with mild emphysema.

► **Barr R**, Bluemke D, Ahmed F, *et al*. Percent emphysema, airflow obstruction, and impaired left ventricular filling. *N Engl J Med* 2010;**362**:217–27.

Brendan Mallia-Milanes

Correspondence to Brendan Mallia-Milanes, ST5, Hereford County Hospital, Hereford, HR1 2ER, UK; bmilanes@doctors.org.uk

Published Online First 29 September 2010

Thorax 2010;**65**:970. doi:10.1136/thx.2010.145599