quarters of the year there was no change in the frequency (%) of oxygenation assessment (96.0, 96.1, 97.6, 98.8), initial antibiotic selection (81.9, 77.9, 80.9, 83.7), or initial antibiotic within 6 h (67.1, 64.0, 63.6, 63.5) or blood cultures performed before antibiotics (58.8, 56.3, 57.9, 62.4) but a small rise in smoking cessation advice (36.4, 32.6, 40.3, 45.5). Overall score, whether composite or appropriate did not change.

Individual trusts Of the 22 Trusts with complete results, overall score over the year was <80% in one, 80–85% in two, 85–90% in three, 90–95% in 13 and >95% in three. Comparing last with first quarter scores only five individual Trusts showed a rise in score of >5%, three a fall of >5% and all others changed less than this amount. The biggest change was in the Trust that performed worst in the first quarter. Practical problems identified included accurate case identification and case note data recording. Despite this the Advancing Quality programme is a practical way of measuring QMs in pneumonia. Further work is required to generate improvements in patient care.

S127 IS HEALTHCARE ASSOCIATED PNEUMONIA A DISTINCT CLINICAL PHENOTYPE?

doi:10.1136/thx.2010.150946.28

¹J Taylor, ¹J Chalmers, ¹A Singanayagam, ¹A Akram, ²G Choudhury, ²P Mandal, ²A Hill. ¹University of Edinburgh, Edinburgh, UK; ²Royal Infirmary of Edinburgh, Edinburgh, UK

Introduction US Guidelines define healthcare associated pneumonia (HCAP) as patients with regular contact with healthcare (eg, nursing home residents, patients with recent hospital admissions or regular outpatient clinics). It is argued that such patients are more likely to develop infection with resistant organisms and require broader spectrum antibiotics. UK guidelines do not distinguish between HCAP and community-acquired pneumonia (CAP). The aim of this study was to investigate whether HCAP is a distinct clinical phenotype.

Methods We studied consecutive patients aged 18+ presenting to an emergency department with CAP according to current UK definitions (NHS Lothian 2006–2009). Exclusion criteria included hospital acquired pneumonia, immunosuppressed patients, and patients not actively treated (palliative). Cases were reviewed by two investigators and HCAP patients were defined according to 2005 ATS/IDSA guidelines. Survival was analysed by Kaplan–Meier analysis and logistic regression.

Results This study analysed 1111 consecutive patients. Of these, 224 (20.2%) met the criteria for HCAP (39.3% hospitalised within 3 months, 37.5% nursing home residents, 10.7% recent outpatient appointments, 12.5% 'other'). 96.4% of HCAP patients received standard 'CAP' antibiotic therapy without coverage of *Pseudomonas aeruginosa* or MRSA. Demographic comparison of HCAP and CAP patients showed HCAP patients were significantly older (median age 76 vs 64, $p<0.0001$) and more likely to have co-morbidities, for example, congestive cardiac failure (30% vs 17%, $p<0.0001$), COPD (33.5% vs 20.8%, $p<0.0001$). HCAP patients had higher markers of severity and worse outcomes on univariate analysis. Mean admission CURB65 score was greater (2.32 vs 1.78, $p<0.0001$), median length of stay was longer (7 vs 5 days, $p=0.01$) and 30-day mortality was double that of CAP patients (16.5% vs 8.2%, $p=0.0004$). Kaplan–Meier analysis showed higher mortality for HCAP patients (Log rank test χ^2 13.24 df=1, $p=0.0003$) as shown in Abstract S127 Figure 1. On multivariate analysis, however, after adjustment for age, co-morbidities and initial pneumonia severity, HCAP was not independently associated with increased 30-day mortality AOR 1.13 (0.69–1.84, $p=0.6$).

Abstract S127 Figure 1

Conclusions Healthcare associated pneumonia is common, with around 20% of CAP patients meeting the definition. Although HCAP patients are older and have more co-morbidities, HCAP is not independently associated with 30-day mortality.

S128 ANALYSIS OF VOLATILE BIOMARKERS WITHIN EXHALED BREATH FOR THE DIAGNOSIS OF PNEUMONIA

doi:10.1136/thx.2010.150946.29

P R Boshier, V Mistry, J R Cushnir, S Curtis, S Elkin, O M Kon, N Marczin, G B Hanna. Imperial College, London, UK

Introduction Whilst pneumonia is a common condition affecting both medical and surgical patients there is currently no definitive diagnostic test. One novel approach may be through the non-invasive analysis of volatile metabolites within exhaled breath, released either by pathogens or host cells. The objective of this study was to delineate the metabolic phenotype of the exhaled breath of patients with community acquired (CAP) and post-operative pneumonia.

Methods Breath samples were collected from patients with a confirmed diagnosis of CAP ($n=26$). Targeted analysis of six prominent breath metabolites was performed by selected ion flow tube-mass spectrometry. Metabolites found to be significantly different in the breath of CAP patients were subsequently investigated in patients undergoing both major abdominal and thoracoabdominal surgery ($n=40$). Receiver operating characteristic (ROC) analysis of putative pneumonia biomarkers was performed in an independent cohort of patients with suspected CAP ($n=21$). A diagnosis of pneumonia was established through fulfilment of pre-defined radiological, microbiological, haematological and clinical criteria.

Results Compared to healthy controls, CAP patients had significantly lower breath levels of hydrogen cyanide (6 vs 14ppb, $p=0.001$) and isoprene (72 vs 111ppb, $p=0.014$). Patients who developed postoperative pneumonia ($n=8$) had significantly lower levels of hydrogen cyanide within their breath compared to both those patients who had an uncomplicated recovery (4 vs 13ppb, $p=0.008$) and healthy controls. Whilst patients who underwent thoracoabdominal compared to abdominal surgery had higher breath levels of isoprene at both 96 (114 vs 176ppb, $p=0.02$) and 168 h (119 vs 176ppb, $p=0.02$) following surgery, this did not correlate with the onset of pneumonia. ROC analysis of hydrogen cyanide and isoprene are displayed in Abstract S128 Figure 1. In both medical and surgical patients there was no significant change in the levels of other examined breath metabolites: acetone, ethanol, propanol and acetic acid.