Institute of Environmental Health Sciences. We acknowledge use of genotype data from the British 1958 Birth Cohort DNA collection, funded by the Medical Research Council grant G0000934 and the Wellcome Trust grant 068545/Z/02.

Competing interests None.

Ethics approval Institutional Review Board of University of Pittsburgh, Partners Health Care (Boston), participating NETT centres, Boston VA, Norway, Costa Rica, and Lovelace Respiratory Institute.

Contributors Manuscript preparation: JMB, JCC; data analysis and study design: JMB, YT, SB, TJM, SB, NB, JPZ, MES-O, LA, MHC, BH, AAL, FJ, EF, SD, EKS, JCC; data collection: YT, TJM, MES, LA, AAL, PB, AG, WHA, DAL, EKS, JCC; statistical analysis: JMB, KH, MHC.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

1. **Cho MH**, Boutaoui N, Klanderman BJ, *et al*. Variants in FAM13A are associated with chronic obstructive pulmonary disease. *Nat Genet* 2010;**42**:200–2.
2. **Hirschhorn JN**. Genome-wide association studies—illuminating biologic pathways. *N Engl J Med* 2009;**360**:1699–701.
3. **Pillai SG**, Ge D, Zhu G, *et al*. A genome-wide association study in chronic obstructive pulmonary disease (COPD): identification of two major susceptibility loci. *PLoS Genet* 2009;**5**:e1000421.
4. **Roeder K**, Bacanu SA, Wasserman L, *et al*. Using linkage genome scans to improve power of association in genome scans. *Am J Hum Genet* 2006;**78**:243–52.
5. **Lander ES**, Botstein D. Homozygosity mapping: a way to map human recessive traits with the DNA of inbred children. *Science* 1987;**236**:1567–70.
6. **Lencz T**, Lambert C, DeRosier P, *et al*. Runs of homozygosity reveal highly penetrant recessive loci in schizophrenia. *Proc Natl Acad Sci U S A* 2007;**104**:19942–7.
7. **Morrow EM**, Yoo SY, Flavell SW, *et al*. Identifying autism loci and genes by tracing recent shared ancestry. *Science* 2008;**321**:218–23.
8. **Reich DE**, Schaffner SF, Daly MJ, *et al*. Human genome sequence variation and the influence of gene history, mutation and recombination. *Nat Genet* 2002;**32**:135–42.
9. **Miyazawa H**, Kato M, Awata T, *et al*. Homozygosity haplotype allows a genome-wide search for the autosomal segments shared among patients. *Am J Hum Genet* 2007;**80**:1090–102.
10. **Bell B**, Rose CL, Damon A. The Normative Aging Study: an interdisciplinary and longitudinal study of health and aging. *Aging Hum Dev* 1972;**3**:5–17.
11. **Fishman A**, Martinez F, Naunheim K, *et al*. A randomized trial comparing lung-volume-reduction surgery with medical therapy for severe emphysema. *N Engl J Med* 2003;**348**:2059–73.
12. **Vestbo J**, Anderson W, Coxson HO, *et al*. Evaluation of COPD Longitudinally to Identify Predictive Surrogate End-points (ECLIPSE). *Eur Respir J* 2008;**31**:869–73.
13. **Pauwels RA**, Buist AS, Calverley PM, *et al*. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease. NHLBI/WHO Global Initiative for Chronic Obstructive Lung Disease (GOLD) Workshop summary. *Am J Respir Crit Care Med* 2001;**163**:1256–76.
14. **Pellegrino R**, Viegi G, Brusasco V, *et al*. Interpretative strategies for lung function tests. *Eur Respir J* 2005;**26**:948–68.
15. **Sood A**, Stidley CA, Picchi MA, *et al*. Difference in airflow obstruction between Hispanic and non-Hispanic White female smokers. *COPD* 2008;**5**:274–81.
16. **Bhattacharya S**, Srisuma S, Demeo DL, *et al*. Molecular biomarkers for quantitative and discrete COPD phenotypes. *Am J Respir Cell Mol Biol* 2009;**40**:359–67.
17. **Benjamini Y**, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J Roy Stat Soc B Stat Meth* 1995;**57**:289–300.
18. **DeMeo DL**, Mariani T, Bhattacharya S, *et al*. Integration of genomic and genetic approaches implicates IREB2 as a COPD susceptibility gene. *Am J Hum Genet* 2009;**85**:493–502.
19. **Demeo DL**, Mariani TJ, Lange C, *et al*. The SERPINE2 gene is associated with chronic obstructive pulmonary disease. *Am J Hum Genet* 2006;**78**:253–64.
20. **Klimentidis YC**, Miller GF, Shriver MD. Genetic admixture, self-reported ethnicity, self-estimated admixture, and skin pigmentation among Hispanics and Native Americans. *Am J Phys Anthropol* 2009;**138**:375–83.
21. **Morera B**, Barrantes R, Marin-Rojas R. Gene admixture in the Costa Rican population. *Ann Hum Genet* 2003;**67**:71–80.
22. **Rubin JS**, Osada H, Finch PW, *et al*. Purification and characterization of a newly identified growth factor specific for epithelial cells. *Proc Natl Acad Sci U S A* 1989;**86**:802–6.
23. **Werner S**, Smola H, Liao X, *et al*. The function of KGF in morphogenesis of epithelium and reepithelialization of wounds. *Science* 1994;**266**:819–22.
24. **Ray P**, Devaux Y, Stolz DB, *et al*. Inducible expression of keratinocyte growth factor (KGF) in mice inhibits lung epithelial cell death induced by hyperoxia. *Proc Natl Acad Sci U S A* 2003;**100**:6098–103.
25. <http://www.b58cgenome.sgu.ac.uk/> (accessed 8 Jan 2010).
26. **lafrate AJ**, Feuk L, Rivera MN, *et al*. Detection of large-scale variation in the human genome. *Nat Genet* 2004;**36**:949–51.

Journal club

The increasing importance of innate antimicrobials

The protein short palate, lung and nasal epithelium clone 1 (SPLUNC1) is found in airway epithelium. It is known to have reduced expression in chronic airways diseases and in smokers. This study aimed to demonstrate its antimicrobial properties against *Mycoplasma pneumoniae* infection. The investigators compared the inflammatory and antibacterial responses to *M pneumoniae* infection in transgenic mice deficient in expressing this protein with the responses in mice overexpressing the protein.

The overall results showed SPLUNC1 has antibacterial effects by inhibiting bacterial adherence proteins, thereby inhibiting *M pneumoniae* growth. Following *M pneumoniae* infection, a reduction in tissue inflammation and increase in neutrophil elastase production was seen in those mice with expression of SPLUNC1. Neutrophil elastase is important in relation to infection and was shown to reduce *M pneumoniae* growth when incubated with human sputum neutrophil elastase.

This study shows the potential antibacterial and immunomodulatory functions of SPLUNC1, which may help in the development of novel treatments for chronic airway diseases. In an ever-increasing climate of drug resistance, it emphasises the importance of focusing on host endogenous antimicrobial responses.

► **Fabienne G**, Di YP, Smith SK. SPLUNC1 promotes lung innate defense against mycoplasma pneumoniae in mice. *Am J Pathol* 2011;**178**:2159–67.

A Loughnan

Correspondence to A Loughnan, Homerton University Hospital, London, UK; alice.loughnan@homerton.nhs.uk

Published Online First 31 July 2011

Thorax 2011;**66**:1090. doi:10.1136/thoraxjnl-2011-200801