7. **Sauleda J**, Garcia-Palmer F, Wiesner RJ, *et al*. Cytochrome oxidase activity and mitochondrial gene expression in skeletal muscle of patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1998;**157**:1413–17.
8. **Puente-Maestu L**, Perez-Parra J, Godoy R, *et al*. Abnormal mitochondrial function in locomotor and respiratory muscles of COPD patients. *Eur Respir J* 2009;**33**:1045–52.
9. **Puente-Maestu L**, Perez-Parra J, Godoy R, *et al*. Abnormal transition pore kinetics and cytochrome C release in muscle mitochondria of patients with chronic obstructive pulmonary disease. *Am J Respir Cell Mol Biol* 2009;**40**:746–50.
10. **Marcuello A**, Gonzalez-Alonso J, Calbet JA, *et al*. Skeletal muscle mitochondrial DNA content in exercising humans. *J Appl Physiol* 2005;**99**:1372–7.
11. **Couillard A**, Koechlin C, Cristol JP, *et al*. Evidence of local exercise-induced systemic oxidative stress in chronic obstructive pulmonary disease patients. *Eur Respir J* 2002;**20**:1123–9.
12. **Mercken EM**, Gosker HR, Rutten EP, *et al*. Systemic and pulmonary oxidative stress after single-leg exercise in COPD. *Chest* 2009;**136**:1291–300.
13. **van Helvoort HA**, Heijdra YF, de Boer RC, *et al*. Six-minute walking-induced systemic inflammation and oxidative stress in muscle-wasted COPD patients. *Chest* 2007;**131**:439–45.
14. **Holloszy JO**. Regulation by exercise of skeletal muscle content of mitochondria and GLUT4. *J Physiol Pharmacol* 2008;**59**(Suppl 7):5–18.
15. **Rabe KF**, Hurd S, Anzueto A, *et al*. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med* 2007;**176**:532–55.
16. **Santos JH**, Mandavilli BS, Van HB. Measuring oxidative mtDNA damage and repair using quantitative PCR. *Methods Mol Biol* 2002;**197**:159–76.
17. **Chance B**, Sies H, Boveris A. Hydroperoxide metabolism in mammalian organs. *Physiol Rev* 1979;**59**:527–605.
18. **Beauchamp C**, Fridovich I. Superoxide dismutase: improved assays and an assay applicable to acrylamide gels. *Anal Biochem* 1971;**44**:276–87.
19. **Pilegaard H**, Saltin B, Neufer PD. Exercise induces transient transcriptional activation of the PGC-1 α gene in human skeletal muscle. *J Physiol* 2003;**546**:851–8.
20. **Gosker HR**, Bast A, Haenen GR, *et al*. Altered antioxidant status in peripheral skeletal muscle of patients with COPD. *Respir Med* 2005;**99**:118–25.
21. **Puente-Maestu L**, Tena T, Trascasa C, *et al*. Training improves muscle oxidative capacity and oxygenation recovery kinetics in patients with chronic obstructive pulmonary disease. *Eur J Appl Physiol* 2003;**88**:580–7.
22. **Vina J**, Gomez-Cabrera MC, Borras C, *et al*. Mitochondrial biogenesis in exercise and in ageing. *Adv Drug Deliv Rev* 2009;**61**:1369–74.
23. **Lin J**, Wu H, Tarr PT, *et al*. Transcriptional co-activator PGC-1 α drives the formation of slow-twitch muscle fibres. *Nature* 2002;**418**:797–801.
24. **Pinho RA**, Chiesa D, Mezzomo KM, *et al*. Oxidative stress in chronic obstructive pulmonary disease patients submitted to a rehabilitation program. *Respir Med* 2007;**101**:1830–5.
25. **Silveira LR**, Pilegaard H, Kusuvara K, *et al*. The contraction induced increase in gene expression of peroxisome proliferator-activated receptor (PPAR)- γ coactivator 1 α (PGC-1 α), mitochondrial uncoupling protein 3 (UCP3) and hexokinase II (HKII) in primary rat skeletal muscle cells is dependent on reactive oxygen species. *Biochim Biophys Acta* 2006;**1763**:969–76.
26. **Cadenas E**, Davies KJ. Mitochondrial free radical generation, oxidative stress, and aging. *Free Radic Biol Med* 2000;**29**:222–30.
27. **Giulivi C**, Boveris A, Cadenas E. Hydroxyl radical generation during mitochondrial electron transfer and the formation of 8-hydroxydeoxyguanosine in mitochondrial DNA. *Arch Biochem Biophys* 1995;**316**:909–16.
28. **Richter C**, Gogvadze V, Laffranchi R, *et al*. Oxidants in mitochondria: from physiology to diseases. *Biochim Biophys Acta* 1995;**1271**:67–74.
29. **Shokolenko I**, Venediktova N, Bochkareva A, *et al*. Oxidative stress induces degradation of mitochondrial DNA. *Nucl Acid Res* 2009;**37**:2539–48.
30. **Wang H**, Hiatt WR, Barstow TJ, *et al*. Relationships between muscle mitochondrial DNA content, mitochondrial enzyme activity and oxidative capacity in man: alterations with disease. *Eur J Appl Physiol Occup Physiol* 1999;**80**:22–7.
31. **Mercken EM**, Hageman GJ, Langen RC, *et al*. Decreased exercise-induced expression of NF- κ B regulated genes in muscle of COPD patients. *Chest* Published Online First: 2010 Aug 5. doi:10.1378/chest.10-0275.
32. **Robinson MM**, Richards JC, Hickey MS, *et al*. Acute β -adrenergic stimulation does not alter mitochondrial protein synthesis or markers of mitochondrial biogenesis in adult men. *Am J Physiol Regul Integr Comp Physiol* 2010;**298**:R25–33.

Lung alert

Lactate clearance is not inferior to central venous oxygen saturation as a goal in early sepsis resuscitation

Goal-directed therapy in severe sepsis improves outcome by providing protocol-driven targets for initial resuscitation. Current guidelines use central venous oxygen saturations (ScvO₂) to assess oxygen delivery and metabolism, but this requires specialist equipment and expertise which may limit its use. This study addresses whether a change in serum lactate concentration may act as an alternative.

This prospective multicentre trial randomised 300 patients with severe sepsis or septic shock to one of two resuscitation protocols. Initially, central venous pressure and mean arterial pressure were normalised using crystalloid boluses and vasopressors. Red blood cell transfusion and dobutamine were then used to either achieve a target ScvO₂ of >70% or a lactate clearance of 10% within the first 2 h of resuscitation.

There was no significant difference in absolute in-hospital mortality between the two groups (34 patients (23%) died in the ScvO₂ group vs 25 (17%) in the lactate clearance group). There was no difference in treatment-related adverse events. Only 29 patients received a blood transfusion or dobutamine to achieve these targets.

This study shows that lactate clearance is a simple but effective measure of total body oxygen metabolism which may be comparable to ScvO₂ when resuscitating patients with severe sepsis. However, this study was not blinded and was performed in hospitals with established goal-directed sepsis protocols and hence may not be generalisable to other institutions.

► **Jones AE**, Shapiro NI, Trzeciak S, *et al*. Lactate clearance versus central venous oxygen saturation as goals of early sepsis therapy: a randomized clinical trial. *JAMA* 2010;**303**:739–46.

Amelia Dunscombe

Correspondence to Amelia Dunscombe, Bristol Royal Infirmary, Bristol BS2 8HW, UK; ameliadunscombe@yahoo.co.uk

Published Online First 14 October 2010

Thorax 2011;**66**:127. doi:10.1136/thx.2010.145607