26. **Subbe CP**, Kruger M, Rutherford P, *et al*. Validation of a modified Early Warning Score in medical admissions. *QJM* 2001;**94**:521–6.
27. **Bland JM**, Altman DG. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* 1986;**1**:307–10.
28. **Roberts CM**, Stone RA, Buckingham RJ, *et al*. Acidosis, non-invasive ventilation and mortality in hospitalised COPD exacerbations. *Thorax* 2011;**66**:43–8.
29. **Luo YM**, Moxham J, Polkey MI. Diaphragm electromyography using an oesophageal catheter: current concepts. *Clin Sci (Lond)* 2008;**115**:233–44.
30. **Gandevia SC**, Hudson AL, Gorman RB, *et al*. Spatial distribution of inspiratory drive to the parasternal intercostal muscles in humans. *J Physiol* 2006;**573**:263–75.
31. **Easton PA**, Hawes HG, Doig CJ, *et al*. Parasternal muscle activity decreases in severe COPD with salmeterol-fluticasone propionate. *Chest* 2010;**137**:558–65.
32. **Reilly CC**, Ward K, Jolley CJ, *et al*. Neural respiratory drive, pulmonary mechanics and breathlessness in patients with cystic fibrosis. *Thorax* 2011;**66**:240–6.
33. **Anthonisen NR**, Manfreda J, Warren CP, *et al*. Antibiotic therapy in exacerbations of chronic obstructive pulmonary disease. *Ann Intern Med* 1987;**106**:196–204.
34. **Trappenburg JC**, van Deventer AC, Troosters T, *et al*. The impact of using different symptom-based exacerbation algorithms in patients with COPD. *Eur Respir J* 2011;**37**:1260–8.
35. **Leidy NK**, Wilcox TK, Jones PW, *et al*. Standardizing measurement of chronic obstructive pulmonary disease exacerbations: reliability and validity of a patient-reported diary. *Am J Respir Crit Care Med* 2011;**183**:323–9.
36. **Sharp JT**, Goldberg NB, Druz WS, *et al*. Thoracoabdominal motion in chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1977;**115**:47–56.
37. **Kyroussis D**, Polkey MI, Hammegard CH, *et al*. Respiratory muscle activity in patients with COPD walking to exhaustion with and without pressure support. *Eur Respir J* 2000;**15**:649–55.
38. **Martinez FJ**, Coughlin JI, Celli BR. Factors influencing ventilatory muscle recruitment in patients with chronic airflow obstruction. *Am Rev Respir Dis* 1990;**142**:276–82.
39. **Man WD**, Mustfa N, Nikolettou D, *et al*. Effect of salmeterol on respiratory muscle activity during exercise in poorly reversible COPD. *Thorax* 2004;**59**:471–6.
40. **Legrand A**, Wilson TA, Troyer AD. Mediolateral gradient of mechanical advantage in the canine parasternal intercostals. *J Appl Physiol* 1996;**80**:2097–101.
41. **De Troyer A**, Legrand A, Gayan-Ramirez G, *et al*. On the mechanism of the mediolateral gradient of parasternal activation. *J Appl Physiol* 1996;**80**:1490–4.
42. **De Troyer A**, Leduc D. Effect of diaphragmatic contraction on the action of the canine parasternal intercostals. *J Appl Physiol* 2006;**101**:169–75.
43. **Easton PA**, Hawes HG, Rothwell B, *et al*. Postinspiratory activity of the parasternal and external intercostal muscles in awake canines. *J Appl Physiol* 1999;**87**:1097–101.
44. **Martinez FJ**, Coughlin JI, Celli BR. Respiratory response to arm elevation in patients with chronic airflow obstruction. *Am Rev Respir Dis* 1991;**143**:476–80.
45. **Ward ME**, Eidelman D, Stubbings DG, *et al*. Respiratory sensation and pattern of respiratory muscle activation during diaphragm fatigue. *J Appl Physiol* 1988;**65**:2181–9.
46. **Bellemare F**, Grassino A. Effect of pressure and timing of contraction on human diaphragm fatigue. *J Appl Physiol* 1982;**53**:1190–5.
47. **Zocchi L**, Fitting JW, Majani U, *et al*. Effect of pressure and timing of contraction on human rib cage muscle fatigue. *Am Rev Respir Dis* 1993;**147**:857–64.
48. **Mahler DA**. The measurement of dyspnea during exercise in patients with lung disease. *Chest* 1992;**101**(5 Suppl):242S–7S.
49. **Marin JM**, Montes de Oca M, Rassulo J, *et al*. Ventilatory drive at rest and perception of exertional dyspnea in severe COPD. *Chest* 1999;**115**:1293–300.
50. **Cao Z**, Ong KC, Eng P, *et al*. Frequent hospital readmissions for acute exacerbations of COPD and their associated factors. *Respirology* 2006;**11**:188–95.
51. **Garcia-Aymerich J**, Farrero E, Felez MA, *et al*. Risk factors of readmission to hospital for a COPD exacerbation: a prospective study. *Thorax* 2003;**58**:100–05.
52. **Hurst JR**, Vestbo J, Anzueto A, *et al*. Susceptibility to exacerbation in chronic obstructive pulmonary disease. *N Engl J Med* 2010;**363**:1128–38.

Journal club

Gastric aspirate culture using the microscopic observation drug susceptibility (MODS) technique is superior to other methods for bacteriological confirmation of tuberculosis in children

Bacteriological evidence of tuberculosis in children is difficult to obtain and is not found in the majority of clinically suspected cases. In this study, which was conducted in Peru, gastric aspirates, nasopharyngeal aspirates and stool samples were taken from 218 children with clinically suspected tuberculosis and nasopharyngeal aspirates and stool samples were taken from 238 low-risk controls. These were examined by the microscopic observation drug susceptibility (MODS) technique broth culture, culture on Lowenstein–Jensen medium and PCR to identify the most effective diagnostic techniques.

10% of cases had at least one positive culture, of which 90.9% were detected by MODS culture compared with only 59.1% by Lowenstein–Jensen culture. Isolation by MODS culture was on average 15 days faster and MODS cultures were spoiled by contamination less often. All culture-positive cases had at least one positive result from a gastric aspirate sample. The yield from adding a second gastric aspirate sample to the first was 37% higher. Nasopharyngeal aspirate samples were positive in 54% of culture-positive cases and addition of a second nasopharyngeal aspirate specimen increased the yield by 50%. Stool cultures were positive in only 18% of culture-positive cases. Gastric aspirate PCR identified only half of clinically high-risk culture-positive cases, and 6.6% of controls had at least one positive PCR result. None of these controls developed evidence of clinical tuberculosis during 1 year of follow-up. The rate of false positive PCR results in any individual specimen was 2%.

These data suggest that the technique most likely to provide bacteriological confirmation of pediatric tuberculosis is MODS culture of gastric aspirates.

► **Oberhelman RA**, Soto-Castellares G, Gilman RH, *et al*. Diagnostic approaches for paediatric tuberculosis by use of different specimen types, culture methods, and PCR: a prospective case-control study. *Lancet Infect Dis* 2010;**10**:612–20.

Jane E Ho

Correspondence to Jane E Ho, Foundation Year 2, St Mary's Hospital, London, W2 1NY, UK; jane.ho@doctors.org.uk

Published Online First 22 October 2011

Thorax 2011;**66**:608. doi:10.1136/thx.2010.151902