18. **Ashitani J**, Mukae H, Arimura Y, *et al*. Elevated plasma procoagulant and fibrinolytic markers in patients with chronic obstructive pulmonary disease. *Intern Med* 2002;**41**:181–5.
19. **Davi G**, Basili S, Vieri M, *et al*. Enhanced thromboxane biosynthesis in patients with chronic obstructive pulmonary disease. The Chronic Obstructive Bronchitis and Haemostasis Study Group. *Am J Respir Crit Care Med* 1997;**156**:1794–9.
20. **Ferroni P**, Basili S, Martini F, *et al*. Soluble P-selectin as a marker of platelet hyperactivity in patients with chronic obstructive pulmonary disease. *J Invest Med* 2000;**48**:21–7.
21. **Walter RE**, Wilk JB, Larson MG, *et al*. Systemic inflammation and COPD: the Framingham Heart Study. *Chest* 2008;**133**:19–25.
22. **Fijnheer R**, Frijns CJ, Korteweg J, *et al*. The origin of P-selectin as a circulating plasma protein. *Thromb Haemost* 1997;**77**:1081–5.
23. **Gurbel PA**, Kereiakes DJ, Serebruany VL. Soluble P-selectin is not a surrogate marker for platelet P-selectin: evidence from a multicenter chest pain study group. *J Thromb Thrombolysis* 2000;**10**:15–22.
24. **Kamath S**, Blann AD, Caine GJ, *et al*. Platelet P-selectin levels in relation to plasma soluble P-selectin and beta-thromboglobulin levels in atrial fibrillation. *Stroke* 2002;**33**:1237–42.
25. **Michelson AD**, Barnard MR, Krueger LA, *et al*. Circulating monocyte-platelet aggregates are a more sensitive marker of in vivo platelet activation than platelet surface P-selectin: studies in baboons, human coronary intervention, and human acute myocardial infarction. *Circulation* 2001;**104**:1533–7.
26. **Gawaz M**, Langer H, May AE. Platelets in inflammation and atherogenesis. *J Clin Invest* 2005;**115**:3378–84.
27. **Del Maschio A**, Evangelista V, Rajtar G, *et al*. Platelet activation by polymorphonuclear leukocytes exposed to chemotactic agents. *Am J Physiol* 1990;**258**:H870–9.
28. **Harding SA**, Sommerfield AJ, Sarma J, *et al*. Increased CD40 ligand and platelet-monocyte aggregates in patients with type 1 diabetes mellitus. *Atherosclerosis* 2004;**176**:321–5.
29. **Diodati JG**, Cannon RO 3rd, Epstein SE, *et al*. Platelet hyperaggregability across the coronary bed in response to rapid atrial pacing in patients with stable coronary artery disease. *Circulation* 1992;**86**:1186–93.
30. **Vaidyula VR**, Boden G, Rao AK. Platelet and monocyte activation by hyperglycemia and hyperinsulinemia in healthy subjects. *Platelets* 2006;**17**:577–85.
31. **Smeeth L**, Thomas SL, Hall AJ, *et al*. Risk of myocardial infarction and stroke after acute infection or vaccination. *N Engl J Med* 2004;**351**:2611–18.
32. **Thaulow E**, Erikssen J, Sandvik L, *et al*. Blood platelet count and function are related to total and cardiovascular death in apparently healthy men. *Circulation* 1991;**84**:613–17.
33. **Ly HQ**, Kirtane AJ, Murphy SA, *et al*. Association of platelet counts on presentation and clinical outcomes in ST-elevation myocardial infarction (from the TIMI Trials). *Am J Cardiol* 2006;**98**:1–5.
34. **Cote C**, Zilberberg MD, Mody SH, *et al*. Haemoglobin level and its clinical impact in a cohort of patients with COPD. *Eur Respir J* 2007;**29**:923–9.
35. **Similowski T**, Agustí A, Macnee W, *et al*. The potential impact of anaemia of chronic disease in COPD. *Eur Respir J* 2006;**27**:390–6.
36. **Klinkhardt U**, Bauersachs R, Adams J, *et al*. Clopidogrel but not aspirin reduces P-selectin expression and formation of platelet-leukocyte aggregates in patients with atherosclerotic vascular disease. *Clin Pharmacol Ther* 2003;**73**:232–41.
37. **Seemungal TA**, Donaldson GC, Bhowmik A, *et al*. Time course and recovery of exacerbations in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2000;**161**:1608–13.
38. **Schuerholz T**, Keil O, Wagner T, *et al*. Hydrocortisone does not affect major platelet receptors in inflammation in vitro. *Steroids* 2007;**72**:609–13.
39. **Harding SA**, Din JN, Sarma J, *et al*. Promotion of proinflammatory interactions between platelets and monocytes by unfractionated heparin. *Heart* 2006;**92**:1635–8.

Journal club

SOX9 upregulation in lung adenocarcinoma

Several SOX genes are involved in lung development and may have functional roles within the lung. SOX9 (a transcription factor) has been shown to be involved in several types of cancer. In this lab-based study, SOX9 expression was investigated in 29 lung adenocarcinoma and 28 normal lung samples using silico data mining, quantitative reverse transcription-PCR and immunohistochemistry.

SOX9 was upregulated at both the mRNA and protein level in the majority of lung adenocarcinomas compared with normal tissue. In addition, recurrent lung adenocarcinoma tissue showed significantly higher SOX9 expression than that of a primary lesion. RNA interference technique and microarray gene expression further demonstrated that knockdown of SOX9 inhibited cell growth, downregulated CDK4 and upregulated p21. These are all important cell cycle regulators and their expression could mediate SOX9-related alteration in cancer cell growth.

SOX9 may be an important biomarker in lung adenocarcinoma which could aid in the early detection and diagnosis of NSCLC. This study builds on our existing knowledge of the role of the molecules involved in the pathogenesis of lung adenocarcinoma.

► **Jiang SS**, Fang WT, Hou YH, *et al*. Upregulation of SOX9 in lung adenocarcinoma and its involvement in the regulation of cell growth and tumorigenicity. *Clin Cancer Res* 2010;**16**:4363–73.

L McKinlay

Correspondence to L McKinlay, Clinical Research Fellow, University of Dundee, Level 7 Cardiovascular and Lung Biology, Ninewells Hospital, Dundee DD19SY, UK; l.mckinlay@dundee.ac.uk

Published Online First 3 November 2010

Thorax 2011;**66**:774. doi:10.1136/thx.2010.153536