questionnaire is responsive to a chronic lung disease's population undergoing rehabilitation. A prospective study is now required to expand the application of the CAT to the general assessment of patients with chronic lung diseases.

REFERENCES

1. Jones PW, et al. Development and first validation of the COPD assessment test. *Eur Respir J* 2009;**34**:648–54.
2. Dodd JW, et al. The COPD assessment test (CAT): response to pulmonary rehabilitation. A multicentre, prospective study. *Thorax* 2011;**66**:425–9.

P147 OUTCOMES OF PULMONARY REHABILITATION IN SEVERE ASTHMA

doi:10.1136/thoraxjnl-2011-201054c.147

J Agbetile, S Singh, P Bradding, I Pavord, R Green. *Department of Respiratory Medicine and Thoracic Surgery, Glenfield Hospital, Leicester, UK*

Objectives Structured physical training programmes are a key component of pulmonary rehabilitation (PR) and have consistently been shown to be of significant benefit to patients with COPD. It is unclear whether the application of similar principles provides similar health improvements in patients with severe asthma. We aimed to assess this by studying the effects of pulmonary rehabilitation in a group of patients with severe asthma and significant disability who have completed our standard 7 week PR programme.

Methods 111 patients with a physician diagnosis of severe asthma who have completed our standard PR programme were studied. Assessments including spirometry, chronic respiratory questionnaire (CRQ), hospital anxiety and depression scores (HAD) and incremental and endurance shuttle walk testing (ISWT, ESWT) were performed at baseline and completion to determine the effect of PR. Baseline demographic and spirometric data were compared with a larger population of patients with severe asthma attending our difficult asthma clinic (DAC).

Results Following rehabilitation, statistically significant improvements were seen in MRC dyspnoea score (mean (95% CI) improvement -0.35 (-0.13 to -0.57), $p=0.003$); HADS anxiety (-1.4 (-0.4 to -2.3), $p=0.007$); HADS depression (-1.2 (-0.2 to -2.2), $p=0.026$); CRQ dyspnoea ($+0.72$ ($+0.33$ to $+1.1$), $p=0.001$); CRQ fatigue ($+0.92$ ($+0.65$ to $+1.2$), $p<0.001$); CRQ Emotion ($+0.64$ ($+0.36$ to $+0.92$), $p<0.001$); CRQ mastery ($+0.47$ ($+0.13$ to $+0.81$), $p=0.007$); Incremental shuttle walk test (ISWT) ($+63$ m (49.5 to 76.5), $p<0.001$); and Endurance shuttle walk test (ESWT) ($+339.7$ m, (249.7 to 429.7), $p<0.001$). Interestingly however, patients with severe asthma who completed pulmonary rehabilitation were older (66 vs 48 yrs $p<0.01$), more likely to have smoked (69.5% vs 43.5%, $p<0.01$), and tended to have more severe fixed airflow obstruction (FEV₁% predicted 55.9 vs 71.9, $p=0.28$) than our DAC population. This may be because clinicians were more likely to refer patients with asthma who had a COPD-like phenotype or because patients with severe asthma were more likely to drop-out of the programme.

Conclusion Pulmonary rehabilitation may offer significant benefit to patients with severe asthma though further work is needed to identify the patients mostly likely to benefit and the optimum training required.

P148 THE ADAPTATION AND EVALUATION OF THE LIVING WELL WITH COPD PROGRAMME FOR PULMONARY REHABILITATION

doi:10.1136/thoraxjnl-2011-201054c.148

¹D Earley, ²J MacMahon, ³J Bourbeau, ¹J M Bradley, ¹B O'Neill. *Health and Rehabilitation Sciences Research Institute, University of Ulster, Newtownabbey, Northern*

Ireland; ²Department of Respiratory Medicine, Belfast City Hospital, Belfast, Northern Ireland; ³Respiratory Epidemiology and Clinical Research Unit, Montreal Chest Institute, McGill University Health Centre, Montreal, Canada

Introduction Mechanisms and materials to support the delivery of the education component of pulmonary rehabilitation are not widely available. The aim of this study was to adapt the Living Well with COPD (LWWCOPD) programme for embedding in pulmonary rehabilitation and to conduct a process and outcome evaluation.

Methods *Adaptation:* Modifications to the LWWCOPD programme were informed by focus groups, current practice, relevant research and guidelines, and in collaboration with the authors of the LWWCOPD programme. The study used a cohort before-after design which incorporated the principles of a process evaluation. *Evaluation:* Sites administered their usual pulmonary rehabilitation programme with the exception of the LWWCOPD programme for pulmonary rehabilitation to deliver the education component. Health professionals and patients completed evaluation questionnaires to assess their acceptance. Patients completed the Understanding COPD (UCOPD) questionnaire and the Bristol COPD Knowledge Questionnaire before and after pulmonary rehabilitation. *Analysis:* Feedback and comments on the programme were collated and categorised. Changes in the UCOPD questionnaire and BCKQ were examined using paired t-tests.

Results *Adaptation:* Amendments to the LWWCOPD programme included reducing the number and length of education sessions, incorporating additional information/techniques, materials to link the education and exercise sessions and a COPD action plan. *Evaluation:* 25 health professionals and 57 patients with COPD from eleven pulmonary rehabilitation programmes evaluated the LWWCOPD for pulmonary rehabilitation. The mean (SD) duration of the education sessions was 41 (9) min. The health professionals felt that the education sessions were either excellent ($n=16/65$, 25%) or good ($n=40/65$, 62%), and that they were comprehensive, evidence-based and utilised a good combination of teaching strategies for example, "Easy to follow and deliver". They reported that the programme would require modification for non-COPD patients. Patients commented on improved knowledge and self-efficacy, peer support and relevant content for example, "Better understanding of action to be taken". They requested supplementary information for family members. The UCOPD questionnaire and the BCKQ improved significantly: mean change (95% CI): UCOPD questionnaire: 26.8 (21.7 to 31.8)%, BCKQ: 10.6 (6.9 to 14.4)%.

Conclusion The LWWCOPD programme for pulmonary rehabilitation is a feasible, effective and versatile way to deliver the education component of COPD pulmonary rehabilitation.

P149 PILOTING AND EVALUATING POST-PULMONARY REHABILITATION (PR) LONG-TERM EXERCISE (LTE) FOR COPD PATIENTS

doi:10.1136/thoraxjnl-2011-201054c.149

¹S E Roberts, ¹G Kettle, ²S Rogers, ²A Segal, ³S Purcell, ³G Fabris, ³H Broomfield, ⁴M Calonge-Contreras, ⁴J Billet, ³L Restrict, ³M Stern. *¹Whittington Hospital Physiotherapy, Whittington Health, London, UK; ²Independent Exercise Instructors, London, UK; ³Department Respiratory Medicine, Whittington Health, London, UK; ⁴Department Public Health, Islington, NHS North Central, UK*

Introduction There is evidence that regular exercise profoundly affects both the course and outcome of COPD. While PR is well established for COPD patients limited by breathlessness, there is currently no evidence-base for follow-on LTE, which takes into account expressed desires of patients to continue exercise regularly. This pilot study aimed to evaluate an easily accessible, disease appropriate, regular LTE for COPD patients who complete PR, to

confirm that it is realistic and assess whether PR benefits can be sustained.

Methods Three suitable venues, geographically distributed to maximise access, and two exercise instructors (completed Loughborough Training for Chronic Respiratory Patients) were identified. Patients completing PR were recruited into 1×weekly LTE groups. Outcomes (demographic, disease severity, functional capacity (6 min walk test (6MWT), emotional (Hospital Anxiety and Depression HAD score, CAT and Chronic Respiratory Questionnaire CRQ)) were collected at baseline, 6 and 12 months. Patient and carer satisfaction was recorded at 6 and 12 months. Attendance data were monitored throughout. Hospital admissions for 12/12 before and after LTE commenced, continues to be audited.

Results 60 COPD patients (Mean (SD) age 68.75 (10.31) yrs, FEV₁ 1.27 (0.56) l, 43% male, 22% current smokers, three on LTOT) were referred for LTE June 2010–May 2011. 20/60 (33%) never attended, 14/60 (23%) dropped out within 2 (1–7) months (median (range)). 26/60 (43%) continue to exercise, median (range) since starting 5 (1–12) months with 50 (28)%. (Mean (SD) classes attendance. CAT, CRQ, HADS and 6MWT findings are given in the Abstract P149 table 1. Satisfaction surveys indicate high levels of satisfaction with venue, instructor and content. No adverse events occurred during the classes. Direct cost per class (hall hire and instructor) £63.

Abstract P149 Table 1 Health related quality of life and exercise capacity at end-PR, 6 months and 12 months of long-term exercise

	Baseline, n = 60, mean (SD)	6 Months, n = 11, mean (SD)	12 Months, n = 3, mean (SD)
MRC	3.2 (0.8)		
CAT score	22.2 (6.8)	12.4 (7.4)	17.0 (14.1)
CRQ Dyspnoea	3.3 (1.4)	4.2 (1.6)	4.9 (1.9)
CRQ fatigue	4.1 (1.5)	4.9 (1.2)	4.7 (2.5)
CRQ emotional function	4.8 (1.6)	5.3 (1.3)	4.8 (2.9)
CRQ mastery	4.9 (1.7)	5.8 (1.4)	4.1 (1.9)
HADS anxiety	7.4 (5.1)	4.3 (3.3)	6.0 (6.0)
HADS depression	5.2 (4.2)	3.6 (2.5)	4.7 (4.7)
6 MWT (metres)	353 (116)	338 (83)	330 (122)

Conclusions This pilot demonstrates feasibility of providing community-based LTE groups using instructors trained to exercise people with COPD. Patients who choose to attend LTE have moderate COPD, but high CAT scores indicating important disease impact on daily functioning. Attendance rate was high for a patient group susceptible to exacerbations, possibly reflecting high patient satisfaction. Preliminary findings suggest LTE in group settings promotes maintenance of benefits acquired from PR with further improvement in health-related quality of life (CRQ and CAT).

P150 USE OF AMBULATORY OXYGEN THERAPY IN PULMONARY REHABILITATION

doi:10.1136/thoraxjnl-2011-201054c.150

D Anderson, S Ferrer Valls, S Church, S Agarwal. *St Helens and Knowsley teaching Hospitals NHS Trust, Prescot, UK*

Introduction and Objectives Pulmonary rehabilitation (PR) is proven to improve oxygen exercise capacity, health status and reduce dyspnoea in patients diagnosed with Chronic Obstructive Pulmonary Disease. Previous studies have shown no significant difference in rehabilitation outcomes following use of supplemental oxygen

during pulmonary rehabilitation. Unfortunately, a few patients are unable to exercise due to hypoxia related dyspnoea depriving them of the benefit of PR. We hypothesised that the use of Ambulatory Oxygen Therapy (AOT) in patients with exercise induced desaturation would increase their capability of undergoing PR and have an overall positive outcome.

Methods Requirement of AOT is routinely assessed at our trust prior to commencing PR. Incremental oxygen is used to decide on the flow rate needed during PR. A retrospective study was performed to evaluate the benefits of AOT in patients desaturating on exertion. Data were analysed using Excel.

Results 218 patients were enrolled for the programme from January 2009 to July 2010, 46 excluded as not suitable or did not attend. Numbers studied-172: 59% male, mean age 70 (range 45–90), 25% current smokers. On initial endurance exercise test 27% (n=47) were noted to have exercise induced desaturation. Seven patients were already on long-term oxygen therapy. Median prescribed flow rate was 2 l/min during PR. The mean distance walked on initial endurance test was 150 m. This improved significantly to 263 m (p<0001) with AOT. 68 % (n=32) attended for post PR endurance exercise testing, 25% no longer required AOT (did not desaturate). Of the 22 patients (46%) who continued to require AOT, 14 were prescribed. Following 2 months of the prescription, AOT was withdrawn in four patients as they found the cylinder too heavy or did not benefit.

Conclusions AOT has shown significant improvement in rehabilitation outcomes in patients with exercise induced desaturation. A quarter of these patients did not require any further ambulatory oxygen therapy following pulmonary rehabilitation.

Abstract P150 Table 1

Paired t test	Mean distance on initial endurance exercise test, 150 m	p Values
Mean distance on AOT	263 m	0.0001
Mean distance post PR	277 m	0.0069
Mean distance post PR + AOT	356 m	0.0003

P151 COST OF PULMONARY REHABILITATION IS OFFSET BY REDUCTION IN HEALTHCARE UTILISATION

doi:10.1136/thoraxjnl-2011-201054c.151

¹S Kibe, ¹D Frost, ²S Hart. ¹Scarborough General Hospital, Scarborough, UK; ²Castle Hill Hospital, Hull, UK

151 patients completed pulmonary rehabilitation over a period of over 4 years at our hospital. Retrospective analysis of 71 patients from this group was done. This showed that there was a significant difference between pre and post rehabilitation in FVC, VC, HADS depression score & SWT (shuttle walk test distance). Analysis of a sub group of 37 patients at 6 months after pulmonary rehabilitation showed that there were still significant differences in HADS depression & SWT, but not the other variables. We were able to establish a significant reduction in healthcare utilisation, both in hospital admissions and in out patient attendances. This was independent of the reported level of benefit on CRDQ scores. “Responders” (patients with a more than 1.5 increase in each of their CRDQ scores) had similar reduction in healthcare utilisation compared to “non-responders” (patients with <0.5 difference or deterioration in their CRDQ scores). At our hospital the approximate cost of pulmonary rehabilitation is 375 pounds per person. Out patient (respiratory) appointments and hospital admissions