Occupational lung disease

S1 EVIDENCE-BASED CASE DEFINITION FOR EAA DUE TO METALWORKING FLUID EXPOSURE

doi:10.1136/thoraxjnl-2011-201054b.1

¹C M Barber, ¹C M Burton, ²D J Hendrick, ³C A C Pickering, ⁴A S Robertson, ⁵W Robertson, ⁴P S Burge. ¹Centre for Workplace Health, Buxton, UK; ²University of Newcastle, Newcastle, UK; ³Wythenshawe Hospital, Manchester, UK; ⁴Heart of England Hospital, Birmingham, UK; ⁵University of Warwick, Coventry, UK

Introduction and objectives Although a number of case definitions for MWF EAA have been used in previous outbreaks, none has emerged as an evidence-based standard. This study used data from a large UK MWF outbreak investigation, to develop and validate a new case definition for MWF EAA.

Methods Demographic and clinical data from the 37 workers with suspected EAA during a large outbreak were reviewed by an Expert Panel. A new case definition for MWF EAA was developed using significant differences between workers with and without definite clinical EAA. This MWF EAA Score was modelled to match Expert Panel opinion as closely as possible, and used clinical criteria relatively weighted by their positive predictive value for EAA in the outbreak. The performance of the new scoring system was compared with other existing case definitions, and applied to 50 cases of MWF-EAA from nine published US outbreaks.

Results The MWF EAA Score is shown in Abstract S1 table 1, where the highest score is applied in each section. A score of >26 represents definite EAA, 19–26 possible EAA, and <19 not a case of EAA. When applied to the 37 workers, the MWF EAA Score showed good correlation (coefficient=0.85) with the Expert Panel Score (per cent likelihood of EAA), and agreed with the Expert Panel opinion in 81% of cases. The MWF EAA Score appeared to perform well when compared with other established case definitions, and when applied to 50 US cases of MWF EAA.

Abstract S1 Table 1 MWF EAA Score for workers with suspected EAA during MWF outbreaks

Respiratory symptoms	
Work-related cough/wheeze/sob/chest tightness	+4
Stopping for breath when walking at own pace on level ground	+6
Previous time off work with any chest illness	+7
Constitutional symptoms	
Recurrent flu-like symptoms worse at the end of the working week	+5
Unexplained weight loss	+7
Physiology	
FVC <80% predicted	+3
FVC <70% predicted or Tlco <80% predicted	+5
Tlco <60% predicted	+10
Radiology/clinical examination	
Abnormal CXR (diffuse ground glass or nodularity)	+6
Abnormal HRCT (ground glass, nodularity, mosaic, or UIP fibrosis)	+7
Fine end-inspiratory crepitations on auscultation	+7
Evidence of inflammation	
Neutrophilia >7 or CRP ≥10	+5
BAL lymphocytosis ≥20%	+8
Lung biopsy typical of EAA (sub-acute EAA or UIP)	+10
Total (max 41)	/41

Conclusions Although difficult to truly validate, the MWF EAA Score offers a weighted and evidence-based case definition for workers suspected of suffering from EAA in MWF outbreaks. If adopted as a standard, the MWF EAA score will facilitate comparison of the findings from future outbreaks, in terms of risk factors and causation of this interesting disease.

S2 OCCUPATIONAL EXPOSURE, BREATHLESSNESS AND COPD IN A GENERAL POPULATION OF OLDER UK MEN

doi:10.1136/thoraxjnl-2011-201054b.2

¹E Pickles, ²J Szram, ¹S J Schofield, ¹A P M Woods, ¹P Cullinan. ¹Imperial College, London, UK; ²Royal Brompton and Harefield NHS Foundation Trust, London, UK

Most COPD is attributable to smoking, however evidence from workforce and general population studies suggest that occupational exposure is also associated with the disease. This study examines the relationship between respiratory symptoms, doctor-diagnosed COPD and occupational exposure in a general UK population. Lifetime occupational and smoking history, doctor-diagnosed disease and current respiratory symptoms were collected by postal questionnaire in a cohort of subjects aged 51–60, recruited from 33 general practices in Kent; here we present the results in men (n=3011). Occupations were defined *a priori* as being associated with an increased risk of COPD using 13 job categories as defined for the European Community Respiratory Health Survey¹; all other occupations were designated low risk (referent). Logistic regression in men with complete data and no history of asthma (n=2452) demonstrated a statistically significant increase in both the odds of breathlessness (modified MRC score 1 or greater, Abstract S2 table 1) and breathlessness reported along with symptoms of chronic bronchitis (data not shown) in subjects who had a history of ever having been employed in a risky job, compared to other referent (mainly white collar) workers. This association, between work in

Abstract S2 Table 1 Logistic regression: relationship between breathlessness in men and ever working in an occupational group with an *a priori* risk of COPD (as defined for the ECRHS) compared to referent occupations

ECRHS occupational groups at risk of COPD	Total (n)	Subjects reporting breathlessness (n, %)	Crude OR (95%CI)	Adjusted* OR (95%CI)	p
Referent occupations	654	84 (12.8)	1.0	1.0	na
Ever working in an at risk job	1798	379 (21.1)	1.81 (1.40 to 2.34)	1.57 (1.21 to 2.04)	0.001
Cleaning	120	38 (31.7)	3.14 (2.00 to 4.92)	2.62 (1.64 to 4.17)	<0.001
Painting	88	26 (29.6)	2.85 (1.70 to 4.75)	2.22 (1.30 to 3.80)	0.003
Agriculture	190	53 (27.9)	2.63 (1.78 to 3.88)	2.31 (1.54 to 3.45)	<0.001
Transport	566	151 (26.7)	2.47 (1.84 to 3.32)	2.07 (1.52 to 2.81)	<0.001
Food	128	34 (26.6)	2.45 (1.56 to 3.87)	1.96 (1.20 to 3.2)	0.007
Healthcare	126	30 (23.8)	2.12 (1.33 to 3.39)	1.96 (1.21 to 3.16)	0.006
Other manual work	307	76 (24.8)	2.23 (1.58 to 3.12)	1.81 (1.26 to 2.60)	0.001
Wood	234	54 (23.1)	2.04 (1.39 to 2.98)	1.81 (1.23 to 2.68)	0.003
Construction	389	92 (23.7)	2.10 (1.52 to 2.92)	1.73 (1.23 to 2.43)	0.002
Mining	378	81 (21.4)	1.85 (1.32 to 2.59)	1.44 (1.01 to 2.04)	0.045
Metal industries	586	113 (19.3)	1.62 (1.19 to 2.20)	1.38 (1.00 to 1.89)	0.048
Electrical	293	51 (17.4)	1.43 (0.98 to 2.09)	1.32 (0.90 to 1.94)	0.159
Chemical	110	15 (13.6)	1.07 (0.59 to 1.93)	0.82 (0.51 to 1.71)	0.822

Crude ORs and *adjusted for age and smoking (by pack year history) are shown.

occupations with an *a priori* risk of COPD and chronic respiratory symptoms, remained after adjustment for age and smoking status, using pack year history. The strength of the relationship between symptoms and work varied by occupational type; cleaners, painters and agricultural workers had the highest risk of breathlessness when compared to the referent population (Abstract S2 table 1). An increased risk of doctor-diagnosed COPD (COPD, chronic bronchitis or emphysema) was also found in cleaners, transport workers, wood and construction workers (data not shown); in comparison with the prevalence of respiratory symptoms, the number of men declaring doctor-diagnosed disease was small (21% vs 5%). This study demonstrates an association between occupational exposure, chronic respiratory symptoms and doctor-diagnosed COPD within a general population of older males in the UK independent of smoking history. Further characterisation of the cohort, using the results of spirometry, will allow the relationship between risky job exposure and disease to be examined in more detail.

REFERENCE

1. European Community Respiratory Health Survey.

S3 WORK PERFORMANCE AND AIRFLOW OBSTRUCTION IN A GENERAL UK POPULATION OF OLDER WORKERS

doi:10.1136/thoraxjnl-2011-201054b.3

¹J Szram, ²S J Schofield, ²A P M Woods, ²N Crisp, ²D Harris, ²P Cullinan. ¹Royal Brompton and Harefield NHS Foundation Trust, London, UK; ²Imperial College, London, UK

Work disability as a consequence of COPD has been found in a number of patient cohorts and respiratory symptoms were shown to be significantly associated with work limitation in the Lungs at Work study.¹ The impact of airflow obstruction on impaired work performance in the general population is unknown. The clinical assessment was designed to examine the relationship between respiratory symptoms and work performance in a general population of older workers in more detail. Volunteers in full time employment at the time of the initial postal questionnaire study (of 51 to 60-year olds through general practice) underwent clinical assessment, including spirometry (n=1773). Results are shown in Abstract S3 table 1. Prevalence of declared doctor-diagnosed COPD was low (1.9% of men and 0.6% of women) compared to that of airflow obstruction on spirometry (post-bronchodilator FEV₁<80% predicted and FEV₁/FVC ratio<0.7) was higher, similar to previous published estimates. Men and women with airflow obstruction had a significantly higher prevalence of poor self-reported performance at work than individuals with normal spirometry. Subjects of both sexes with abnormal spirometry who also reported high levels of physical activity in their current job were significantly more likely to report poor work performance than individuals without airflow obstruction with similarly high activity levels; this difference was not seen in low activity work (data not shown). Men were significantly more likely than women to predict that they would stop work due to ill-health. In both sexes, participants with abnormal spirometry were significantly more likely to predict ill-health retirement than individuals with normal lung function. This study has demonstrated an association between airflow obstruction and both work performance within a general population of older workers; the level of physical activity required at work had an important effect on this relationship. Future loss from the workforce due to ill-health was also related to lung function. Detection of airflow obstruction could aid retention in employment, provided that suitable interventional strategies are in place to support older workers.

Abstract S3 Table 1 Self-reported work performance, physical activity and current work exposures in a general population of adults aged 51–60 in full time employment (n=1773), stratified by sex and airflow obstruction

	Men (n=1101) n, %		Women (n=672) n, %		p Value
Poor work performance	112 (10.4)		54 (8.2)		0.128
High level of physical activity at work	496 (46.1)		227 (34.6)		<0.001
Airflow obstruction*	77 (7.3)		25 (3.9)		0.004
Health will limit ability to work	568 (52.1)		271 (40.7)		<0.001

	Airflow obstruction* n, %			Airflow obstruction* n, %		
	Yes (n=77)	No (n=977)	p Value	Yes (n=25)	No (n=617)	p Value
Poor work performance	13 (16.9)	90 (9.4)	0.035	5 (20.0)	48 (7.9)	0.032
Poor work performance and high level of physical activity at work	11(30.6)	58 (13.4)	0.005	3 (23.1)	15 (7.5)	0.084
Health will limit ability to work	50 (64.9)	495(51.2)	0.020	16 (64.0)	9 (36.0)	0.016

*Post-bronchodilator FEV₁ <80% predicted and FEV₁/FVC ratio <0.7.

REFERENCE

1. Szram J, Schofield SJ, Woods APM, et al. Breathlessness, respiratory disease and work performance in older adults. *British Thoracic Society Conference* 2010. Poster: P4.

S4 OCCUPATIONAL ASTHMA; IS THIS THE CAUSE OF EXCESS RESPIRATORY SYMPTOMS AND COPD DESCRIBED IN BITUMEN EXPOSED WORKERS?

doi:10.1136/thoraxjnl-2011-201054b.4

N Surange, J Hoyle. North Manchester General Hospital, Penine Acute NHS Trust, Manchester, UK

Introduction Epidemiological studies suggest increased risk of asthma and COPD in asphalt exposed workers.¹ Bitumen is used in this industry. In this case we describe occupational asthma caused by bitumen exposure in a lab environment. This is the first such report to our knowledge.

Case History A 49-year old male with no history of asthma or atopy and <5 pack years smoking history presented with airflow obstruction on surveillance spirometry, cough and wheeze. He analysed hot bitumen samples (1900 °C) in a lab environment without using respiratory protective equipment. Spirometry FEV₁ 57%, FVC 85% ratio 55%. Respiratory symptoms improved after 3 weeks off work, returning soon after he rejoined the lab. Peak flow rates were variable and lower at work. Analysis with OASYS scored 3.08 (Abstract S4 figure 1). Histamine challenge test was positive (PC₂₀ 2.216 mg/ml). Skin prick test to paraldehyde was positive. After 8 weeks away from the exposed environment the subject was asymptomatic, continued to have obstructive spirometry but improved bronchial reactivity (PC₂₀ 7.489 mg/ml) without medication and improved peak flows with little diurnal variation. One week after restarting work the respiratory symptoms returned. Repeat OASYS charts scored 3.14 with histamine reactivity similar to baseline (PC₂₀ 2.81 mg/ml) after 4 weeks. A specific challenge test was not possible due to the problems with heating bitumen to 1900 °C in the hospital lab.

Conclusion The progression of symptoms and lung function in relation to work history supports the diagnosis of occupational asthma induced by bitumen fume exposure. This has not been reported previously. The possible mechanisms include sensitisation to short chain aldehydes, produced by partial combustion of