

occupations with an *a priori* risk of COPD and chronic respiratory symptoms, remained after adjustment for age and smoking status, using pack year history. The strength of the relationship between symptoms and work varied by occupational type; cleaners, painters and agricultural workers had the highest risk of breathlessness when compared to the referent population (Abstract S2 table 1). An increased risk of doctor-diagnosed COPD (COPD, chronic bronchitis or emphysema) was also found in cleaners, transport workers, wood and construction workers (data not shown); in comparison with the prevalence of respiratory symptoms, the number of men declaring doctor-diagnosed disease was small (21% vs 5%). This study demonstrates an association between occupational exposure, chronic respiratory symptoms and doctor-diagnosed COPD within a general population of older males in the UK independent of smoking history. Further characterisation of the cohort, using the results of spirometry, will allow the relationship between risky job exposure and disease to be examined in more detail.

## REFERENCE

1. European Community Respiratory Health Survey.

## S3 WORK PERFORMANCE AND AIRFLOW OBSTRUCTION IN A GENERAL UK POPULATION OF OLDER WORKERS

doi:10.1136/thoraxjnl-2011-201054b.3

<sup>1</sup>J Szram, <sup>2</sup>S J Schofield, <sup>2</sup>A P M Woods, <sup>2</sup>N Crisp, <sup>2</sup>D Harris, <sup>2</sup>P Cullinan. <sup>1</sup>Royal Brompton and Harefield NHS Foundation Trust, London, UK; <sup>2</sup>Imperial College, London, UK

Work disability as a consequence of COPD has been found in a number of patient cohorts and respiratory symptoms were shown to be significantly associated with work limitation in the Lungs at Work study.<sup>1</sup> The impact of airflow obstruction on impaired work performance in the general population is unknown. The clinical assessment was designed to examine the relationship between respiratory symptoms and work performance in a general population of older workers in more detail. Volunteers in full time employment at the time of the initial postal questionnaire study (of 51 to 60-year olds through general practice) underwent clinical assessment, including spirometry (n=1773). Results are shown in Abstract S3 table 1. Prevalence of declared doctor-diagnosed COPD was low (1.9% of men and 0.6% of women) compared to that of airflow obstruction on spirometry (post-bronchodilator FEV<sub>1</sub><80% predicted and FEV<sub>1</sub>/FVC ratio<0.7) was higher, similar to previous published estimates. Men and women with airflow obstruction had a significantly higher prevalence of poor self-reported performance at work than individuals with normal spirometry. Subjects of both sexes with abnormal spirometry who also reported high levels of physical activity in their current job were significantly more likely to report poor work performance than individuals without airflow obstruction with similarly high activity levels; this difference was not seen in low activity work (data not shown). Men were significantly more likely than women to predict that they would stop work due to ill-health. In both sexes, participants with abnormal spirometry were significantly more likely to predict ill-health retirement than individuals with normal lung function. This study has demonstrated an association between airflow obstruction and both work performance within a general population of older workers; the level of physical activity required at work had an important effect on this relationship. Future loss from the workforce due to ill-health was also related to lung function. Detection of airflow obstruction could aid retention in employment, provided that suitable interventional strategies are in place to support older workers.

**Abstract S3 Table 1** Self-reported work performance, physical activity and current work exposures in a general population of adults aged 51–60 in full time employment (n=1773), stratified by sex and airflow obstruction

	Men (n=1101)		Women (n=672)		p Value
	n, %		n, %		
Poor work performance	112 (10.4)		54 (8.2)		0.128
High level of physical activity at work	496 (46.1)		227 (34.6)		<0.001
Airflow obstruction*	77 (7.3)		25 (3.9)		0.004
Health will limit ability to work	568 (52.1)		271 (40.7)		<0.001

	Airflow obstruction* n, %			Airflow obstruction* n, %		
	Yes	No		Yes	No	
	(n=77)	(n=977)	p Value	(n=25)	(n=617)	p Value
Poor work performance	13 (16.9)	90 (9.4)	0.035	5 (20.0)	48 (7.9)	0.032
Poor work performance and high level of physical activity at work	11(30.6)	58 (13.4)	0.005	3 (23.1)	15 (7.5)	0.084
Health will limit ability to work	50 (64.9)	495(51.2)	0.020	16 (64.0)	9 (36.0)	0.016

\*Post-bronchodilator FEV<sub>1</sub> <80% predicted and FEV<sub>1</sub>/FVC ratio <0.7.

## REFERENCE

1. Szram J, Schofield SJ, Woods APM, et al. Breathlessness, respiratory disease and work performance in older adults. *British Thoracic Society Conference* 2010. Poster: P4.

## S4 OCCUPATIONAL ASTHMA; IS THIS THE CAUSE OF EXCESS RESPIRATORY SYMPTOMS AND COPD DESCRIBED IN BITUMEN EXPOSED WORKERS?

doi:10.1136/thoraxjnl-2011-201054b.4

N Surange, J Hoyle. North Manchester General Hospital, Penine Acute NHS Trust, Manchester, UK

**Introduction** Epidemiological studies suggest increased risk of asthma and COPD in asphalt exposed workers.<sup>1</sup> Bitumen is used in this industry. In this case we describe occupational asthma caused by bitumen exposure in a lab environment. This is the first such report to our knowledge.

**Case History** A 49-year old male with no history of asthma or atopy and <5 pack years smoking history presented with airflow obstruction on surveillance spirometry, cough and wheeze. He analysed hot bitumen samples (1900 °C) in a lab environment without using respiratory protective equipment. Spirometry FEV<sub>1</sub> 57%, FVC 85% ratio 55%. Respiratory symptoms improved after 3 weeks off work, returning soon after he rejoined the lab. Peak flow rates were variable and lower at work. Analysis with OASYS scored 3.08 (Abstract S4 figure 1). Histamine challenge test was positive (PC<sub>20</sub> 2.216 mg/ml). Skin prick test to paraldehyde was positive. After 8 weeks away from the exposed environment the subject was asymptomatic, continued to have obstructive spirometry but improved bronchial reactivity (PC<sub>20</sub> 7.489 mg/ml) without medication and improved peak flows with little diurnal variation. One week after restarting work the respiratory symptoms returned. Repeat OASYS charts scored 3.14 with histamine reactivity similar to baseline (PC<sub>20</sub> 2.81 mg/ml) after 4 weeks. A specific challenge test was not possible due to the problems with heating bitumen to 1900°C in the hospital lab.

**Conclusion** The progression of symptoms and lung function in relation to work history supports the diagnosis of occupational asthma induced by bitumen fume exposure. This has not been reported previously. The possible mechanisms include sensitisation to short chain aldehydes, produced by partial combustion of