

The example of collateral ventilation shown in figure 2 of our paper² is intersegmental, however, ³He MRI is able to visualise both interlobar and intralobar collateral and delayed ventilation. The technique images gas as it moves within the lung whether it traverses lobar or segmental boundaries.

Interlobar lung fissure condition and emphysema type are anatomical radiological markers, and while structure is important there are risks to predicting function from form alone. The assessment of collateral and delayed ventilation pathways with hyperpolarised gas MRI is intrinsically functional because we directly visualise the gas itself.

For further information about a region being considered for bronchoscopic lung volume reduction, time-resolved ³He MRI could be used in conjunction with ¹H blood perfusion MRI³ to assess ventilation–perfusion matching, and with ³He diffusion MRI⁴ to measure emphysematous destruction at the acinar level.

More research is needed to determine if hyperpolarised gas MRI reliably correlates with catheter measurements in treatment planning, and we look forward to exploring such studies in the future, and welcome this correspondence.

Helen Marshall,¹ Rod Lawson,² Jim M Wild¹

¹Department of Academic Radiology, University of Sheffield, Sheffield, South Yorkshire, UK; ²Respiratory Medicine, Sheffield Teaching Hospitals NHS Trust, South Yorkshire, UK

Correspondence to Professor Jim M Wild, Department of Academic Radiology, University of Sheffield, C Floor, Royal Hallamshire Hospital, Glossop Road, Sheffield, S10 2JF, UK; j.m.wild@sheffield.ac.uk

Competing interests None.

Ethics approval Ethics approval was provided by South Yorkshire REC (REC reference: 10/H1310/11).

Provenance and peer review Not commissioned; internally peer reviewed.

Accepted 24 July 2012

Published Online First 15 August 2012

Thorax 2012;**67**:1111.

doi:10.1136/thoraxjnl-2012-202455

REFERENCES

1. **Anile M**, Diso D, Venuta F. Assessment of intraparenchymal lung collateral ventilation. *Thorax* 2012;**67**:1111.
2. **Marshall H**, Deppe MH, Parra-Robles J, *et al*. Direct visualisation of collateral ventilation in COPD with hyperpolarised gas MRI. *Thorax* 2012;**67**:613–17.
3. **Ohno Y**, Hatabu H, Murase K, *et al*. Quantitative assessment of regional pulmonary perfusion in the entire lung using three-dimensional ultrafast dynamic contrast-enhanced magnetic resonance imaging: preliminary experience in 40 subjects. *J Magn Reson Imaging* 2004;**20**:353–65.
4. **Saam BT**, Yablonskiy DA, Kodibagkar VD, *et al*. MR imaging of diffusion of (³)He gas in healthy and diseased lungs. *Magn Reson Med* 2000;**44**:174–9.

Authors' response to assessment of intraparenchymal lung collateral ventilation

We thank the writers of the letter¹ for their interest and agree that the potential placement of this novel imaging method in the planning of patient treatment, such as bronchoscopic and surgical lung volume reduction, is not yet determined.