ORIGINAL ARTICLE

Glucocorticoid receptor β and histone deacetylase 1 and 2 expression in the airways of severe asthma

Claire A Butler,¹ Stephen McQuaid,² Cliff C Taggart,¹ Sinead Weldon,¹ Richard Carter,³ Grzegorz Skibinski,¹ Timothy J Warke,¹ David F Choy,⁴ Lorcan P McGarvey,¹ Peter Bradding,³ Joe R Arron,⁴ Liam G Heaney¹

► An additional material is published online only. To view this file please visit the journal online (<http://thorax.bmj.com/content/67/5.toc>).

¹Centre for Infection and Immunity, Health Sciences Building, Queens University Belfast, Belfast, UK

²Tissue Pathology, Belfast Health and Social Care Trust, Belfast, UK

³Department of Infection, Immunity and Inflammation, Maurice Shock Medical Sciences Building, Leicester, UK

⁴ITGR Biomarker Discovery, Genentech, Inc., South San Francisco, California, USA

Correspondence to

Dr Liam Heaney, Centre for Infection and Immunity, Queens University Belfast, Level 8, Belfast City Hospital, Lisburn Road, Belfast BT9 7AB, Northern Ireland, UK; l.heaney@qub.ac.uk

Received 7 July 2011

Accepted 11 November 2011

Published Online First

8 December 2011

ABSTRACT

Rationale Upregulation of glucocorticoid receptor β (GR β) has been implicated in steroid resistance in severe asthma, although previous studies are conflicting. GR β has been proposed as a dominant negative isoform of glucocorticoid receptor α (GR α) but it has also been suggested that GR β can cause steroid resistance via reduced expression of histone deacetylase 2 (HDAC2), a key regulator of steroid responsiveness in the airway.

Objectives To examine GR β , GR α , HDAC1 and HDAC2 expression at transcript and protein levels in bronchial biopsies from a large series of patients with severe asthma, and to compare the findings with those of patients with mild to moderate asthma and healthy volunteers.

Methods Bronchoscopic study in two UK centres with real-time PCR and immunohistochemistry performed on biopsies, western blotting of bronchial epithelial cells and immunoprecipitation with anti-GR β antibody.

Measurements and main results Protein and mRNA expression for GR α and HDAC2 did not differ between groups. GR β mRNA was detected in only 13 of 73 samples (seven patients with severe asthma), however immunohistochemistry showed widespread epithelial staining in all groups. Western blotting of bronchial epithelial cells with GR β antibody detected an additional 'cross-reacting' protein, identified as clathrin. HDAC1 expression was increased in patients with severe asthma compared with healthy volunteers.

Conclusions GR β mRNA is expressed at low levels in a minority of patients with severe asthma. HDAC1 and HDAC2 expression was not downregulated in severe asthma. These data do not support upregulated GR β and resultant reduced HDAC expression as the principal mechanism of steroid resistance in severe asthma. Conflicting GR β literature may be explained in part by clathrin cross-reactivity with commercial antibodies.

INTRODUCTION

Patients with severe asthma respond poorly to available therapies, which results in high healthcare costs due to unscheduled healthcare visits, treatment of co-morbidities and work absence.¹ The cornerstone of asthma treatment is glucocorticoid therapy; however patients with severe asthma are relatively resistant to inhaled glucocorticoids, often resulting in a requirement for systemic steroids to maintain disease control.²

Proposed molecular mechanisms for glucocorticoid resistance include abnormalities of glucocorticoid

Key messages

What is the key question?

► Increased glucocorticoid receptor β (GR β) with resultant reduction in histone deacetylase 2 (HDAC2) expression has been implicated in the pathogenesis of severe steroid-resistant asthma, but previous data are conflicting.

What is the bottom line?

► GR β mRNA was expressed at very low levels in the airways of a minority of patients with severe asthma. GR β immunostaining was prominent in bronchial biopsies from all patients, but this was subsequently shown to be due to antibody cross-reactivity with heavy chain clathrin. No reduction in HDAC2 expression was identified in severe asthma.

Why read on?

► These data do not support altered GR β /HDAC expression as the principal underlying mechanism of steroid resistance in severe asthma. Some of the discrepancies in the GR β literature may be explained by clathrin cross-reactivity with commercial antibodies.

binding,^{2–4} reduced glucocorticoid receptor nuclear translocation in response to steroids,^{5,6} increased expression of pro-inflammatory transcription factors (eg, activator protein-1 and nuclear factor- κ B)^{7,8} and altered expression of co-repressor proteins (eg, histone deacetylases).⁹ Two commonly proposed mechanisms for steroid resistance in severe asthma are increased expression of the dominant negative isoform of the glucocorticoid receptor, glucocorticoid receptor β (GR β), and reduced expression and activity of histone deacetylase 2 (HDAC2).^{3,5} A recent study suggested the reduced HDAC2 in severe asthma may be regulated by GR β overexpression.¹⁰

Existing data on GR β expression in severe asthma have been variable. GR β -immunoreactive peripheral blood mononuclear cells (PBMCs), CD3 T cells and bronchoalveolar lavage cells were increased in steroid-resistant asthma.^{3,11,12} Furthermore, Sousa *et al* reported increased GR β -positive cells in steroid-resistant asthmatic tuberculin responses.¹³ In the lungs of patients who have died from asthma increased GR β immunoreactivity has been reported,¹⁴ and in the only published

bronchial biopsy study, GR β expression detected by immunohistochemistry was increased in the epithelium and submucosa of patients with severe asthma.¹⁵ However, other studies in PBMCs in severe asthma have been negative.^{16–17} Thus, it remains unclear if GR β is elevated in the airways of severe asthma (and other 'steroid resistant' conditions) and why there is such variability in the published literature.

Our aims were to examine GR β and GR α expression at transcript and protein levels in bronchial biopsies from a large series of patients with severe asthma, and to compare the findings with those for patients with mild to moderate asthma and healthy volunteers. We also studied HDAC1 and HDAC2 expression in matched biopsies.

METHODS

Subjects

This study was performed in two UK centres, Queens University Belfast and University of Leicester. Patients with severe asthma were recruited from the Belfast City Hospital and Glenfield Hospital, Leicester Difficult Asthma Services, where systematic evaluation protocols ensure patients have well characterised severe asthma as defined by current American Thoracic Society guidelines.¹⁸ All patients had persisting symptoms despite treatment at step 4 or 5 of the Global Initiative for Asthma (GINA) guidelines.¹⁹ Patients with mild to moderate asthma and healthy volunteers were recruited from hospital clinics or by advertisement.

In total, 27 participants were recruited from Belfast (10 healthy volunteers (HV), 7 patients with mild to moderate asthma (MA) and 10 patients with severe asthma (SA)) and 56 participants from Leicester (12 HV, 23 MA and 21 SA) (table 1).

Bronchoscopy

Bronchoscopy was performed using standard techniques (see online supplement). Biopsies were either placed immediately in RNA preservative (RNAlater, Ambion, Austin, Texas, USA) and submitted for real-time PCR (qPCR) or placed into dry acetone containing protease inhibitors (iodoacetamide and phenylmethylsulfonyl fluoride) and fixed overnight at -20°C for immunohistochemistry. Brushings were placed in Promocell medium containing penicillin/streptomycin and Primocin for cell culture (Promocell, Heidelberg, Germany).

Gene expression analysis

qPCR analysis was performed using RNA from homogenised bronchial biopsies (74 biopsies; 22 HV, 28 MA and 24 SA) using TaqMan $^{\circledR}$ Gene Expression Assays (Applied Biosystems, Foster City, California, USA). RNA purification and qPCR quality control are detailed in the online supplement. Relative expression levels were determined by the $2(-\Delta\Delta C_T)$ method, as described in Applied Biosystems User Bulletin No. 2 (P/N 4303859).²⁰

Bronchial biopsy immunohistochemistry and confocal microscopy

For immunohistochemistry, biopsy samples were processed and stained using local protocols. Fifty-five biopsies (17 HV, 21 MA and 17 SA) were processed for immunohistochemistry. Infiltration and embedding with JB4 resin was performed.²¹ The immunohistochemical technique and confocal analysis are detailed in the online supplement.

Immunostained sections were evaluated by two independent assessors with consensus scoring for any conflicting results (scoring system given in the online supplement). Two polyclonal anti-GR β antibodies were compared, ab3581 (Abcam,

Cambridge, UK) and PA3-514 (Affinity Bioreagents, Golden, CO, USA via Pierce Biotechnology, Rockford, IL, USA) along with anti-GR α (sc-1002), HDAC1 (sc-7872) and HDAC2 (sc-7899) (Insight Biotechnology, Wembley, UK).

Epithelial cell culture and western blotting

Bronchial epithelial brushings (Belfast cohort) were cultured (Promocell) and lysates (NE-PER lysis kit, Pierce Biotechnology, MSC, Dublin) used for western blot experiments (see online supplement). For clathrin identification blots, anti-clathrin antibody (ab2731, Abcam) was used. For immunoprecipitation, BEAS-2B cells (ATCC, Manassas, VA, USA) were also used following culture in supplemented Dulbecco's Modified Eagle's Medium (Gibco-Invitrogen Ltd, Paisley, UK).

Immunocytochemistry

Primary bronchial epithelial cells (PBECs) were grown onto collagen-coated Thermanox coverslips (catalogue no. 174950, Nalge Nunc International, Rochester, NY, USA), permeabilised and stained (see online supplement), and mounted on slides.

Immunoprecipitation

Immunoprecipitation was performed using IgA Magnetic Beads (Pierce Biotechnology, Rockford, IL, USA). PBECs and BEAS-2B cells were used. The method is detailed in the online supplement and the resultant blots were probed with anti-GR β and anti-clathrin (ab2731, Abcam) antibodies. In addition, increasing amounts (250 ng, 500 ng, 1000 ng and 2500 mg) of recombinant clathrin was run on a gel and the blot probed with anti-GR β and anti-clathrin antibodies.

Proteomics

Protein identification was performed by the Proteomics Facility, University of Dundee, UK. Following immunoprecipitation, gels were stained with Coomassie blue and the higher molecular weight band cut out of the gel, digested with trypsin prior to analysis by one-dimensional nano-scale liquid chromatography tandem mass spectrometry (nLC-MS-MS) and protein identification performed with Mascot (www.matrixscience.com/).

Statistical analysis

The statistical package Prism (GraphPad Prism version 4.00 for Windows, GraphPad Software, San Diego, California, USA) was used to generate graphs and perform appropriate statistical analysis. Data are presented as median (IQR) unless otherwise stated. Between-group data were analysed using one-way analysis of variance (Kruskal–Wallis) with Dunn's post hoc test or Wilcoxon Signed Rank test as appropriate. A p value <0.05 was considered significant.

RESULTS

GR β , GR α , HDAC1 and HDAC2 mRNA expression

In the samples available for qPCR ($n=74$), GR β mRNA was detectable in only 13 of the 74 (18%) participants (seven SA—four of whom were on oral steroids, four with moderate persistent asthma on inhaled steroids, one with steroid-naïve asthma and one HV) (figure 1A). There was a similar rate of detection in the SA group ($p=0.10$, Fisher's exact test). Despite the use of different specific primer-probe sets for GR β , GR β mRNA was not detected in any of the other samples. GR α mRNA was detectable in 70 of the biopsy samples (figure 1B, median (IQR) relative fold expression of GR α mRNA expression was 343.7 (137.4–928.2), 498.4 (200.4–1439) and 463.1 (186.3–1349) in HV, MA and SA groups respectively, $p=0.62$,

Table 1 Demographic characteristics of patients with asthma and healthy volunteers (Belfast and Leicester cohorts combined)

	Healthy volunteers (n=22)	Patients with mild to moderate asthma (n=30)	Patients with severe asthma (n=31)	p Value*
Age (years), median (IQR)	30 (22–45)	29 (24–39)	48 (38–54)†	<0.0001
Sex, men/women (n)	10/12	14/16	22/9	0.09
Duration of asthma (years), mean±SD	N/A	12±10	22±15‡	0.007
Atopy (n)	11	21	23	0.16
FEV ₁ (litre/min), mean±SD	3.67±0.78	3.32±0.77	2.48±0.67§	<0.0001
FEV ₁ (% predicted), mean±SD	102.8±11.9	93.5±16.0	75.9±16.9¶	<0.0001
FEV ₁ /FVC, mean±SD	82.1±9.8	75.9±10.6	64.6±11.1**	<0.0001
Inhaled steroid (n)	0	19	31	<0.0001
BDP equivalent (µg), median (IQR)	0	400 (0–800)	1600 (1200–1600)††	
Maintenance systemic steroids (n)	0	0	15 (median dose prednisolone 5 mg (0–10 mg) plus 1 subject maintenance intramuscular triamcinolone)	
Theophylline (n)	0	1	17	<0.0001
Long-acting β ₂ -agonists (n)	0	14	31	<0.0001
Prior smoking history (pack years) (n, median range)	1 (4)	5 (1, 1–10)	13 (2.5, 1–20)	0.004
Eosinophils (10 ⁹ /litre), median (IQR)	0.1 (0.07–0.21)	0.26 (0.14–0.45)‡‡	0.49 (0.25–0.74)§§	<0.0001
IgE (kU/litre), geometric mean (95% confidence interval)	24.6 (13.3 to 45.8)	159.8 (89.8 to 284.6)¶¶	238.5 (136.2 to 417.5)***	<0.0001

Comparison with the healthy volunteer group: †p<0.001, §p<0.001, ¶p<0.001, **p<0.001, §§p<0.001, ‡‡p<0.05, ¶¶p<0.001. Comparison with patients with mild to moderate asthma: ‡p<0.001, ‡‡p=0.007, §p<0.001, ¶p<0.001, **p<0.001, ††p<0.0001, §§p<0.05, ***p<0.001.

*Calculated by analysis of variance (ANOVA) with Bonferroni's multiple comparisons test (parametric) or Kruskal–Wallis with Dunn's multiple comparison test (non-parametric) or t-test when only asthma groups compared or χ^2 for categorical variables (sex, atopy, theophylline, long-acting β₂ agonists, smoking history).

BDP, beclomethasone dipropionate; FEV₁, forced expiratory volume in 1 s; FVC, forced vital capacity.

Kruskal–Wallis). HDAC1 mRNA was significantly upregulated in the SA group compared with the HV group (figure 1C, HDAC1—median (IQR) relative fold expression was 0.08 (0.04, 0.21), 0.22 (0.09, 0.49) and 0.27 (0.10, 0.94) in the HV, MA and SA groups respectively (p=0.02, Kruskal–Wallis, post hoc analysis p<0.05 between HV and SA). HDAC2 expression did not

differ between participant subgroups (figure 1D, HDAC2—median (IQR) relative fold expression was 0.52 (0.14–0.97), 0.63 (0.33–4.29) and 0.76 (0.36–1.97) in the HV, MA and SA groups respectively (p=0.23, Kruskal–Wallis). There was a significant correlation between HDAC2 mRNA and GRα mRNA (figure 3D, Spearman r=0.75, p<0.001).

Figure 1 Gene expression in bronchial biopsies (n=74). Values represent fold change in target gene expression with GAPDH as housekeeping gene ($2^{-\Delta\Delta Ct}$). (A) Glucocorticoid receptor β (GRβ) mRNA was detectable in only 13 of 74 samples (seven patients with oral-steroid-dependent asthma, four with moderate persistent asthma on inhaled steroids, one with steroid-naïve asthma and one healthy volunteer). (B) GRα mRNA widely expressed in bronchial biopsies. There were no significant between-group differences (p=0.62, Kruskal–Wallis). (C) Histone deacetylase 1 (HDAC1) mRNA expression in bronchial biopsies. HDAC1 was increased in patients with severe asthma compared with healthy volunteers (HDAC1 p=0.02, Kruskal–Wallis, post hoc analysis p<0.05 between healthy volunteers and severe asthma). (D) HDAC2 mRNA expression in bronchial biopsies. There were no significant between-group differences (HDAC2 p=0.23, Kruskal–Wallis).

Immunohistochemistry for GR β , GR α , HDAC1 and HDAC2 and relation to asthma severity in bronchial biopsies

Similar location and intensity of immunostaining for GR β was seen using both immunohistochemistry protocols (figure 2A). Identical staining was noted with both GR β antibodies (data not shown). The most striking observation was the intensity of immunostaining in contrast to low mRNA expression. GR β immunostaining was mainly localised to the epithelium with scant submucosal expression in glands, smooth muscle cells and inflammatory cells (figure 2A,B). Confocal microscopy revealed that epithelial immunostaining with GR β antibodies was predominantly cytoplasmic, though some cells did demonstrate nuclear staining (figure 2D).

There was no significant between-group difference in epithelial GR β immunostaining (figure 2C, $p=0.18$, Kruskal–Wallis). Figure 2B shows representative sections from each of the participant groups. Among the 10 patients with mRNA above the limit of detection and matched immunohistochemistry samples, there was no correlation between mRNA levels and epithelial expression of GR β (data not shown).

Glucocorticoid receptor α (GR α) immunostaining was diffusely present in the epithelium and submucosa with no difference between participant subgroups (data not shown), but GR α immunostaining was more abundant than GR β immunostaining in all groups.

HDAC1 and HDAC2 immunostaining was also more prominent in the epithelium, with occasional submucosal cells. There was no difference between subgroups (figure 3A–C; HDAC1, $p=0.55$, Kruskal–Wallis and HDAC2, $p=0.89$, Kruskal–Wallis). In the SA group, there was no significant difference in mRNA or protein expression for HDAC1 and HDAC2 between those taking theophyllines and those who were not (data not shown).

Western blot analysis and immunocytochemical analysis of PBEC lysates

Immunocytochemical analysis of cultured PBECs revealed predominantly cytoplasmic location of GR β with some cells

demonstrating concomitant nuclear expression (figure 4A). In view of the ‘dissociation’ between mRNA levels of GR β and immunohistochemical staining, further analysis of GR β -antibody specificity was performed using western blots of cultured PBECs. This consistently demonstrated two bands (figure 4B); one band was identified at the anticipated molecular weight of GR β at 90 kDa but a second higher molecular weight band (~ 180 kDa, band 2) was consistently present on the PBEC blots and always stronger than the GR β band in both nuclear and cytoplasmic compartments (figure 5, online supplement).

To evaluate the 180 kDa band, immunoprecipitation using both GR β antibodies was performed using BEAS-2B cells (where the 180 kDa band was also identified) and PBEC lysates. The higher molecular weight bands were cut out of the gel and sequenced. In both BEAS-2B cells and PBECs, this identified heavy chain clathrin as the 180 kDa protein.

To further confirm that clathrin was responsible for band 2, recombinant clathrin was run on a gel and probed separately with anti-GR β and anti-clathrin antibodies (figure 4C). Both antibodies identified the clathrin protein, confirming this band was clathrin and importantly both anti-GR β antibodies displayed similar bands on western blotting. There was no significant difference in clathrin mRNA expression in bronchial biopsies between groups (figure 4D).

DISCUSSION

This study examined GR β , HDAC1 and HDAC2 gene expression and protein in the largest cohort of patients with severe asthma to date. We demonstrated low levels of GR β mRNA expression in bronchial biopsies from a minority of patients with severe asthma despite prominent immunostaining in matched tissue sections. GR β is upregulated at the transcriptional level by steroid exposure in an airway epithelial cell line²² and of the 13 participants in the present study in whom GR β mRNA was detectable, 11 were on inhaled steroids; however, in the majority of patients with asthma on steroid therapy, GR β was not

Figure 2 Glucocorticoid receptor β (GR β) expression in bronchial biopsies; representative immunohistochemistry shown at $\times 400$ magnification. (A) Biopsy from patients with severe asthma, demonstrating the pattern and anatomical localisation of GR β expression to be identical in both clinical centres. (B) GR β expression in individual patient groups. (C) Quantitative immunohistochemical epithelial expression for GR β in bronchial biopsies. There was no significant difference between groups ($p=0.18$, Kruskal–Wallis). (D) Immunostaining with GR β antibody (ab3581) using confocal microscopy in bronchial biopsy from a patient with severe asthma, demonstrating diffuse epithelial staining with scattered submucosal reactivity ($\times 400$ magnification).

Figure 3 Immunohistochemical epithelial expression for histone deacetylase 1 (HDAC1) and HDAC2 in bronchial biopsies. (A) HDAC1: there was no significant difference between groups ($p=0.55$, Kruskal–Wallis); (B) HDAC2: there was no significant difference between groups ($p=0.89$, Kruskal–Wallis). (C) HDAC2 expression in individual patient groups—representative immunohistochemistry shown at $\times 400$ magnification. (D) Highly significant correlation between glucocorticoid receptor α (GR α) and HDAC2 mRNA.

detectable. No previous study has reported the existence of a 180 kDa band on western blot with commercially available GR β antibodies, which we have subsequently identified as heavy chain clathrin. Therefore it is impossible to comment with confidence on the distribution of GR β using these antibodies.

Previous data suggested predominantly nuclear expression of GR β , though cytoplasmic expression has been reported.²³ Clathrin is a ubiquitous protein complex involved in transport vesicle formation to mediate endocytosis of transmembrane receptors,²⁴ and given the low GR β mRNA expression in a few patients, we believe that cross-reactivity with clathrin resulted in prominent biopsy staining. The only other bronchial biopsy study in severe asthma reported significant upregulation of GR β in bronchial biopsies from patients with severe asthma.¹⁵ Their pattern of staining was similar to that reported here but tissue expression of GR β at the mRNA level was not examined. Other studies examining GR β expression in severe asthma using these antibodies have reported variable associations. GR β immunoreactivity was increased in PBMCs and bronchoalveolar lavage (BAL) cells from patients with severe asthma.^{3 12 25} However, another PBMC study was unable to identify any GR β immunoreactivity in healthy volunteers and patients with asthma that is steroid sensitive or resistant.¹⁶ Of note, that study was unable to identify GR β mRNA in the first round of PCR, though it did on the second cycle of nested PCR.¹⁶ A further PBMC study by Torrego *et al* reported 600-fold lower GR β mRNA expression than GR α , but that group was also unable to detect any GR β expression on western blot.¹⁷ Goleva *et al* demonstrated readily appreciable GR β expression using immunohistochemistry in BAL macrophages but demonstrated only femtograms of GR β mRNA.¹²

In other inflammatory disease states, such as glaucoma, ulcerative colitis and rheumatoid arthritis, GR β expression has

been reported to be upregulated in glucocorticoid-unresponsive disease, although again there is controversy.^{26–30} Some of these studies have looked at mRNA expression from PBMCs rather than immunohistochemistry. For example, GR β mRNA was detectable in PBMCs from patients with glucocorticoid-resistant ulcerative colitis compared with low detection in glucocorticoid-responsive disease and healthy volunteers.²⁷ This same group also demonstrated elevated PBMC GR β mRNA in patients with active ulcerative colitis compared with inactive disease or Crohn's disease, though this was disputed by Hausmann *et al*, who did not find elevated PBMC GR β mRNA in relation to steroid resistance.^{26 29} PBMC GR β mRNA and GR β -positive PBMCs were increased in steroid-resistant rheumatoid arthritis.²⁸ Zhang *et al* were able to confirm that glaucomatous trabecular meshwork cells expressed lower levels of GR β protein by western blotting and these cells were more sensitive to the effects of steroids compared with normal trabecular meshwork cells.³⁰ A recent immunohistochemistry study by Ishida *et al* reported increased GR β expression in patients with severe allergic rhinitis compared with healthy controls, and of note, GR β mRNA was not assessed in this study.³¹

Our data suggest that tissue immunohistochemistry with these GR β antibodies is unreliable due to concomitant clathrin staining. This illustrates the problems associated with using antibodies for immunohistochemistry in the absence of validation with other methods, for example, mRNA expression and target protein specificity using western blots. If multiple bands are seen on western blot from the target tissue, we would suggest that this antibody may be inappropriate for use in immunofluorescence, immunocytochemistry or immunohistochemistry. These techniques have been widely used in the GR β literature and this study questions the validity of previous immunohistochemical and immunocytochemical data utilising

Figure 4 (A) Immunocytochemical analysis of primary bronchial epithelial cells from a healthy volunteer showing predominantly cytoplasmic glucocorticoid receptor β (GR β) expression with occasional cells demonstrating nuclear expression (arrowed). (B) Typical western blot showing GR β band (at around 90 kDa) and higher molecular weight band (band 2) at 180 kDa in primary bronchial epithelial cells. (C) Recombinate clathrin (250–2500 ng) was immunoblotted with (i) anti-clathrin antibody (1:1000 in 5% milk) and (ii) anti-GR β antibody (1:500 in 3% bovine serum albumin (BSA)). Similar bands were noted in the anti-GR β and anti-clathrin probed gels at 180 kDa. (D) Gene expression for clathrin in bronchial biopsies. There was no significant difference between patient groups noted.

these antibodies and supports the need for western blotting techniques. In view of the low mRNA signal compared with that for GR α , we would further conclude that GR β is not highly expressed in the airways of patients with severe asthma. Given the substantially higher expression of GR α , GR β is unlikely to function as a dominant-negative isoform of GR α and this mechanism has little to do with the relative steroid resistance seen in this group.

Another proposed mechanism for the induction of steroid resistance by GR β is reduced tissue expression of HDAC2 mRNA and recent data suggested reduced HDAC2 in severe asthma may be regulated by GR β overexpression. If this mechanism of HDAC2 regulation is correct, then our data provide no support that it is relevant in severe asthma because, consistent with the low levels of GR β expression in severe asthma biopsies, we identified no reduction in HDAC2 expression in matched biopsies from the same patients when compared with normal controls and patients with mild to moderate asthma. Reduced HDAC2 tissue expression has been described in patients with chronic obstructive pulmonary disease (COPD) and those who smoke and is associated with reduced HDAC function,^{32–33} but the data in asthmatic airway samples are conflicting. One study suggested decreased expression of HDAC1 and HDAC2 in mild asthma which increased with steroid treatment.³⁴ In more severe asthma, Bergeron *et al* reported similar HDAC2 expression between patients with asthma that is steroid resistant and steroid responsive,¹⁵ whereas another study reported reduced HDAC2 mRNA expression in BAL cells.¹⁰ Importantly, in previous studies in which HDAC mRNA data have been

presented along with functional HDAC activity, when there has been a reduction in HDAC activity, there has been a concomitant reduction in HDAC mRNA.³⁴ Thus, there has been no suggestion that reduced HDAC function in asthma or COPD disease is due to any specific alteration in HDAC function, but rather reduced activity is due to reduced tissue expression. Therefore, we believe it is reasonable to conclude that an absence of reduction of HDAC1 and HDAC2 in this large cohort of patient with refractory asthma argues against this being the mechanism of steroid resistance in this patient group.

At a transcriptional level, we found HDAC1 was increased in patients with severe asthma on higher dose steroid treatment compared with healthy volunteers but not those with mild to moderate asthma, which is consistent with the previously described steroid effect of upregulating HDAC1.³⁵ However, this difference appeared small compared with milder disease and no difference was identified in HDAC1 protein expression. Thus, even if HDAC1 is upregulated by steroid treatment, the fact that many patients with severe asthma remain dependent on oral steroids suggests that this upregulation does not enhance steroid efficacy in this group. There was no association between HDAC2 and theophylline treatment, suggesting that the normal HDAC2 levels in severe asthma are not due to theophylline treatment, as has previously been suggested.^{36–37} The close correlation of HDAC2 with GR α expression at the transcriptional level suggests that these molecules are co-regulated, with increased expression of both molecules tending to increase steroid responsiveness. We did not specifically examine bronchoalveolar macrophages in the present study and thus

a reduction in HDAC2 activity in macrophages contributing to steroid resistance cannot be completely excluded. However, other important and relevant immunological mechanisms in asthma pathogenesis have been identified in proximal airway samples obtained at bronchoscopy.

In conclusion, we have demonstrated that commercially available GR β antibodies cross-react with clathrin and are unreliable for the analysis of GR β expression in situ. GR β mRNA was expressed at low levels in the airways with no difference between steroid-sensitive and severe disease. Our findings may explain the discrepant reports for the role of GR β in steroid-resistant states in a variety of diseases, and do not support the view that GR β plays a role in severe asthma. In addition, our data suggest that downregulation of HDAC1 or HDAC2 does not occur in severe asthma, and so the mechanism of relative steroid resistance in severe asthma remains unclear.

Funding CB was supported by a grant from the Northern Ireland Chest, Heart and Stroke Association (reference no. 200645). Research in Leicester was conducted in laboratories and was part funded by ERDF #05567 and supported by a grant-in-aid from Genentech Inc., South San Francisco, California, USA.

Correction notice This article has been corrected since it was published Online First. The corresponding author is affiliated to Centre for Infection and Immunity, Health Sciences Building, Queens University Belfast, Belfast, UK only.

Competing interests None.

Contributors CB and SMCQ performed immunostaining in the Belfast cohort, analysed the data and contributed to the writing of the manuscript. DC and JA performed and analysed the gene expression data, analysed the data and contributed to the writing of the manuscript. CB and LH recruited subjects, performed bronchoscopy and acquired the demographic data for the Belfast cohort and PB recruited subjects, performed bronchoscopy and acquired the demographic data for the Leicester cohort. TJW and LMcG assisted with bronchoscopy in the Belfast cohort. RC undertook immunohistochemistry for the Leicester cohort and assisted with data analysis under the supervision of PB. CT and SW performed the clathrin western blots and immunostaining. LH and GS conceived and designed the research, contributed to data analysis and interpretation, drafted the original manuscript and supervised CB. All authors approved the final manuscript.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- Masoli M, Fabian D, Holt ST, *et al.* The global burden of asthma: executive summary of the GINA Dissemination Committee report. *Allergy* 2004;**59**:469–78.
- Sher ER, Leung DY, Surs W, *et al.* Steroid-resistant asthma: cellular mechanisms contributing to inadequate response to glucocorticoid therapy. *J Clin Invest* 1994;**93**:33–9.
- Hamid QA, Wenzel SE, Hauk PJ, *et al.* Increased glucocorticoid receptor beta in airway cells of glucocorticoid-insensitive asthma. *Am J Respir Crit Care Med* 1999;**159**:1600–4.
- Nimmagadda SR, Szefer SJ, Spahn JD, *et al.* Allergen exposure decreases glucocorticoid receptor binding affinity and steroid responsiveness in atopic asthmatics. *Am J Respir Crit Care Med* 1997;**155**:87–93.
- Ito K, Chung KF, Adcock IM. Update on glucocorticoid action and resistance. *J Allergy Clin Immunol* 2006;**117**:522–43.
- Matthews JG, Ito K, Barnes PJ, *et al.* Defective glucocorticoid receptor nuclear translocation and altered histone acetylation patterns in glucocorticoid-resistant patients. *J Allergy Clin Immunol* 2004;**113**:1100–8.
- Adcock IM, Lane SJ, Brown CR, *et al.* Abnormal glucocorticoid receptor-activator protein 1 interaction in steroid-resistant asthma. *J Exp Med* 1995;**182**:1951–8.
- Adcock IM, Barnes PJ. Molecular mechanisms of corticosteroid resistance. *Chest* 2008;**134**:394–401.
- Ito K, Yamamura S, Essilfie-Quaye S, *et al.* Histone deacetylase 2-mediated deacetylation of the glucocorticoid receptor enables NF- κ B suppression. *J Exp Med* 2006;**203**:7–13.
- Li LB, Leung DY, Martin RJ, *et al.* Inhibition of histone deacetylase 2 expression by elevated glucocorticoid receptor beta in steroid-resistant asthma. *Am J Respir Crit Care Med* 2010;**182**:877–83.
- Leung DY, Hamid Q, Vottero A, *et al.* Association of glucocorticoid insensitivity with increased expression of glucocorticoid receptor beta. *J Exp Med* 1997;**186**:1567–74.
- Goleva E, Li LB, Eves PT, *et al.* Increased glucocorticoid receptor beta alters steroid response in glucocorticoid-insensitive asthma. *Am J Respir Crit Care Med* 2006;**173**:607–16.
- Sousa AR, Lane SJ, Cidlowski JA, *et al.* Glucocorticoid resistance in asthma is associated with elevated in vivo expression of the glucocorticoid receptor beta-isoform. *J Allergy Clin Immunol* 2000;**105**:943–50.
- Christodouloulopoulos P, Leung DY, Elliott MW, *et al.* Increased number of glucocorticoid receptor-beta-expressing cells in the airways in fatal asthma. *J Allergy Clin Immunol* 2000;**106**:479–84.
- Bergeron CB, Fukakusa M, Olivenstein R, *et al.* Increased glucocorticoid receptor- β expression, but not decreased histone deacetylase 2, in severe asthma. *J Allergy Clin Immunol* 2006;**117**:703–5.
- Gagliardo R, Chanez P, Vignola AM, *et al.* Glucocorticoid receptor alpha and beta in glucocorticoid dependent asthma. *Am J Respir Crit Care Med* 2000;**162**:7–13.
- Torrego A, Pujols L, Roca-Ferrer J, *et al.* Glucocorticoid receptor isoforms alpha and beta in in vitro cytokine-induced glucocorticoid insensitivity. *Am J Respir Crit Care Med* 2004;**170**:420–5.
- Anon. Proceedings of the ATS Workshop on Refractory Asthma. Current understanding, recommendations, and unanswered questions. *Am J Respir Crit Care Med* 2000;**162**:2341–51.
- Bateman ED, Hurd SS, Barnes PJ, *et al.* Global strategy for asthma management and prevention GINA executive summary. *Eur Respir J* 2008;**31**:143–78.
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 2001;**25**:402–8.
- Britten KM, Howarth PH, Roche WR. Immunohistochemistry on resin sections: a comparison of resin embedding techniques for small mucosal biopsies. *Biotech Histochem* 1993;**68**:271–80.
- Kim SH, Kim DH, Lavender P, *et al.* Repression of TNF-alpha-induced IL-8 expression by the glucocorticoid receptor-beta involves inhibition of histone H4 acetylation. *Exp Mol Med* 2009;**41**:297–306.
- de Castro M, Elliot S, Kino T, *et al.* The non-ligand binding beta-isoform of the human glucocorticoid receptor (hGR beta): tissue levels, mechanism of action and potential physiologic role. *Mol Med* 1996;**2**:597–607.
- Smith CJ, Pearce BM. Clathrin: anatomy of a coat protein. *Trends Cell Biol* 1999;**9**:335–8.
- Leung DY, Hamid Q, Vottero A, *et al.* Association of glucocorticoid insensitivity with increased expression of glucocorticoid receptor beta. *J Exp Med* 1997;**186**:1567–74.
- Hausmann M, Herfarth H, Scholmerich J, *et al.* Glucocorticoid receptor isoform expression does not predict steroid treatment response in IBD. *Gut* 2007;**56**:1328–9.
- Honda M, Oriti F, Ayabe T, *et al.* Expression of glucocorticoid receptor beta in lymphocytes of patients with glucocorticoid-resistant ulcerative colitis. *Gastroenterology* 2000;**118**:859–66.
- Kozaci DL, Chernajovsky Y, Chikanza IC. The differential expression of corticosteroid receptor isoforms in corticosteroid-resistant and -sensitive patients with rheumatoid arthritis. *Rheumatology (Oxford)* 2007;**46**:579–85.
- Oriti F, Ashida T, Nomura M, *et al.* Quantitative analysis for human glucocorticoid receptor [alpha]/[beta] mRNA in IBD. *Biochem Biophys Res Commun* 2002;**296**:1286–94.
- Zhang X, Clark AF, Yorio T. Regulation of glucocorticoid responsiveness in glaucomatous trabecular meshwork cells by glucocorticoid receptor- β . *Invest Ophthalmol Vis Sci* 2005;**46**:4607–16.
- Ishida A, Ohta N, Koike S, *et al.* Overexpression of glucocorticoid receptor- β in severe allergic rhinitis. *Auris Nasus Larynx* 2010;**37**:584–8.
- Ito K, Ito M, Elliott WM, *et al.* Decreased histone deacetylase activity in chronic obstructive pulmonary disease. *N Engl J Med* 2005;**352**:1967–76.
- Ito K, Lim S, Caramori G, *et al.* Cigarette smoking reduces histone deacetylase 2 expression, enhances cytokine expression, and inhibits glucocorticoid actions in alveolar macrophages. *FASEB J* 2001;**15**:1110–12.
- Ito K, Caramori G, Lim S, *et al.* Expression and activity of histone deacetylases in human asthmatic airways. *Am J Respir Crit Care Med* 2002;**166**:392–6.
- Cosio BG, Mann B, Ito K, *et al.* Histone acetylase and deacetylase activity in alveolar macrophages and blood monocytes in asthma. *Am J Respir Crit Care Med* 2004;**170**:141–7.
- Cosio BG, Tsaprouni L, Ito K, *et al.* Theophylline restores histone deacetylase activity and steroid responses in COPD macrophages. *J Exp Med* 2004;**200**:689–95.
- Ito K, Lim S, Caramori G, *et al.* A molecular mechanism of action of theophylline: induction of histone deacetylase activity to decrease inflammatory gene expression. *Proc Natl Acad Sci U S A* 2002;**99**:8921–6.