events. The effect of macrolides was most evident in patients with pre-existing cardiovascular disease or at high risk of cardiovascular disease according to the QRISK2 score.

Conclusions The use of clarithromycin in the setting of AECOPD or CAP is associated with increased cardiovascular events.

Abstract S27 Figure 1

S28 INCREASED INCIDENCE OF COPD EXACERBATIONS FOLLOWING INFLUENZA VACCINATION

doi:10.1136/thoraxjnl-2012-202678.034

GC Donaldson, BS Kowlessar, ARC Patel, AJ Mackay, R Singh, S Brill, JA Wedzicha. *University College London, London, United Kingdom.*

Introduction Influenza vaccination prevents substantial morbidity, and mortality, and the spread of infection, but patients still sometimes refuse vaccinations due to concerns over adverse reactions. Our aim was to determine the relative risk of COPD exacerbations post-vaccination.

Methods We analysed data before and after 162 vaccinations given to 112 patients in the London COPD cohort between 2009 and 2011. Patients recorded on daily diary cards increase in respiratory symptoms. Exacerbations were defined as ≥ 2 days of 2 major symptoms (new or increased breathlessness, sputum volume or purulence) or 1 major and 1 minor symptom (cold, increased cough, increased wheeze, sore throat). Patients were telephoned in Oct 2011 and asked about recent or forthcoming vaccinations; other dates were known from previous years when recorded diary cards or reported at clinic visits.

The risk of COPD exacerbation in the two post-vaccination periods, days 1–2 and 3–14 were calculated relative to the patient's baseline risk assessed over the year before and after vaccination using a self-controlled case series approach. The data was also simply analysed with a Chi-squared test of the proportion of exacerbations in the two weeks before and after vaccination.

Results There was a 3.31 fold (95% CI 1.5–7.4; $P=0.004$) increased risk of COPD exacerbation 1–2 days post influenza vaccination, and a 1.8 fold risk (1.2–2.9; $p=0.007$) 3–14 days afterwards. In the two weeks prior to vaccination there were 5/162 exacerbations and 26/162 in the two weeks post vaccination ($p<0.001$). All 5 of the exacerbations before and 17/26 exacerbations after vaccination were treated with antibiotics and/or oral steroids. Using this healthcare utilisation definition of COPD exacerbations, there were still significantly more events post vaccination ($p=0.008$). Figure 1 shows the average number of new or increased respiratory symptoms recorded by the patients during the 3 weeks before and after vaccination.

Conclusion The incidence of COPD exacerbation increases immediately after influenza vaccination. COPD patients should be warned about increased respiratory symptoms after their influenza vaccination. Future studies should investigate the mechanisms underlying these increased symptoms in order to intervene and prevent them effectively.

Abstract S28 Figure 1

S29 THE PROPOSED NATIONAL EARLY WARNING SYSTEM (NEWS) COULD BE HAZARDOUS FOR PATIENTS WHO ARE AT RISK OF HYPERCAPNIC RESPIRATORY FAILURE

doi:10.1136/thoraxjnl-2012-202678.035

B Kane, S Decalmer, P Murphy, P Turkington, BR O'Driscoll. *Salford Royal Foundation Trust, Manchester, UK*

Introduction Early Warning Scoring systems (EWS) identify critical illness at an early stage. The Royal College of Physicians has proposed use of a 'National Early Warning Score' (NEWS) across UK hospitals. The NEWS allocates EWS points for oxygen saturation $<96\%$ (see Table) and does not take account of target oxygen saturation range which should be set for each patient according to the BTS oxygen guidelines. This is particularly important for patients at risk of hypercapnic respiratory failure (T2RF) who are safest with a target range of 88–92% or less. We compared the NEWS with a proposed alternative based on the Salford Royal Hospital EWS system (AltNEWS, see Table) to identify how many patients with COPD would be placed at risk of hyperoxaemia using the NEWS system.

Methods We calculated EWS scores using the NEWS and AltNEWS for 108 unselected acute medical patients at a single time point.

Results 34/108 general medical patients (31%) had risk factors for T2RF (30 COPD, 4 obstructive sleep apnoea). Nineteen of these 34 patients had saturations within their target range of 88–92% either on air or oxygen. The NEWS system allocated these patients 2 or 3 EWS points for "low" oxygen levels which could prompt nursing staff to increase supplemental oxygen, potentially precipitating dangerous hypercapnia. Three of these 34 patients had saturations $>92\%$ on oxygen. The NEWS did not alert nursing staff that supplemental oxygen should be reduced for these patients; saturations of 93–94% were actually scored as "too low". This could prompt nursing staff to further increase supplemental oxygen which could harm these patients. The AltNEWS allocated EWS points according to whether patients were in or out of range and no patients were placed at risk of T2RF.

Conclusion The NEWS system makes no allowance for patients at risk of T2RF. This may lead to potentially dangerous use of oxygen in this substantial group of patients. We recommend that a target oxygen saturation range should be set for all hospital patients on admission and oxygen scores within EWS systems should be adjusted to alert clinicians to scores above and below the target range.