

P34 RESPIRATORY TEAM LEADING SECONDARY CARE STOP SMOKING SERVICE

doi:10.1136/thoraxjnl-2012-202678.175

W Preston, K Hennessy, E Williams. *George Eliot Hospital, Nuneaton, Warwickshire*

There are over a billion smokers in the world and tobacco addiction is one of the biggest health treats for the world. Smoking tobacco kills half of its users and is estimated to cause up to one billion deaths in the 21st century (WHO, 2010). There is a public health drive to provide behavioural and medication support to reduce the amount of current smokers and prevent future smokers. Respiratory clinicians have an opportunity to lead change and facilitate services closer to patients; smoke cessation is the key evidence based treatment for COPD that will delay progression of disease (NICE, 2010). 15% of smokers will be diagnosed with COPD and alters the immune system; there is a 2.5 increased risk of contracting pneumonia, quadrupled risk of contracting tuberculosis (Bernhard, 2011).

This case study identifies how a respiratory team have implemented a secondary care stop smoking service lead by a BTS Stop Smoking Champion. The service design is discussed, lessons learnt and results. An innovative model attracts tariff payment for in reach into schools, colleges and workplaces which in turn funds the inpatient service.

The service has been operational for 24 months and has treated over 700 patients. Respiratory nurses and AHP's are fully qualified advisers and facilitate provision with the help of a full time co-ordinator. Support is available on wards and in outpatient with follow up for at least 12 weeks, nicotine replacement therapy available for inpatients 24 hours per day. All pre-op patients receive carbon monoxide testing with an opt out referral system. Referral systems in place from A/E to outpatients.

Evaluation has shown an average quit rate of 50%; 65% in COPD clinics with COPD CNS advisor. 14–18 year old complex smokers –20% quit rate and 85% in work place (routine and manual workers). NHS staff quit attempts have been consistently high (50%) with family groups also facilitated (80%).

To conclude respiratory staff leading on stop smoking services has been beneficial and rewarding; implementation enhanced by credibility and status of BTS role.

P35 A PHYSIOLOGIST-LED INTERSTITIAL LUNG DISEASE FOLLOW UP SERVICE: EXPERIENCES AND OUTCOMES IN A UK DISTRICT GENERAL HOSPITAL

doi:10.1136/thoraxjnl-2012-202678.176

MG Jones, K Butterfield, W McConnell. *Dorset County Hospital, Dorchester, Dorset*

Introduction and Objectives A proportion of patients with interstitial lung disease (ILD) maintain disease stability but require ongoing physiological and symptomatic monitoring. With increasing service demands and targets to reduce the proportion of follow up appointments, new approaches to use physician time efficiently whilst ensuring patient safety are required.

In 2006, under the supervision of a Consultant Respiratory Physician, a Physiologist Led ILD service was established in our hospital. Based on the clinical decision of the physician, patients with stable ILD at follow up are transferred for physiologist led follow up for ongoing observation. All patients attend for pulmonary function tests under the supervision of a physiologist at an interval determined by the physician. At this visit a patient symptom questionnaire is completed, and, if requested by the physician, a chest radiograph is performed. The information is subsequently reviewed by the physician who determines whether ongoing physiologist led follow up, further physician input, or discharge from follow up is required.

The Aim of this study was to retrospectively review service demand and patient outcomes over a 6 year period.

Methods Retrospective database and case review of all patients transferred by one respiratory consultant for physiologist led follow up from January 2006 until December 2011.

Results 106 patients were transferred for physiologist led follow up. Mean age at commencement of follow up was 71.2. 29 patients were stable throughout follow up and discharged (mean number of visits per discharged patient 2.46). 8 patients required physician input based on change in symptoms or physiological parameters. 6 of these patients were diagnosed with progression of ILD and 3 commenced on treatment. 2 subsequently died from ILD. No patient died from ILD whilst under physiologist led follow up. In 2011 48 follow up appointments were saved from one consultant's clinic lists.

Conclusions With appropriate patient selection, physiologist-led ILD follow up provides a safe and effective service whilst reducing the demand placed on physician follow up appointments.

Respiratory physiology and oxygen therapy

P36 AN ADVANCED PHYSIOLOGICAL MONITORING TOOL TO DETECT TREATMENT FAILURE IN HOSPITALISED PATIENTS WITH ACUTE EXACERBATION OF CHRONIC OBSTRUCTIVE PULMONARY DISEASE (AECOPD)

doi:10.1136/thoraxjnl-2012-202678.177

¹ES Suh, ¹S Mandal, ¹MC Ramsay, ²R Harding, ²A Thompson, ¹J Moxham, ³N Hart. ¹Division of Asthma, Allergy and Lung Biology, King's College London, London, UK; ²Lane Fox Respiratory Unit, Guy's St. Thomas' NHS Foundation Trust, London, UK; ³Guy's and St Thomas' NHS Foundation Trust/King's College London, National Institute Health Research Comprehensive Biomedical Research Centre, London, UK

Background Acute exacerbations of COPD are associated with significant mortality and morbidity. Hospitalised patients with AECOPD often deteriorate symptomatically despite treatment. We have developed 2nd intercostal space parasternal EMG (EMG_{para}) as a measure of neural respiratory drive (NRD), with the aim of detecting clinical change in such patients. We hypothesised that change in NRD would be able to detect treatment failure in an unselected cohort of AECOPD patients.

Methods Patients with AECOPD were recruited at a central London teaching hospital within 12 hours of admission. Patients underwent EMG_{para}, spirometry and measurement of inspiratory capacity at least daily until medically fit for discharge. Modified Borg scale was recorded with each EMG_{para} acquisition. We assessed improvement and deterioration by two measures: 1) the responsible medical team was asked to report any deterioration or improvement in the patients' condition during the course of admission (*Leidy et al, AJRCCM 2011*); and 2) 2-point changes in Borg score were recorded as representing significant deterioration or improvement in dyspnoea (*Ries et al, COPD; 2005*). Changes in EMG_{para%max} and neural respiratory drive index (Δ NRDI, where $\text{NRDI} = \text{EMG}_{\text{para}\% \text{max}} \times \text{respiratory rate}$) (*Murphy et al, Thorax; 2011*) during episodes of deterioration or improvement were analysed.

Results 65 patients were recruited. At admission, age was 71 ± 10 years, male 46.2%, FEV₁%predicted $32 \pm 15\%$ (n=34), body mass index (BMI) $25.6 \pm 7.7 \text{ kg/m}^2$, length of hospital stay 4 (IQR 2–5.75) days, admission COPD assessment tool (CAT) score 29 (IQR 25–33). There were 66(83%) episodes of improvement and 14(17%) episodes of deterioration as defined by *physician opinion*. There was a difference in the change in EMG_{para%max} between episodes of improvement and deterioration ($\Delta \text{EMG}_{\text{para}\% \text{max}} = -2.00\% \text{ vs } +4.99\%$, $p=0.01$; $\Delta \text{NRDI} = -53/\text{min vs } 117/\text{min}$, $p=0.02$). There were 54(65%) episodes of improvement and 29(35%) episodes of deterioration as defined by a 2-point change on the Borg scale. Again there was a difference in neural drive measures between improvers and deteriorators ($\Delta \text{EMG}_{\text{para}\% \text{max}} = -3.94\% \text{ vs } +2.72\%$, $p=0.002$; $\Delta \text{NRDI} = -108/\text{min vs } 74/\text{min}$, $p<0.001$) (Figure 1).