Further evidence supporting programmatic screening for, and treatment of latent TB Infection (LTBI) in new entrants to the UK from high TB prevalence countries

Lawrence Peter Ormerod^{1,2,3}

A systematic comparison of tuberculosis (TB) cases and rates between the USA and E&W (England and Wales), shows diametrically opposed trends. Between 1993 and 2010 in the USA, active TB cases fell from 25 107 to 11 182,¹ a fall of 55%. By contrast in E&W, notified TB cases rose from 5104 in 1993^{2 3} to 7901 in 2010,⁴ an increase of 55%.

Case management is not the explanation for the difference, as both countries use rifampicin and isoniazid based 6-month short course chemotherapy, with high completion rates, 92.7% for the USA¹ and 83.6% of survivors in E&W.⁴ Equally, different BCG policies do not account for the difference as the USA has never used BCG as a health control measure.

Both countries now have a majority of TB cases in foreign-born individuals. In the USA, in 2010, 60% of all cases were foreign born, with a rate of 18.1/100 000 for this group. They were predominantly Hispanic and Latin American, and from the Phillipines, with only 9% from India and only 1% from sub-Saharan Africa.¹ By contrast in E&W, 73% of cases were non-UK born, with 55% from South Asia (Pakistani rate 132/1 000 000; Indian rate 151/1 000 000), and 26% from Sub-Saharan Africa (rate 165/100 000), with only 23% of cases diagnosed within 2 years of initial entry.⁴ So both countries have a majority of cases in foreign born sub-populations, with those from E&W from different and even higher incidence countries. This of itself does not explain the divergence in trends, but differences in the management of new entrants, particularly in the diagnosis and treatment of latent TB Infection (LTBI) probably does.

The USA has had quite rigorous guidelines on the diagnosis and treatment of LTBI, in both recent entrants and other

risk groups for many years,⁵ recently updated to include Interferon- γ Release Assays (IGRA).⁶ By contrast in E&W although treatment of LTBI in children was mandatory, in young adult new entrants, this was discretionary from 1984 onwards.^{7 8} In 2006, with the switch to NICE guidance, this although showing on economic appraisal that treatment of LTBI was cost-beneficial if the incidence of LTBI was at 8% in a cohort, mainly because of the absence of data, limited new entrant screening age 16–34 for LTBI, to those from Sub-Saharan Africa, and other countries with an incidence of 500/100 000pa or more,⁹ which excluded the significant proportion of E&W new entrants from South Asia.

In 2010, more rational new entrant screening was advocated,¹⁰ this and early data on the near 30% IGRA positive rate for new entrants predominantly from South Asia,¹¹ were taken into account in a further revision of NICE guidance, with emphasis on the use of IGRA tests. This led to revised NICE recommendations in 2011,¹² which reduced the threshold for new entrant screening to all countries with an incidence of 40/100 000pa or more, thus covering the main ethnic new entrant groups seen,⁴ while also confirming support on health economic grounds. The paper by Pareek *et al*¹³ in this edition, provides further significant evidence supporting programmatic screening for, and treatment of LTBI in new entrants to the UK from high TB prevalence countries, whilst also questioning whether the threshold of 40/100 000 currently advised¹² is economically appropriate. Additional data of this type should allow refinement of the screening for LTBI to maximum efficacy and coverage.

Those involved in UK TB strategy believe than only implementation of much wider, effective and consistently applied screening for, and treatment of LTBI will change our current rising TB trend, into a persistent downward trend like the USA. The next challenge however will be ensuring that contracting and commissioning of TB services, ensures that the funding and staffing

of such TB services is sufficient for its systematic implementation.^{14 15}

Correction notice This article has been corrected since it was published Online First. The author name Parikh has been amended to read Pareek in the text and the references.

Competing interests None.

Provenance and peer review Commissioned; internally peer reviewed.

To cite Ormerod LP. *Thorax* 2013;**68**:201.

Received 7 August 2012

Accepted 21 August 2012

Published Online First 19 September 2012

► <http://dx.doi.org/10.1136/thoraxjnl-2011-201542>

Thorax 2013;**68**:201.

doi:10.1136/thoraxjnl-2012-202184

REFERENCES

- 1 Reported TB in the USA. 2010. <http://www.cdc.gov/tb/statistics/pdf>
- 2 Kumar D, Watson JM, Charlett A, *et al*. Tuberculosis in England and Wales in 1993: results of a national survey. *Thorax* 1997;**52**:1060–7.
- 3 Tuberculosis Section Health Protection Agency. personal communication. 03.08.12.
- 4 Tuberculosis in the UK: 2011 Report. London HPA Dec 2011.
- 5 Centres for Disease Control. Targeted tuberculin testing and treatment of Latent tuberculosis infection. *MMWR* 2000;**(49)**:RR6 1–43.
- 6 Centres for Disease Control. Updated guidelines for using interferon gamma release assays to detect *Mycobacterium tuberculosis* infection. *MMWR* 2010;**59**:R5 1–25.
- 7 Joint Tuberculosis Committee of the British Thoracic Society. Control and prevention of tuberculosis in the United Kingdom: Code of Practice 1994. *Thorax* 1994;**49**:193–1200.
- 8 Joint Tuberculosis Committee of the British Thoracic Society. Control and prevention of tuberculosis in the United Kingdom: Code of Practice. *Thorax* 2000;**5**:887–901.
- 9 NICE. Guideline 33. Clinical diagnosis and management of tuberculosis and measures for its control. 2006. <http://guidance.nice.org.uk/CG33>
- 10 Moore-Gillon J, Davies PDO, Ormerod LP. Rethinking TB screening: politics, practicalities and the press. *Thorax* 2010;**65**:663–5.
- 11 Pareek M, Watson JP, Ormerod LP, *et al*. Screening of immigrants in the UK, for imported latent tuberculosis, a multicentre cohort study and cost-effectiveness analysis. *Lancet Infect Dis* 2011;**6**:435–44.
- 12 NICE Guideline 117. Clinical diagnosis and management of tuberculosis and measures for its prevention and control. 2011. <http://guidance.nice.org.uk/CG 117>
- 13 Pareek M, Bond M, Shorey J, *et al*. Community-based evaluation of immigrant tuberculosis screening using interferon gamma release assays and tuberculin skin testing: observational study and economic analysis. *Thorax*, Published Online First: 12 June 2012. doi:10.1136/thoraxjnl-2011-201542.
- 14 NICE Public Health Guidance. Identifying and managing tuberculosis among hard to reach groups. 2012. <http://www.nice.org.uk/ph37>
- 15 Royal College of Nursing. Tuberculosis case management and cohort review. Guidance for health professionals. <http://www.rcn.org.uk/data/assets/pdf file/0010/439129/004204.pdf>

¹Chest Clinic, Royal Blackburn Hospital, Blackburn, Lancs, UK; ²Lancashire Postgraduate School of Medicine, University of Central Lancs, Preston, Lancs, UK; ³University of Manchester, Manchester, UK

Correspondence to Professor Lawrence Peter Ormerod, Department of Chest Clinic, Royal Blackburn Hospital, Blackburn, Lancs BB2 7AE, UK