developing countries do not and many rely on documents like GOLD to form policies. Does the world—including developing countries—really need promotion of more widespread use of antibiotics that in a year will lead to 80% of its users having throat swabs with macrolide-resistant bacteria and no information whatsoever on the impact on the spread of resistance in the community? What drug and dose should we have recommended: azithromycin 250 mcg once daily for years? Should we advise that everybody had ECGs, hearing tests and monitoring of liver enzymes—and how do the editors think this would work in the real world? Should we follow advice from US colleagues in *NEJM*² advocating 250 mg three times a week as a follow-up to the paper by Albert *et al*³ without any firm evidence that this dose has comparable efficacy? We felt that these questions precluded currently recommending use of macrolides despite some evidence of efficacy.

We would appreciate the assistance of the editors here—as would most people whose decisions were labelled ‘bizarre’ in *Thorax*.

Jørgen Vestbo,^{1,2} Roberto Rodriguez-Roisin³

¹Respiratory Research Group, University of Manchester, Manchester, UK

²Department of Respiratory Medicine, Odense University Hospital and University of Southern Denmark, Odense, Denmark

³Department of Pneumologia, Hospital Clinic, Barcelona, Spain

Correspondence to Professor Jørgen Vestbo, Respiratory Research Group, University of Manchester, Manchester M23 9LT, UK; jorgen.vestbo@manchester.ac.uk

Contributors JV and RR-R wrote this letter together.

Competing interests None.

Provenance and peer review Not commissioned; internally peer reviewed.

To cite Vestbo J, Rodriguez-Roisin R. *Thorax* 2013;**68**:387–388.

Received 25 November 2012

Accepted 4 December 2012

Published Online First 9 January 2013

► <http://dx.doi.org/10.1136/thoraxjnl-2012-203080>

Thorax 2013;**68**:387–388.

doi:10.1136/thoraxjnl-2012-203044

REFERENCES

- 1 Bush A, Pavord I. Highlights from this issue. *Thorax* 2012;**67**. doi:10.1136/thoraxjnl-2012-202790
- 2 Wenzel RP, Edmond MB. Antibiotic prevention of acute exacerbations in COPD. *N Engl J Med* 2012;**367**:340–7.

Letter to the editors

We read with interest your ‘Highlights from this Issue’ in the November issue of *Thorax*.¹ We noticed that you had a fresh comment on Global Initiative for Obstructive Lung Diseases (GOLD) not recommending macrolides to prevent exacerbations of chronic obstructive pulmonary disease (COPD).

We also noted that you hinted that it was probably too cheap for GOLD—suggesting that we were in the pockets of ‘big pharma’. We find the remark hurtful and wrong. GOLD does receive funding from the pharmaceutical industry—as do most of the major respiratory scientific societies and journals through advertising—but none of the people working with the GOLD document receives money for the work performed. In addition, GOLD has a steady income from sales of GOLD material and this income is sufficient to fund the work by the scientific committee, mainly expenses related to meetings at airport hotels.

However, we are much more interested in how the editors would have dealt with the evidence for macrolides in a global COPD strategy document.

As the UK has National Institute of Clinical Excellence (NICE), other well-off countries have bodies that advise and regulate on drug use. However, most

- 3 Albert RK, Connett J, Bailey WC, *et al.* for the COPD Clinical Research Network. Azithromycin for prevention of exacerbations of COPD. *N Engl J Med* 2011;365:689–98.