ORIGINAL ARTICLE

Efficacy and safety of the oral p38 inhibitor PH-797804 in chronic obstructive pulmonary disease: a randomised clinical trial

William MacNee,¹ Richard J Allan,² Ieuan Jones,² Maria Cristina De Salvo,³ Lisa F Tan²

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/thoraxjnl-2012-202744>).

¹Queen's Medical Research Institute, University of Edinburgh, Edinburgh, UK

²Department of Clinical Research, Pfizer Global Research and Development, Sandwich, UK

³Centro Medico Dra. De Salvo, Buenos Aires, Argentina

Correspondence to

Dr Lisa F Tan, Department of Clinical Research, Pfizer Global Research and Development, Field House, Field Road, Kingston, Cambridge CB23 2NQ, UK; lisa.tan@pfizer.com

Received 17 September 2012

Revised 1 March 2013

Accepted 4 March 2013

Published Online First

28 March 2013

► <http://dx.doi.org/10.1136/thoraxjnl-2013-203498>

To cite: MacNee W, Allan RJ, Jones I, et al. *Thorax* 2013;**68**: 738–745.

ABSTRACT

Rationale Chronic obstructive pulmonary disease (COPD) is a common lung disease leading to progressive decline in lung function. Inhibition of release of inflammatory mediators by p38 inhibitors may be a useful treatment for chronic inflammation of the airways thought to underlie the pathogenesis of the disease.

Objectives To evaluate the efficacy and safety of PH-797804, a potent and selective p38 inhibitor, in adults with moderate to severe COPD (Global Initiative for Chronic Obstructive Lung Disease stage II/III).

Methods This was a randomised, adaptive design, double-blind, placebo-controlled, parallel-group, multicentre trial. Patients were initially randomised to placebo, 0.5, 3, 6 or 10 mg PH-797804 once daily and treated for 6 weeks following a 2-week run-in.

Measurements and main results The primary endpoint was change from baseline in trough forced expiratory volume in 1 s (FEV₁) compared with placebo after 6 weeks of treatment. Secondary endpoints included other spirometric parameters, transition dyspnoea index, rescue medication use, high sensitivity C-reactive protein and symptoms. A total of 230 patients were assigned to treatment; placebo (n=45), 0.5 mg (n=20), 3 mg (n=47), 6 mg (n=70) and 10 mg (n=48). PH-797804 showed a statistically significant improvement in trough FEV₁ at week 6 compared with placebo of 0.086 litre (95% Bayesian CI 0.008 to 0.164) and 0.093 litre (95% CI 0.018 to 0.166) at 3 and 6 mg PH-797804, respectively. PH-797804 3 mg and 6 mg showed an improvement in the baseline dyspnoea index/transition dyspnoea index total focal score at week 6. PH-797804 was well tolerated at all doses studied.

Conclusions PH-797804 demonstrated improvements over placebo in lung function parameters and dyspnoea in patients with moderate to severe COPD.

TrialRegNo NCT00559910.

INTRODUCTION

Chronic obstructive pulmonary disease (COPD) is a common disease leading to a progressive decline in lung function. COPD is associated with high morbidity and death, and the direct and indirect socio-economic costs of COPD are high.¹ COPD is currently ranked as the fourth leading cause of death worldwide and is predicted to rise.^{2,3}

Treatments are limited, providing inadequate symptomatic relief and reduction in exacerbations. No therapeutic class has consistently shown

Key messages

What is the key question?

- What is the efficacy and safety of 6 weeks of treatment with PH-797804, an oral p38 inhibitor, in patients with moderate to severe chronic obstructive pulmonary disease (COPD)?

What is the bottom line?

- Six weeks of treatment with PH-797804 led to improvements over placebo in lung function and dyspnoea in patients with moderate to severe COPD with few adverse events observed.

Why read on?

- This study supports further investigation of PH-797804 as a potential treatment for patients with COPD.

anti-inflammatory properties across the spectrum of disease severity. Hence, there is still a need for treatments with demonstrable anti-inflammatory activity and potential clinical benefit.⁴

Signalling through p38 mitogen-activated protein kinase (p38-MAPK) is required for the expression of a range of inflammatory mediators associated with the chronic lung inflammation characteristic of COPD, such as tumour necrosis factor α , interleukin-1 (IL-1), IL-6 and IL-8.⁵ p38-MAPK is expressed on inflammatory cells associated with COPD, and there is increased lung expression and activation of p38-MAPK in patients with COPD compared with smoking and non-smoking controls.⁶ p38-MAPK inhibition, but not corticosteroids, attenuated oxidative stress-induced cytokine release in an in vitro assay modelling some aspects of COPD-associated inflammation.⁷ Therefore, p38-MAPK inhibition is an attractive target for COPD.

PH-797804 is a potent, selective p38-MAPK inhibitor,^{8,9} being evaluated as a potential oral anti-inflammatory COPD treatment. The primary objective of this trial was to assess the efficacy and safety of PH-797804 in adults with moderate to severe COPD defined by the guidelines applicable at the time (Global Initiative for Chronic Obstructive Lung Disease (GOLD) stages II/III).¹⁰ Secondary objectives included understanding the dose–response relationship and the time course of response to PH-797804.

These data were first presented at the European Respiratory Society 2010 Annual Congress in Barcelona, Spain.¹¹

METHODS

Patients

This randomised, double-blind, placebo-controlled, parallel-group, phase II trial was conducted between February 2008 and December 2009 in patients (40–80 years) with moderate to severe COPD in 38 centres across 13 countries. Patients had a diagnosis for at least 6 months and had a post-bronchodilator forced expiratory volume in 1 s (FEV₁)/forced vital capacity (FVC) ratio <0.7 with a post-bronchodilator FEV₁ of 30–80% of predicted. Patients with stable disease (no exacerbation in the preceding month) and stable FEV₁ during the run-in phase were eligible for enrolment. Key exclusion criteria included significant comorbidities or laboratory abnormalities, indicating significant concomitant disease. Long-acting bronchodilators and inhaled corticosteroids were not allowed 2 or 4 weeks prior to screening, respectively. To achieve this, withdrawal of treatment was necessary in some patients. Throughout the trial, patients received ipratropium bromide metered dose inhalers (MDIs; 40 µg four times a day) as maintenance therapy and salbutamol MDIs (100–200 µg as required) as rescue medication. Inclusion/exclusion criteria are described in the online data supplement. Independent ethics committees approved the trial and patients provided written informed consent. The trial was conducted in accordance with the principles of the Declaration of Helsinki and Good Clinical Practice guidelines.

Trial design

The trial involved a screening visit, a run-in phase (weeks –2 and –1 to ensure stability of lung function), a baseline/randomisation visit (week 0), 6 weeks of double-blind treatment with PH-797804 or placebo (visits at weeks 1, 2, 3, 4 and 6), a 2-week run-out phase and a final follow-up visit (week 8). Patients withheld ipratropium bromide and salbutamol for at least 8 h before clinic visits.

The primary endpoint was the change from baseline in trough FEV₁ compared with placebo after 6 weeks of treatment. Secondary efficacy endpoints included forced expiratory volume in 6 s (FEV₆), FVC and inspiratory capacity (IC). Trough (prior to trial medication) measurements for these spirometry parameters were recorded at all visits. Measurements were also recorded post trial medication (weeks 0 and 6) and post salbutamol (screening and weeks –1, –2, 0 and 6). Baseline dyspnoea index^{12–13} was evaluated at week 0, and transition dyspnoea index (TDI) was evaluated at weeks 1, 2, 3, 4 and 6. Patients maintained a diary throughout the trial to record COPD symptoms, maintenance and rescue bronchodilator use and peak expiratory flow rate. Spirometry was performed using standardised equipment with centralised over-read/interpretation in accordance with guidelines.¹⁴

Safety endpoints included adverse events, laboratory data, ECGs and vital signs. High sensitivity C-reactive protein (hsCRP) was measured at each visit. Exploratory biomarkers Clara cell protein-16 (CC16), IL-6, surfactant protein D (SPD) and fibrinogen were measured at weeks 0 and 6.

Randomisation and masking

Patients were initially randomised (according to a computer-generated randomisation code) to one of five treatment groups in the ratio 1:1:1:2:1 for placebo, 0.5 mg, 3 mg, 6 mg, 10 mg once daily PH-797804, respectively, until the interim analysis.

An independent Data Monitoring Committee (DMC) conducted an unblinded interim analysis once 96 patients had

completed 6 weeks of treatment. The DMC could drop a dose(s) in the event of statistical evidence of predefined futility or if unacceptable adverse events occurred, and could adapt the randomisation to increase the sample size for the 10 mg dose.

Trial treatments were supplied as PH-797804 capsules and matching placebo. Patients, investigator staff and the sponsor's project team were masked to treatment assignment throughout the trial until the database was locked. The randomisation code was provided to the DMC for the interim analysis. During the trial, the DMC could review unblinded data, at any time, in the event of safety concerns.

Statistical analysis

The analysis of the primary endpoint used Bayesian statistics. This approach has advantages over classical hypothesis testing because direct probabilistic statements about the effect size could be made (not addressed by p values) and information on placebo could be utilised from previous trials, reducing the number of placebo patients required.¹⁵ The Bayesian estimation of the dose–response relationship also means that the sample size is reduced compared with pairwise testing because information is used from all doses. Bayesian spline (normal dynamic linear model analyses)¹⁶ was applied that adjusted for baseline covariate.

It was of interest to characterise the dose–response relationship and calculate the probability of any dose achieving statistical significance and ascertaining whether the average treatment effect was minimally clinically relevant, defined as ≥ 75 ml improvement over placebo in FEV₁.¹⁷ A statistically significant result was defined if there was $\geq 95\%$ probability that the treatment effect over placebo was positive. An additional hurdle required that the average effect over placebo was >75 ml. This hurdle was for internal decision-making purposes to give confidence that the magnitude of effect was sufficient to warrant further investigation.

The anticipated clinical dose was 6 mg. Simulations showed that with 48 patients per group and 64 for 6 mg, the power of the study was such that there was 94% probability of a dose achieving significance if the true effect size in FEV₁ was 75 ml, and a 9% probability if the true effect size was only 20 ml (akin to type 1 error). The analysis of FEV₁ used Bayesian statistics with an uninformative prior for placebo (WinBUGS¹⁸). No multiplicity adjustments were required because this issue was addressed implicitly in the Bayesian model fitted.

Secondary endpoints were analysed using a classical repeated measures analysis of covariance approach using SAS version 9.1.¹⁹ To adjust for multiplicity of testing several dose groups against placebo, for secondary endpoints, a controlled testing procedure was used to control the overall α level at 5%. The mixed-effect model includes all available information over time and provides appropriate, consistent estimates of model parameters under the assumption of ‘missing at random’.

Further details concerning the statistical methodology and models used can be found in the online data supplement.

All primary and secondary efficacy endpoints were analysed using the full analysis set, defined as all randomised patients with at least one valid FEV₁ measurement in the double-blind phase. All safety analyses were conducted on the safety analysis set, defined as all patients who received at least one dose of PH-797804 or placebo.

Predefined futility rules

At the interim analysis, the dose–response relationship was estimated. The probability of a dose being futile was calculated. Doses could be stopped for futility if there was $<10\%$ chance

Figure 1 Trial profile. The full analysis set was used for all primary and secondary endpoints and was defined as all randomised patients who had at least one valid forced expiratory volume in 1 s measurement during the double-blind phase of the trial. The safety analysis set comprised all patients who received at least one dose of PH-797804 or placebo. *DBT=double-blind treatment.

of there being ≥ 75 ml improvement over placebo in FEV₁ change from baseline. Discontinued doses still contributed information to the dose-response curve at the end-of-study analysis.

RESULTS

A total of 230 patients were randomised to treatment and 201 completed double-blind treatment. Forty-five patients received placebo and 185 received PH-797804: 0.5 mg (n=20), 3 mg (n=47), 6 mg (n=70) or 10 mg (n=48). Figure 1 shows the trial profile. Demographics and baseline characteristics were similar among the five treatment groups (table 1).

Following the interim analysis, the 0.5 mg treatment group was dropped for futility and the randomisation ratio was modified as planned with post-interim patients randomised in the ratio 1:0:1:1:1 for placebo, 0.5 mg, 3 mg, 6 mg, 10 mg PH-797804, respectively.

The Bayesian analyses show that in this trial there is 99.2% probability that the true effect of 6 mg PH-797804 over placebo is greater than zero (ie, statistically significant). Additionally, the

analysis shows that the probability of this effect being clinically relevant (>75 ml improvement) is 67.3%. The Bayesian CIs, called credible intervals (CrI), are wide and show that there is a probability of 0.95 that the true effect over placebo for 6 mg lies between 18 and 166 ml. The results are similar for 3 mg PH-797804 (table 2).

There was only one outlier; a subject in the 3 mg PH-797804 group had an improvement in trough FEV₁ of approximately 1 litre, influencing the mean effect; when removed from the analyses the mean effect at 3 mg was 61 ml (CrI=−9 to 132 ml).

The probability of 10 mg PH-797804 being better than placebo was 94.5% (almost classed as statistically significant), but this trial also required that the magnitude of effect was large enough (the probability that the true effect is 75 ml was $<50\%$).

By contrast, 0.5 mg showed only numerical improvements over placebo of 59 ml.

Figure 2 shows the mean change from baseline in trough FEV₁ for each treatment group over the 6 weeks of treatment. FEV₁ declined by 52 ml in the placebo group, while it increased

Table 1 Demographic and baseline characteristics

	Placebo	PH-797804 dose group			
		0.5 mg	3 mg	6 mg	10 mg
N	45	20	47	70	48
Age (years)	64 (7.6)	64 (9.5)	62 (7.8)	65 (7.0)	65 (6.8)
Men	26 (58%)	12 (60%)	32 (68%)	55 (79%)	38 (79%)
Weight (kg)	72.8 (16.3)	70.4 (11.3)	74.5 (16.3)	77.0 (13.9)	77.1 (13.6)
Current smoker	18 (40%)	2 (10%)	23 (49%)	21 (30%)	16 (33%)
Time since diagnosis (years)	5.3 (0.8, 19.3)	7.1 (0.6, 28.0)	5.1 (0.9, 30.8)	6.0 (0.7, 21.0)	5.7 (0.6, 19.0)
Trough FEV ₁ at baseline (ml)	1080 (370)	1230 (400)	1270 (500)	1400 (420)	1380 (540)
FEV ₁ /FVC ratio at screening	0.44 (0.11)	0.46 (0.10)	0.46 (0.11)	0.47 (0.11)	0.47 (0.10)
% Predicted FEV ₁ at screening	47 (10.0)	56 (12.0)	53 (10.9)	56 (12.7)	56 (12.2)

Data are mean (SD), number (%) or mean (min–max). Data are from the safety analysis set (ie, all patients who received at least one dose of PH-797804/placebo), except for trough FEV₁ at baseline, FEV₁/FVC ratio and % predicted FEV₁ at screening which are from the full analysis set. FEV₁, forced expiratory volume in 1 s; FVC, forced vital capacity.

Table 2 Trough FEV₁: treatment comparisons versus placebo at week 6

FEV ₁ (ml)	Placebo	PH-797804 dose			
		0.5 mg	3 mg	6 mg	10 mg
N	42	18	44	65	42
Raw mean change from baseline	−43	18	38	37	4
SD of change from baseline	160	128	235	192	201
Estimated change from baseline*	−52	7	34	41	15
95% CrI for change from baseline	(−110 to 6)	(−62 to 82)	(−17 to 86)	(−2 to 85)	(−42 to 73)
Estimated difference from placebo		59	85	92	66
95% CrI of effect over placebo		(−23 to 149)	(8 to 164)	(18 to 166)	(−17 to 151)
Probability of effect >0 ml†		0.927	0.984	0.992	0.945
Probability of effect >75 ml‡		0.331	0.601	0.673	0.408

*The estimates are derived from the fitted statistical Bayesian model (normal dynamic linear model).

†A value >0.95 indicates statistical significance.

‡A value >0.5 indicates that the magnitude of effect is of interest.

CrI, credible interval (ie, Bayesian CI); FEV₁, forced expiratory volume in 1 s.

in each of the PH-797804 groups. There was an improvement in FEV₁ over placebo as a function of time with clear evidence of a treatment effect given the separation of all active treatments from placebo. The observed maximum estimated effect was in the 10 mg group at week 2, this difference then reduced over time. The results of the statistical analysis of trough FEV₆, FVC and IC at week 6 are presented in table 3.

PH-797804 at doses of 3, 6 and 10 mg produced a statistically significant improvement in trough FEV₆ at week 6 versus placebo, with mean changes from placebo of 115 ml (SE=54), 87 ml (SE=50) and 94 ml (SE=55), respectively. PH-797804 at doses of 6 and 10 mg produced a statistically significant improvement in trough FVC at week 6 versus placebo, with mean changes from placebo of 111 ml (SE=54) and 128 ml (SE=59), respectively. PH-797804 at doses of 3, 6 and 10 mg produced a statistically significant improvement in trough IC at week 6 versus placebo, with mean changes from placebo of 119 ml (SE=57), 119 ml (SE=53) and 96 ml (SE=58), respectively.

Figure 2 Mean change from baseline in trough forced expiratory volume in 1 s (FEV₁) (ml) during treatment. The symbols and bars represent the means and 95% CIs.

No statistically significant differences from placebo in FEV₁, FEV₆, FVC or IC were observed when post-dose (PH-797804 or bronchodilator) measurements were compared with pre-dose measurements.

PH-797804 at doses of 3 and 6 mg showed an improvement in the TDI total focal score at week 6 at the 10% and 5% significance level, with a 0.95 point (SE=0.646) and 0.99 point (SE=0.596) improvement, respectively (table 4). This improvement was driven by the magnitude of effort component as opposed to the magnitude of task or functional impairment components. Mean TDI for each treatment over time is presented in figure 3.

Table 5 summarises treatment comparisons with placebo for the change from baseline in rescue medication (salbutamol) usage. A statistically significant decrease in rescue medication use was observed with 6 mg PH-797804 with a mean daily use of 0.66 less actuations of salbutamol at week 6.

Log hsCRP values were analysed and adjusted for log base-lines. A decrease in hsCRP compared with placebo occurred with the 3, 6 and 10 mg doses of PH-797804, with statistically significant differences from placebo observed in the ratio of the means of 0.633 (p=0.033), 0.588 (p=0.011) and 0.594 (p=0.021), respectively after 6 weeks treatment. There were no statistically significant differences from placebo for CC16, IL-6, SPD or fibrinogen.

COPD symptoms were assessed using an internal unvalidated daily paper diary. Individual item analyses found no significant differences over placebo at the end of treatment compared with baseline. A total dyspnoea incidence score was calculated to determine the number of dyspnoea-experienced and dyspnoea-free days. Trends were seen for 6 mg to affect dyspnoea days.

The most frequently reported adverse events were COPD exacerbation, rash and nasopharyngitis (table 6). Rash was the most frequently reported treatment-related adverse event. All reported rashes resolved on cessation of treatment. Three patients (6 mg PH-797804) experienced serious adverse events considered to be treatment related by the investigator; two had gastrointestinal (GI) haemorrhages (one with a history of GI complications including past GI bleed, partial gastrectomy and heavy alcohol intake; the event occurred on day 2 of the trial). The other had been self-medicating with over-the-counter aspirin. One patient had left and right bundle branch block (also present on pretrial ECGs). No significant trends were

Table 3 Trough FEV₆, FVC and IC: treatment comparisons versus placebo at week 6

		PH-797804 dose			
	Placebo	0.5 mg	3 mg	6 mg	10 mg
FEV ₆ (ml)					
N (baseline)	42	18	44	65	42
Baseline, mean (SD)	2170 (610)	2410 (560)	2480 (830)	2680 (630)	2590 (820)
N (week 6)	40	17	40	58	39
Adjusted mean change	−77	−2	38	11	17
SE of the mean	39	58	38	32	39
95% CI	(−153 to −1)	(−117 to 113)	(−37 to 113)	(−52 to 73)	(−59 to 93)
Difference between means*		75	115	87	94
SE of difference*		70	54	50	55
95% one-sided lower limit for difference*		−41	26	4	4
p Value (difference between means)*		0.143	0.017	0.042	0.044
FVC (ml)					
N (baseline)	42	18	44	65	42
Baseline, mean (SD)	2630 (770)	2850 (680)	2960 (1020)	3190 (790)	2960 (940)
N (week 6)	40	17	40	58	39
Adjusted mean change	−91	−2	−1	20	38
SE of the mean	42	63	41	34	42
95% CI	(−173 to −8)	(−127 to 123)	(−83 to 80)	(−48 to 88)	(−45 to 120)
Difference between means*		89	89	111	128
SE of difference*		76	59	54	59
95% one-sided lower limit for difference*		−37	-8	21	30
p Value (difference between means)*		0.122	0.065	0.022	0.016
IC (ml)					
N (baseline)	42	18	44	65	42
Baseline, mean (SD)	1920 (560)	2040 (570)	2130 (620)	2310 (620)	2210 (690)
N	40	17	40	58	39
Adjusted mean change	−53	52	66	66	43
SE of the mean	41	62	40	34	41
95% CI	(−133 to 27)	(−70 to 174)	(−13 to 145)	(0 to 133)	(−38 to 123)
Difference between means*		105	119	119	96
SE of difference*		74	57	53	58
95% one-sided lower limit for difference*		−17	25	32	0
p Value (difference between means)*		0.078	0.019	0.013	0.049

*Comparisons are between each dose of PH-797804 and placebo using the full analysis set, which included all patients who were randomised and had a valid FEV₁ measurement during the double-blind treatment phase.

IC, inspiratory capacity; FEV₁, forced expiratory volume in 1 s; FEV₆, forced expiratory volume in 6 s; FVC, forced vital capacity.

Table 4 TDI total focal score: treatment comparisons versus placebo for change from baseline at week 6

TDI total focal score	Placebo	PH-797804 dose			
		0.5 mg	3 mg	6 mg	10 mg
N (baseline)	42	18	44	65	42
Baseline, mean (SD)	6.74 (2.20)	6.89 (1.64)	7.08 (1.92)	6.80 (2.14)	6.50 (2.38)
N (week 6)	40	17	40	58	40
Adjusted mean change	1.580	1.090	2.527	2.571	1.443
SE of the mean	0.459	0.701	0.455	0.379	0.460
95% CI	(0.674 to 2.485)	(−0.292 to 2.472)	(1.630 to 3.424)	(1.823 to 3.320)	(0.535 to 2.351)
Difference between means*		−0.490	0.947	0.992	−0.137
SE of difference*		0.838	0.646	0.596	0.650
95% one-sided lower limit for difference*		−1.874	−0.121	0.007	−1.212
p Value (difference between means)*		0.720	0.072	0.049	0.583

*Comparisons are between each dose of PH-797804 and placebo using the full analysis set, which included all patients who were randomised and had a valid FEV₁ measurement during the double-blind treatment phase.

FEV₁, forced expiratory volume in 1 s; TDI, transition dyspnoea index.

Figure 3 Mean change from baseline in transitional dyspnoea index (TDI) total focal score during treatment. The symbols and bars represent the means and 95% CIs.

noted in any laboratory safety parameters or vital signs. There were no significant trends in mean ECG parameters, although there was an increase in the percentage of patients with changes from baseline in QT interval corrected for heart rate using Fridericia's formula (QTcF) >30 ms. No patients had a QTcF of >500 ms.

DISCUSSION

To the authors' knowledge, this is the first clinical trial to demonstrate a positive effect of a p38 inhibitor on a range of clinical outcomes in COPD. Six weeks of treatment with PH-797804 in patients with COPD (GOLD II/III) maintained on a background of short-acting bronchodilators led to statistically significant improvements in lung function associated with improvements in dyspnoea and rescue medication usage.

By week 6, the 6 mg PH-797804 group showed an improvement over placebo in prebronchodilator FEV₁ of 93 ml and a 0.99-point improvement in the TDI (one point change is considered clinically meaningful).^{12 13} The minimum target effect on FEV₁ was predefined as 75 ml over placebo, based on data obtained with the phosphodiesterase 4 inhibitor roflumilast, which showed placebo-corrected increases of 49 ml and 80 ml in prebronchodilator FEV₁ in two 6-month studies.¹⁷ This

magnitude of the change in FEV₁ is below the defined minimally clinically important difference of 100 ml derived from the improvement in FEV₁ bronchodilator trials.²⁰ It is recognised that FEV₁ improvements may be less with drugs having an anti-inflammatory rather than a bronchodilator effect. However, the change in TDI and decrease in rescue medication alongside the other changes in lung function suggest that the effects may be of clinical relevance.

PH-797804 at doses of 3 and 6 mg consistently showed an improvement in FEV₁ over placebo throughout the trial. In addition, FEV₁ improved compared with baseline throughout the trial in all PH-797804 groups, whereas the placebo group showed a decline in lung function over the duration of the trial. This decline is greater than expected in a 6-week trial and may have been contributed to by the fact that subjects were maintained on short-acting bronchodilators alone.

Other lung function parameters, such as FVC, FEV₆ and IC, showed improvements consistent with the FEV₁ improvement; this was maintained in post-bronchodilator measures, indicating an additive effect of PH-797804 on top of salbutamol. There was no acute effect of PH-797804 on lung function, supporting its mode of action as anti-inflammatory rather than a direct bronchodilator.

PH-797804 reduced hsCRP at doses above 0.5 mg, indicating a systemic anti-inflammatory effect. This effect was maintained throughout the 6-week dosing period. Statistically significant effects were not observed on other systemic biomarkers measured in this study. A 12-week study in patients with COPD, with another p38 inhibitor, losmapimod, examined the effect of p38 inhibition on sputum neutrophils. No statistically significant effect was seen on this endpoint. However, a statistically significant reduction in plasma fibrinogen and trends towards an effect on systemic IL-6, IL-8 and CRP were observed but no statistically significant changes in lung function.²¹

The most consistently efficacious dose of PH-797804 was 6 mg, not the highest dose of 10 mg; the reason for this is unknown. The SEs of the mean data are large, therefore this may be due to variability. An alternative explanation could be tachyphylaxis, although the mechanism of this is unknown. This phenomenon will be investigated in longer-term studies. Potential differences in baseline severity of COPD, systemic exposure compared with other dose groups, and preponderance of adverse events potentially leading to a higher dropout rate or functional unblinding do not explain the apparent lower effect size of the 10 mg group.

Table 5 Treatment comparisons versus placebo for change from baseline in mean rescue medication usage at week 6

Rescue medication usage (daily no. of puffs averaged over each week)	PH-797804 dose				
	Placebo	0.5 mg	3 mg	6 mg	10 mg
N (baseline)	40	16	43	63	42
Baseline, mean (SD)	3.40 (2.84)	2.03 (2.30)	3.50 (2.47)	3.42 (4.81)	2.55 (2.34)
N (week 6)	38	15	40	54	38
Adjusted mean change (puffs per day)	0.309	0.025	-0.165	-0.351	0.115
SE of the mean	0.249	0.395	0.241	0.204	0.246
95% CI	(-0.183 to 0.801)	(-0.754 to 0.805)	(-0.641 to 0.311)	(-0.753 to 0.052)	(-0.372 to 0.601)
Difference between means*		-0.283	-0.473	-0.659	-0.194
SE of difference*		0.467	0.346	0.322	0.351
95% one-sided lower limit for difference*		0.490	0.100	-0.127	0.386
p Value (difference between means)*		0.273	0.087	0.021	0.291

*Comparisons are between each dose of PH-797804 and placebo using the full analysis set, which included all patients who were randomised and had a valid FEV₁ measurement during the double-blind treatment phase.
FEV₁, forced expiratory volume in 1 s;

Table 6 All-cause treatment-emergent adverse events in at least two patients in any treatment group

Adverse event	Patients, N (%)				
	Placebo	PH-797804			
		0.5 mg	3 mg	6 mg	10 mg
Patients (N)	45	20	47	70	48
Patients with adverse events (N)	17 (38)	4 (20)	19 (40)	37 (53)	23 (48)
COPD exacerbation	5 (11)	0	3 (6)	1 (1)	6 (13)
Rash*	1 (2)	1 (5)	3 (6)	9 (13)	1 (2)
Nasopharyngitis	2 (4)	0	5 (11)	3 (4)	0
Diarrhoea	0	0	1 (2)	3 (4)	2 (4)
Headache	1 (2)	0	0	5 (7)	0
Tremor	0	0	0	1 (1)	4 (8)
Pharyngitis	1 (2)	0	0	2 (3)	1 (2)
Urinary tract infection	1 (2)	0	0	1 (1)	2 (4)
Increased ALT†	0	0	1 (2)	1 (1)	2 (4)
Dizziness	2 (4)	0	0	0	1 (2)
Dyspepsia	1 (2)	0	0	2 (3)	0
Fatigue	0	0	0	0	2 (4)
Nausea	0	0	0	2 (3)	1 (2)
Arthralgia	0	0	0	2 (3)	0
Left bundle branch block	0	0	0	2 (3)	0
Right bundle branch block	0	0	0	2 (3)	0
Chest pain	0	0	0	0	2 (4)
Dry skin	0	0	0	2 (3)	0
Syncope	0	0	0	2 (3)	0

If the same patient in a given treatment had more than one occurrence in the same preferred term event category (or more than one of the rash terms in the combined category), the patient was only counted once per treatment group.

Data are from the safety analysis set.

*Rash includes the following terms: rash, rash generalised, rash macular, rash pustular, dermatitis, dermatitis acneiform and pruritus.

†The maximum ALT increase was less than twice the upper limit of normal range.

ALT, alanine aminotransferase; COPD, chronic obstructive pulmonary disease.

This study was on a background of short-acting bronchodilators alone and it recognised that these are not the mainstay of treatment in COPD. It's anticipated that PH-797804 would also show additive effects to long-acting bronchodilators. This is being studied in ongoing trials (clinicaltrials.gov NCT0153919 and NCT01321463).

A further confounding feature in this study is the effect of withdrawal of treatment with long-acting β agonists in some patients in the run-in period, which was considered to have influenced the results of many pharmacological trials.²² In our study a similar number of subjects (<50%) in each of the arms of the study had treatment withdrawn, which makes it less likely that differences between placebo and treatment groups resulted from differences in treatment withdrawal between the groups.

Data from trials of agents such as roflumilast, tiotropium, salmeterol and fluticasone suggest that, although the changes from baseline in FEV₁ may decrease with time in patients with COPD, the difference between active treatment and placebo generally appears to be maintained across the duration of the studies.^{23–27} The duration of this trial (6 weeks of dosing) was relatively short compared with the likely long-term use of efficacious COPD medications and the long-term effects of PH-797804 will be explored in longer studies.

An effect on COPD exacerbations could not be adequately assessed during a 6-week trial, however we propose that the anti-inflammatory properties of PH-797804 which led to the improvements in lung function and dyspnoea measures may translate to reductions in COPD exacerbations over a longer period.

The overall safety profile of PH-797804 in this trial was good, with most patients assigned to PH-797804 treatment groups completing the double-blind treatment phase. The most commonly reported treatment-related adverse event was acneiform rash, which resolved upon cessation of treatment. This has been reported with other p38 inhibitors.²⁸ There were no significant effects on liver function, unlike with other p38 inhibitors, when increases in alanine transaminase and aspartate transaminase have been reported.²⁹

In conclusion, PH-797804 showed improvements in lung function and dyspnoea scores in patients with moderate to severe COPD treated with short-acting bronchodilators. These data are encouraging and support further investigation of this compound in patients with COPD.

Acknowledgements The authors would like to thank the investigators and patients who participated in the trial. Medical writing support was provided by Samantha Abel, Valley Writing Solutions Ltd. This support included drafting the initial manuscript based on discussions with the authors and subsequent updates based on author comments. This assistance was funded by Pfizer Global Research and Development.

Funding This trial and all trial-related activities including medical writing support for the manuscript were funded by Pfizer Global Research and Development, Sandwich, UK.

Competing interests R Allan, L Tan and I Jones are or were employees of Pfizer and own or owned a limited amount of Pfizer stock. W MacNee has received funding for patient recruitment and participation in this trial and has received funding for research projects unrelated to this study from Pfizer. W MacNee has also received funding for work unrelated to this study from Almirall, Boehringer-Ingelheim, GlaxoSmithKline, Grifols, Novartis and Janssen. MC De Salvo has received funding for patient recruitment and participation in this trial.

Ethics approval This was a multicentre study and was approved by multiple independent ethics committees as appropriate for each centre/country.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- Barnes PJ. *Managing chronic obstructive pulmonary disease*. London: Sci Press, 1999.
- World Health Statistics. 2008. http://www.who.int/whosis/whostat/EN_WHS08_Full.pdf (accessed 12 Mar 2013).
- The global burden of disease. 2004 update. http://www.who.int/healthinfo/global_burden_disease/2004_report_update/en/index.html (accessed 12 Mar 2013).
- Barnes PJ. Chronic obstructive pulmonary disease: new opportunities for drug development. *Trends Pharmacol Sci* 1998;19:415–23.
- Underwood DC, Osborn RR, Bochnowicz S, et al. SB 239063, a p38 MAPK inhibitor, reduces neutrophilia, inflammatory cytokines, MMP-9 and fibrosis in lung. *Am J Physiol Lung Cell Mol Physiol* 2000;279:L895–902.
- Renda T, Baraldo S, Pelaia G, et al. Increased activation of p38 MAPK in COPD. *Eur Respir J* 2008;31:62–9.
- Kilty IC, Lucas J, Rhodes J, et al. Do p38 alpha inhibitors represent a potential anti-inflammatory treatment for COPD? *European Respiratory Society Lung Inflammation Conference*; May 2005, Taormina, Sicily: European Respiratory Society, 2005.
- Xing L, Shieh HS, Selness SR, et al. Structural bioinformatics-based prediction of exceptional selectivity of p38 MAP kinase inhibitor PH-797804. *Biochemistry* 2009;48:6402–11.
- Hope HR, Anderson GD, Burnette BL, et al. Anti-inflammatory properties of a novel N-phenyl pyridinone inhibitor of p38 mitogen-activated protein kinase: preclinical-to-clinical translation. *J Pharmacol Exp Ther* 2009;331:882–95.
- GOLD guidelines. (2010 update). http://www.goldcopd.org/uploads/users/files/GOLDReport_April112011.pdf (accessed 12 Mar 2013).
- MacNee W, Allan R, Jones I, et al. A randomized, placebo controlled trial of 6 weeks' treatment with a novel oral p38 inhibitor in patients with COPD. *Eur Respir J* 2010;36(Suppl 54):P4000.
- Mahler DA, Weinberg DH, Wells CK, et al. The measurement of dyspnoea. Contents, interobserver agreement, and physiologic correlates of two new clinical indexes. *Chest* 1984;85:751–8.
- Mahler DA, Witek TJ. The MCID of the transition dyspnoea index is a total score of one unit. *COPD* 2005;2:99–103.
- Miller MR, Hankinson J, Brusasco V, et al. ATS/ERS Task Force: standardisation of lung function testing. Standardisation of spirometry. *Eur Respir J* 2005;26:319–38.
- FDA. Guidance for the use of Bayesian statistics in medical device clinical trials. 2010. <http://www.fda.gov/medicaldevices/deviceregulationandguidance/guidancedocuments/ucm071072.htm> (accessed 12 Mar 2013).

- 16 Smith MK, Jones I, Morris MF, *et al.* Implementation of a Bayesian adaptive design in a proof of concept study. *Pharm Stat* 2006;5:39–50.
- 17 Fabbri LM, Calverley PMA, Izquierdo-Alonso JL, *et al.* M2–127 and M2–128 study groups. Roflumilast in moderate-to-severe chronic obstructive pulmonary disease treated with longacting bronchodilators: two randomised clinical trials. *Lancet* 2009;374:695–703.
- 18 Lunn DJ, Thomas A, Best N, *et al.* WinBUGS—a Bayesian modelling framework: concepts, structure, and extensibility. *Stat Comput* 2000;10:325–37.
- 19 Brown H, Prescott R. *Applied mixed models in medicine*. 2nd edn. New York: John Wiley & Sons, Ltd, 2006.
- 20 Cazzola M, MacNee W, Martinez FJ, *et al.* American Thoracic Society, European Respiratory Society Task Force on Outcomes of COPD. Outcomes for COPD pharmacological trials: from lung function to biomarkers. *Eur Respir J* 2008;31:416–69.
- 21 Lomas DA, Lipson DA, Miller BE, *et al.* An oral inhibitor of p38 MAP kinase reduces plasma fibrinogen in patients with chronic obstructive pulmonary disease. *J Clin Pharmacol* 2012;52:416–24.
- 22 Suissa S, Ernst P, Vandemheen KL, *et al.* Methodological issues in therapeutic trials of COPD. *Eur Respir J* 2008;31:927–33.
- 23 Hanania NA, Darken P, Horstman D, *et al.* The efficacy and safety of fluticasone propionate (250 µg)/salmeterol (50 µg) combined in the Diskus inhaler for the treatment of COPD. *Chest* 2003;124:834–43.
- 24 Rabe KF, Bateman ED, O'Donnell D, *et al.* Roflumilast—an oral anti-inflammatory treatment for chronic obstructive pulmonary disease: a randomised controlled trial. *Lancet* 2005;366:563–71.
- 25 Calverley PMA, Rabe KF, Goehring UM, *et al.* Roflumilast in symptomatic chronic obstructive pulmonary disease: two randomised clinical trials. *Lancet* 2009;374:685–94.
- 26 Calverley P, Pauwels R, Vestbo J, *et al.* Trial of Inhaled Steroids and Long-acting beta2 Agonists Study Group. Combined salmeterol and fluticasone in the treatment of chronic obstructive pulmonary disease: a randomised controlled trial. *Lancet* 2003;361:449–56.
- 27 Tashkin DP, Celli B, Senn S, *et al.* A 4-year trial of tiotropium in chronic obstructive pulmonary disease. *N Engl J Med* 2008;359:1543–54.
- 28 Cohen SB, Cheng TT, Chindalore V, *et al.* Evaluation of the efficacy and safety of pamapimod, a p38 MAP kinase inhibitor, in a double-blind, methotrexate-controlled study of patients with active rheumatoid arthritis. *Arthritis Rheum* 2009;60:335–44.
- 29 Damjanov N, Kauffman RS, Spencer-Green GT. Efficacy, pharmacodynamics, and safety of VX-702, a novel p38 MAPK inhibitor, in rheumatoid arthritis: results of two randomized, double-blind, placebo-controlled clinical studies. *Arthritis Rheum* 2009;60:1232–41.