IMAGES IN THORAX

An unusual case of haemoptysis

Jovin John Mathew,¹ Nicholas Paul Hollings,² Jonathan Daniel Myers¹

¹Respiratory Medicine, Royal Cornwall Hospital, Truro, Cornwall, UK

²Department of Radiology, Royal Cornwall Hospital, Truro, Cornwall, UK

Correspondence to

Dr Jovin John Mathew,
Respiratory Medicine, Royal
Cornwall Hospital, Truro,
Cornwall TR1 3LJ, UK;
drjjmathew@gmail.com

Received 14 December 2012

Accepted 17 December 2012

Published Online First

24 January 2013

An 18-year-old female subject with a history of decortication for right sided, postpneumonic pleural

Figure 1 Axial maximum intensity projection image from CT pulmonary angiogram showing hypoplastic pulmonary artery (black arrow) and prominent bronchial artery (white arrow) on the right side.

thickening presented with haemoptysis. A CT pulmonary angiogram showed marked hypoplasia of the right main, lobar and segmental pulmonary arteries, along with reduced volume of the right lung. Prominent bronchial artery supply to this lung was noted (figure 1) with absence of pulmonary veins. Perfusion scintigraphy showed only 4% perfusion to the right lung and normal perfusion of the left lung (figure 2). Pulmonary artery hypoplasia has been reported to be associated with ipsilateral pulmonary venous atresia.¹ It can present with symptoms mimicking pulmonary embolism.²

Learning points

- ▶ Pulmonary artery hypoplasia is a rare cause of haemoptysis. The source of bleeding is usually from hypertrophic bronchial arteries.
- ▶ CT pulmonary angiogram combined with perfusion scintigraphy is very useful in making the diagnosis non-invasively.

Figure 2 Perfusion scintigraphic images demonstrating scant perfusion to the right lung.

To cite: Mathew JJ,
Hollings NP, Myers JD.
Thorax 2013;**68**:796–797.

Acknowledgements Mr Ivor Dixon, Medical Physicist, Royal Cornwall Hospital, Truro, Cornwall, UK for providing the new figure.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; internally peer reviewed.

REFERENCES

- 1 Heyneman LE, Nolan RL, Harrison JK, *et al.* Congenital unilateral pulmonary vein atresia: radiologic findings in three adult patients. *AJR Am J Roentgenol* 2001;177:681–5.
- 2 Vohra N, Alvarez M, Abramson AF, *et al.* Hypoplastic pulmonary artery: an unusual entity mimicking pulmonary embolism during pregnancy. *Obstet Gynecol* 1992;80(3 Pt 2):483–5.