Abstract P78 Table 1. Results of the Asthma and COPD Reviews

Exacerbations in previous 12 months	Asthma	COPD	
	Oral antibiotics and/or oral steroids per patient	2.2 per patient	3.1 per patient
Number of A&E attendances (n)	14	33	
Number of hospital admissions (n)	5	31	
Inhaler technique	Good technique (n)	21% (11)	42% (13)
	Moderate technique	19% (10)	29% (9)
	Poor technique	60% (31)	29% (9)
Beclomethasone dipropionate equivalence	pre review	1656.6mcg	NA
	post review	1000.0mcg	NA
Smoking history (n)	Current	26.4% (14)	29.0% (9)
	Agreed to stop following review	78.6% (11)	77.8% (7)
	Ex smoker	17.0% (9)	71.0% (22)
	Mean pack year history	29.4	37.4
Adherence to medicines in previous 12 months (mean)	Maintenance	6.4 inhalers	10.3 inhalers
	ICS/LABA or ICS		
	LAMA	NA	10.8 inhalers
On correct therapy based on symptoms, diagnosis and disease severity (n)	Reliever inhaler	8.3 inhalers	8.7 inhalers
		26.9% (14)	45.2% (14)
If not on correct therapy, intervention made (n)	Step down	59.6% (31)	NA
	Stopping a part of treatment	3.8% (2)	25.8% (8)
	Querying the correct diagnosis	5.8% (3)	22.6% (7)
	Change drug class	NA	9.7% (3)
FEV1 % of predicted mean		NA	52.2%
ACT score mean		16.9	
CAT score mean		21.5	

had asthma (n = 53) with 37% COPD (n = 31), with a mean age 56.3 and 67.0 years respectively.

Table 1 shows a significant proportion of patients had uncontrolled disease based on their quality of life scores (QoL), use of reliever inhalers and frequency of exacerbations. Despite use of high dose medicines, this was likely to be due to poor adherence to maintenance inhalers and poor inhaler technique in a large proportion of patients. Interventions were made in most patients, including reducing the beclomethasone dipropionate (BDP) equivalence in the asthma group by 60.4%. Follow up showed that despite a reduction in inhaler therapy, QoL, peak flow measurements and reliever use all improved.

Conclusions Tailored reviews by the pharmacist resulted in significant interventions that improved QoL, adherence to therapy, reduced unnecessary over prescribing of high dose medicines and resulted in a large proportion of patients to successfully stop smoking.

From screening to treatment of children with chronic lung disease

P79 THE IMPACT OF SOCIAL DEPRIVATION ON CLINICAL OUTCOMES IN CHILDREN WITH CYSTIC FIBROSIS (CF) IN A DEPRIVED AREA OF SCOTLAND

J Barge, Y Cheong, L Thomson, NA Gibson, PL Davies; *Royal Hospital for Sick Children, Glasgow, UK*

10.1136/thoraxjnl-2013-204457.229

Introduction Lung function tests and BMI are widely-used outcome measures in children with CF, and their preservation requires adherence to exacting treatment regimens.

We hypothesised that deprivation might affect the adherence to therapy of children in a deprived area of Scotland, thus reducing BMI and FEV₁ scores and increasing clinical input required.

Schechter (2003) demonstrated a link between deprivation and mortality in children with CF, although the mechanism remained unclear, and the study has not been repeated elsewhere. In an adult CF population in an affluent area of England, Jarad (2005) found no correlation between deprivation and FEV₁. This question has not previously been examined in a deprived paediatric population.

Method In February 2013, RHSC Glasgow had 95 children (4946) with CF under its care. All children over 2 years had their BMI centile calculated. In all children over four years, mean FEV1% predicted was calculated. The number of outpatient clinics and inpatient admissions over a three year period was noted for all children over three years old.

The Scottish Index of Multiple Deprivation (SIMD) ranks the 6505 postcode districts in Scotland in order of deprivation, and a rank was obtained for each child's postcode.

SMID rank was plotted against BMI, mean FEV1% predicted, outpatient appointments and inpatient admissions, with any correlations noted.

Result 83 children had BMI centile recorded (mean 45.4, range 0.4–98), 66 children had FEV1 % predicted calculated (mean 92.4%, range 50.4–146%). Mean SMID ranking was 2763 (range 33–6499.). The mean number of respiratory outpatient clinics attended was 23.1 (range 18–33), total outpatient clinics attended was 25.8 (range 18–48) and inpatient admissions was 4.3 (range 0–20) over three years.

No correlation was shown between SMID rank and BMI centile ($r^2 = 0.0025$), mean FEV1% predicted ($r^2 = 0.0025$), respiratory clinics attended ($r^2 = 0.0121$), total outpatients attended ($r^2 = 0.0263$) and inpatient admissions ($r^2 = 0.0078$).

Conclusion Relative deprivation does not correlate with BMI centile or FEV1% predicted in this population of CF children and is not associated with the level of clinical input required.

P80 ANTIBIOTIC USAGE AS A RISK FACTOR FOR NON-TUBERCULOUS MYCOBACTERIUM INFECTION IN CHILDREN WITH CYSTIC FIBROSIS

¹JL Bowker, ²MF Thomas, ¹CJ O'Brien.³; ¹Newcastle University, Newcastle, England; ²Great North Children's Hospital, Newcastle, England

10.1136/thoraxjnl-2013-204457.230

Background There is a growing incidence of Non-Tuberculous Mycobacterium (NTM) in children with Cystic Fibrosis. This audit aimed to determine relevant risk factors for NTM acquisition in paediatric cystic fibrosis patients.

Methods Data was obtained from clinical notes for 66 patients aged 5–18 years old in a North-East CF clinic who had sputum or bronchoalveolar lavage screened for NTM between 2011 and 2012.