Poster sessions

“irresistible” (SL1V)) and adequacy of CPAP compliance. They were also asked about their use of objective tests.

Results 178 respondents completed the survey. There was poor agreement among clinicians regarding the presence of residual drowsiness (McNemar's test, figure-1). In response to the DVLA question about “excessive” drowsiness, the patient had a 1 in 2.57 (range-1.12 to 5.66), and about “irresistible” drowsiness, a 1 in 1.32 (range- 1.04 to 2.84), chance of being given a different answer depending on the clinician seen. Furthermore in each vignette the same clinician was more likely to say “yes” to “excessive” than to “irresistible” (71 +/-12% v/s 42 +/-10%, P=0.0045). There was also a lack of consensus as to what constitutes “adequate CPAP compliance”. Across the vignettes there was minimum of 1 in 1.7 and a maximum of 1 in 6.7 chance of disagreement amongst the clinicians (median- 3, range 1.7 to 6.7). 1% of clinicians always and 4% frequently use objective tests to help in their assessment. They are more likely to use in professional drivers as compared to non professional drivers (52% v/s 38%, P=0.0002, OR=1.75). Tests used were MSLT (34%), OSLER (29%), MWT (28%) and divided attention driving simulator (9%).

Conclusions The information that the DVLA is given may vary markedly depending upon which clinician completes the form. Furthermore the same clinician may give a different impression to the DVLA depending on which form they are asked to complete. Objective testing is not undertaken routinely. Better guidance and better objective tests are needed to ensure consistency of the information that the DVLA is given.

P253 OBSTRUCTIVE SLEEP APNOEA SYNDROME: PATIENTS' EXPERIENCE OF THE DRIVER AND VEHICLE LICENSING AGENCY

M Mason, N Oscroft, M Davies, T Quinnell, I Smith; *Respiratory Support and Sleep Centre, Papworth Hospital NHS Foundation Trust, Papworth Everard, Cambridge, UK*

10.1136/thoraxjnl-2013-204457.405

Introduction Driver's licence holders have a legal obligation to inform the Driver and Vehicle Licensing Authority (DVLA) if diagnosed with Obstructive Sleep Apnoea Syndrome (OSAS). This requirement may cause considerable anxiety but few data are available to advise patients on the likely outcomes following such notification.

Methods Patients diagnosed with OSAS and offered CPAP between 1st October 2009 and 31st March 2010 were surveyed

Abstract P253 Table 1. Characteristics of the patient population sample

Characteristic	
Mean Age years (range)	54 (31-82)
Gender (male/female)	2.7:1
BMI (kg/m ²)	38 (9)
ESS at diagnosis	13 (5)
ESS post treatment	7 (5)
ODI at diagnosis	21 (20)
ODI post treatment	4 (12)
Compliance hours/night (measured by CPAP device)	5.8 (2)
Compliance with CPAP treatment >3.5 hours/night	72%
Recorded evidence in medical notes of advice regarding the DVLA	79%

BMI, body mass index; ESS, Epworth Sleepiness Score; ODI, oxygen desaturation index (4% drop in saturation/hour); CPAP, continuous positive airway pressure
Data presented as mean (SD) unless specified

anonymously and asked to report on: a) symptoms of sleepiness whilst driving before and after treatment; b) if they recalled being given advice about contacting the DVLA by healthcare professionals; c) whether they contacted the DVLA; d) the response of the DVLA if notified. The survey was performed between 1st June and 31st August 2010. To provide a description of the population surveyed 67 (10%) case notes were chosen at random from the population surveyed, reviewed and descriptive data extracted.

Results Six hundred and seventy three patients were surveyed and 297 (44%) responded. 92% were category B licence holders. The data outlining the surveyed population are in Table 1.

Sixty percent and 16% respectively reported no and moderate to severe sleepiness whilst driving, prior to treatment. Two hundred and six patients (69%) recalled being given advice about driving by a health care professional and of those 161 patients (78%) had informed the DVLA of their diagnosis. In total 197 patients (66%) had informed the DVLA of their OSAS. The DVLA asked 8% (16/197) to stop driving temporarily of which 80/197 contacted the DVLA prior to starting treatment. Five patients (2.5%) were deemed not fit to drive by the DVLA in the long term. The mean (SD) time for the DVLA to reach a decision was 29 (33) days.

Conclusion In summary, recognising its limits, in particular the risk of responder bias, this survey shows most OSAS patients offered CPAP do not experience problems with driving licence retention if they contact the DVLA. The DVLA infrequently ask patients to stop driving. Approximately one third of patients had not informed the DVLA of their OSAS during the timeframe of this survey.

P254 PREDICTING WHO NEEDS A HUMIDIFIER WITH CPAP

L Thiruganasothy, J Littlemore, J Oliver, P Close, SD West; *Freeman Hospital, Newcastle-upon-tyne, England*

10.1136/thoraxjnl-2013-204457.406

Introduction Some patients using continuous positive airway pressure (CPAP) for the treatment of Obstructive Sleep Apnoea (OSA) require the addition of a heated humidifier to their CPAP machine. This is used to combat symptoms of dry nose and mouth and blocked nose which may limit CPAP adherence. We sought to establish whether there are any patient predictors for whether a humidifier will subsequently be needed. This might allow provision of more economical integrated CPAP/humidifiers.

Methods All patients commencing CPAP over a one year period completed a questionnaire with a member of the Sleep team at the time of CPAP set-up. The questionnaire asked about symptoms prior to CPAP including blocked, dry or runny nose and dry mouth on waking. Details of sleep study parameters such as ODI and AHI were obtained from hospital notes, along with anthropometric measures of body mass index (BMI) and neck size. The questionnaire also enquired about previous medical history (including ENT surgery), medications, smoking history and bedroom environment. Patients were given humidifiers according to usual practice as required after CPAP commencement, according to symptoms. The CPAP database was reviewed at the time of analysis to determine which patients had received humidifiers.

Results Questionnaires were completed by 185 people commencing CPAP from January 2012. The mean (SD) age of this group was 53 years (11.7), BMI 36.6 kg/m²(7.1), neck size 43.9cm (4.4), ODI 29.3 (23). The proportions of different severities of OSA were 19% mild, 34% moderate and 47% severe. The frequencies of symptoms prior to CPAP were 87% dry mouth, 54% blocked nose, 40% dry nose and 22% runny nose.