Conclusions Findings show that greatest improvements in terms of exercise capacity may be seen from 8 week pulmonary rehabilitation programmes, but that improvement for the ISWT can be obtained from 6, 7 or 8 week programmes.

REFERENCES

National Institute for Clinical Excellence. Commissioning a pulmonary rehabilitation service for patients with COPD. 2010. [http://www.nice.org.uk/usingguidance/commissioningguides/pulmonaryrehabilitationforpatientswithcopd/commissioning.jsp] Accessed 13/07/13

P117 PULMONARY REHABILITATION IN THE EAST OF ENGLAND – 2.5-FOLD VARIATION IN COMPLETION RATES

¹L Jongepier, ²R Barlow. ¹EoE Respiratory Strategic Clinical Network, Cambridge, UK; ²PROVIDE, Chelmsford, UK

10.1136/thoraxjnl-2014-206260.258

Introduction and objectives Pulmonary Rehabilitation (PR) should be made available to all suitable people with COPD and various other chronic respiratory conditions.¹ An abundance of guidelines has been produced making recommendation on the quality of both the provision and commissioning of PR. Limited data is available on PR programme adherence rates and most study rates post COPD exacerbation. Recent IMPRESS Guidance suggested a target completion rate of 75% of offered sessions and stated the national average being less than 50%.² Our aim was to get an accurate regional perspective of completion rates of all PR service providers to use as a lever for improvement.

Methods Prior to 2013/14 a regional PR group was formed to promote best practice, offer peer support and enable improvements through the collection of meaningful regional data. A data set was agreed and defined and during 2013/14 quarterly data was collected from 17/18 (95%) providers across all CCGs. The number of PR starters and completers (attending a minimum of 75% of the offered sessions) was collected. Data from all providers was compared.

Results During 2013/14 4,737 patients started PR and 2935 patients attended at least 75% of the offered sessions. The average completion rate across the EoE was 62% with programmes varying more than 2.5-fold from 37% to 91%. 24% of providers had a completion rate of $\leq 50\%$, however another 24% had a completion rate of $\geq 75\%$.

Conclusions Completion rate varied widely with a 2.5-fold variation between the best and worst performers. 24% had a completion rate of $\leq 50\%$. The regional average of 62% was below the suggested target rate of 75%. The reasons for this variation are not known, but will be investigated by the EoE PR group.

Abstract P117 Figure 1

REFERENCES

- NICE. COPD: management of COPD in adults in primary and secondary care. 2010 [http://www.nice.org.uk/guidance/CG101]
- Impress (BTS and the primary care respiratory society – UK). Impress Guide to Pulmonary Rehabilitation. 2011. [www.impressresp.com/index.php?option=com_docam&task=doc_view&grid=41&Itemid=82] Accessed on 07/07/13

P118 QUANTITY AND QUALITY OF REFERRALS TO PULMONARY REHABILITATION FROM PRIMARY CARE FOLLOWING INCLUSION IN THE QUALITY AND OUTCOMES FRAMEWORK (QOF) IN WALES

¹J Ayling-Smith, ²M Owen, ¹J Byers, ¹D Menzies. ¹Glan Clwyd Hospital, Betsi Cadwaladr Health Board, UK; ²Wrexham Maelor Hospital, Betsi Cadwaladr Health Board, UK

10.1136/thoraxjnl-2014-206260.259

Introduction Referring suitable patients for Pulmonary Rehabilitation (PR) has formed part of the Quality and Outcomes Framework (QOF) reward for General Practitioners (GPs) in Wales since April 2013. Patients with Chronic Obstructive Pulmonary Disease (COPD) that have a Medical Research Council (MRC) dyspnoea score of 3 or above, or those with an MRC score of 2 and recently discharged from hospital for COPD are eligible. We sought to determine the impact of this change on the referral pattern to our PR programme, which traditionally had only accepted referrals from secondary care respiratory consultants.

Methods A standardised form was prospectively designed and distributed to enable primary care providers to refer suitable patients to the PR programme. Data on patient demographics, respiratory diagnosis, co-morbid conditions, MRC dyspnoea score, exacerbation frequency and current medication were collected. Referring GPs were also asked to provide recent spirometry values for patients that were referred. An assessment was made based on the information provided as to whether the patient could be listed immediately for pulmonary rehabilitation, whether further assessment was required, or whether the referral was inappropriate.

Results A sample of 250 GP referral forms out of a total of 545 were evaluated of which 51% of patients were male and the mean age was 69 years (range 31–90). 40.8% of GP referrals could be listed immediately for PR, 34.8% of referrals needed further assessment and 24.4% were inappropriate. 22.4% of all patients referred did not have COPD based on the spirometry results supplied. Compared with the baseline referral rate to PR from secondary care consultants, the rate of referral from GPs

Abstract P118 Figure 1 Referrals to pulmonary rehabilitation before and after QoF