Abstract P120 Table 1

Shared Decision Making Descriptor	Result
% of patients felt that the clinician addressed what was important to them at their appointment	90%
% of patients felt that their expectations were met during their appointment	90%
% of patients felt able to make a decision that was right for them with their clinician	90%
% of patients felt that the clinician shared their expertise with them enough to help the patient feel that they were making the right choice for them	90%
% of patients felt that they fully understood the pros and cons of each treatment option	88%
% of patients were happy that there was enough time to help them feel confident in making their treatment choice	90%
% of patients reported that the information they received led to them changing their decision regarding treatment choices	45%
% of patients felt more confident to manage their condition after attending an education and self-management group	90%
% of patients reporting that they now do things differently as a result of their consultations showing changes in lifestyle and health behaviours	95%
% of patients reporting high confidence scores in self-managing their condition at the start of the group	5%
% of patients reporting high confidence scores in self-managing their condition at the end of the group.	90%

shared decision making within group education for COPD patients to date.

Methods 20 semi-structured interviews were performed to obtain quantitative and qualitative data from COPD patients who had recently attended an education and supported self-management group held over six weeks. Data collection was performed by allied health professionals who do not work in the COPD clinic. Questionnaires were reviewed and amended by a Questionnaire Users, Interviews and Surveys group prior to use.

Results (see Table) Qualitative feedback provided by patients supported the quantitative results and ranged from neutral to highly positive in nature, with several patients reporting significant impact on their quality of life, confidence in supported self-management, increased exercise participation, physical function, and social participation.

Conclusion COPD patients attending a six weeks education and supported self-management group reported significant understanding of information, increased understanding of treatment options, and increased education and ability to self-manage.

REFERENCES

- Collins A (2011) The Kings Fund Report: 'Making shared decision-making a reality: No decision about me, without me'
- Coulter A (2009) 'Implementing Shared Decision Making in the UK': A Report for The Health Foundation

P121

SPEECH AND LANGUAGE THERAPY IN PULMONARY REHABILITATION: THE IMPLICATION OF EDUCATION SESSIONS ON DYSPHAGIA MANAGEMENT

SF Lillie, J Haines, A Vyas, SJ Fowler. *Lancashire Teaching Hospitals Trust, Preston, UK*

10.1136/thoraxjnl-2014-206260.262

Introduction Pulmonary rehabilitation (PR) programs use multi-disciplinary teams to optimise physical and social functioning of patients with chronic respiratory impairment. Such patients demonstrate an increased prevalence of oropharyngeal dysphagia as a consequence of impaired co-ordination between respiration and swallowing function. Often patients will not be aware of the warning signs of dysphagia and unfortunately will not be seen by a speech and language therapist until they are admitted to hospital. We report the outcomes of a pilot scheme whereby such patients underwent education, assessment and treatment for dysphagia as part of their PR programme..

Methods The pilot scheme ran between June 2013 and May 2014. Intervention consisted of: (1) a one hour group education

session on the signs, symptoms and risks of dysphagia; (2) screening for oropharyngeal dysphagia; and (3) individual outpatient management in Airways Clinic. The majority of patients attending the education sessions had a diagnosis of Chronic Obstructive Pulmonary Disease (COPD).

Results The education programme was delivered to 72 patients, and resulted in a significant improvement in dysphagia knowledge. The average score pre education was 3/11 and post education was 8/11. Fourteen patients (19%) exhibited or reported symptoms of dysphagia. Of these two patients were overtly aspirating and required food/fluid modification and seven patient's required instrumental assessment in the form of fibre endoscopic evaluation of swallowing (FEES). During FEES, three patients showed penetration of food/ fluids and were at risk of silent aspiration. These patients attended for further SLT where diet/ fluids were modified, posture was assessed and dysphagia therapy was introduced.

Conclusions Dysphagia education and management of patients in PR can contribute the early identification, patient awareness and self-management of dysphagia. We have confirmed that undiagnosed but clinically important dysphagia is present in patients undergoing PR. We are investigating whether improved dysphagia knowledge and early identification of dysphagia symptoms leads to reduced exacerbations and improved quality of life.

P122

A SURVEY OF PULMONARY REHABILITATION (PR) SERVICES IN KENT, SURREY, SUSSEX (KSS)

J-P Crofton-Biwer, E Lazar, J Bott. *Kent Surrey Sussex Academic Health Science Network, Crawley, UK*

10.1136/thoraxjnl-2014-206260.263

Introduction and objectives There is no agreed model for Pulmonary Rehabilitation (PR) and wide variation in services exists. A regional PR network was established 4 years ago, with the aim to drive up standards and reduce variation. An audit was undertaken of all PR services in the region to determine costs of services and factors influencing variance.

Method In June 2013 e-questionnaires were sent to all 16 known PR providers; fifty questions requested average annual/ weekly data including: staff pay bands, time spent on exercise, education, administration, travel and other identifiable costs, numbers failing to complete (drop-out) and clinical outcomes. All costs were calculated in terms of cost-per-patient. Providers