all patients with a *DNAH11* defect ($n = 5$) compared to healthy controls ($n = 3$) and patients with PCD due to a defect of the central pair or nexin link ($n = 3$) (Figure 1). A reduction in outer dynein arm volume of 30% was identified compared to healthy and PCD controls.

In conclusion, a mutation in *DNAH11* results in a subtle abnormality in ultrastructure. The defects are specific to the 'forearm' of the outer dynein arm and only detectable at the base of the cilium where *DNAH11* is located. Electron tomography is highly effective in visualising this defect.

S89 IDENTIFICATION OF POTENTIALLY PATHOGENIC MICROORGANISMS BY SELECTED ION FLOW TUBE – MASS SPECTROMETRY (SIFT-MS)

¹JM Jackson, ²VL Kim, ³AC Tuck, ¹SA Wootton, ⁴TM Wilkinson. ¹NIHR Southampton Biomedical Research Centre, Southampton, UK; ²NIHR Southampton Respiratory Biomedical Research Unit, Southampton, UK; ³Infectious Disease Epidemiology Group, Sir Henry Wellcome Laboratories, Academic Unit of Clinical and experimental Sciences, Faculty of Medicine, University of Southampton, Southampton, UK; ⁴University Hospital Southampton NHS Foundation Trust, Southampton, UK

10.1136/thoraxjnl-2014-206260.95

Introduction Improved targeted therapy and reduced clinical burden from respiratory bacterial infection could result from early identification of specific Potential Pathogenic Microorganisms (PPM). Sputum culture for identification usually requires several days. SIFT-MS has the potential to reduce the required incubation time by sampling the culture headspace to generate an ionic spectrum from volatile organic compounds that may be characteristic of the PPM. Additionally, these signatures may be detectable in breath taken directly from patients. This pilot study investigates the potential of SIFT-MS to identify 5PPMs, incubated separately for 24 hr.

Methods Training set: *Haemophilus influenzae* (HI), *Moraxella catarrhalis* (MC), *Streptococcus pneumoniae* (SP), *Staphylococcus aureus* (SA) and *Pseudomonas aeruginosa* (PA) cultures and a negative control incubated at 37°C on chocolate agar plates in sealed bags for 24 hr, 48 hr and 72 hr. Plates were opened for 10 min at room temperature before ionic spectra of the gas above the culture dishes in the range 15 to 200 mass units were recorded using SIFT-MS and standardised to operating conditions. Test set: the same five PPMs and a negative control were incubated in triplicate for 24 hr only and analysed as above.

Results Using the spectra generated with H₃O⁺ ionisation, 6 ion sets were identified. The sum of ions within each set, expressed as a percentage of the total ion sum of masses 15 to 200 (excluding reagent ions) fell into ranges that, in combination, differentiated between the PPMs. This set of conditions was incorporated into an algorithm that was then applied to the test set of triplicate plates. The algorithm correctly differentiated all 24 hr plates with MC, SP, SA and PA from each other and from the negative control and HI plates with 100% accuracy. Negative control and HI could not be differentiated.

Conclusion This pilot study illustrates the potential for SIFT-MS to identify monocultures of 4 common PPMs within a short incubation time and encourages further study with a wider range of pathogens alone and in combination. Early identification of PPMs in culture, and translation to potentially detect carriage or infection with specific pathogens in breath may improve management of respiratory infections.

S90 ROLE FOR IL-1ALPHA IN VIRAL-INDUCED INFLAMMATORY RESPONSES IN A CO-CULTURE MODEL OF THE AIRWAY MUCOSA

¹A Hill, ¹L Tezera, ¹C Blume, ²C Grainge, ¹DE Davies, ¹EJ Swindle. ¹University of Southampton (Clinical and Experimental Sciences, Faculty of Medicine), Southampton, UK; ²University of Newcastle (School of Medicine and Public Health), Newcastle, Australia

10.1136/thoraxjnl-2014-206260.96

Introduction Asthma is an inflammatory disease of the conducting airways which is exacerbated by environmental exposures, such as viral infections. Bronchial epithelial cells (BECs) together with underlying fibroblasts form an epithelial mesenchymal trophic unit (EMTU) that maintains normal tissue homeostasis. In asthma the EMTU is dysregulated. Recent evidence suggests that viral infections activate the epithelial barrier resulting in mediator release which could potentially activate fibroblasts. Therefore, we hypothesised that exposure to viruses activates inflammatory and anti-viral responses in the EMTU.

Methods The EMTU was modelled using a co-culture system of polarised BECs (16HBE14o-) and fibroblasts (MRC5s) maintained on the apical and basolateral surface of a nanoporous membrane respectively. After 6 days the model was challenged apically with poly (I:C) (a viral mimetic) and barrier responses were determined by measuring transepithelial resistance (TER) while cytokine release was determined by ELISA.

Results Following poly (I:C) stimulation a significant reduction in TER was observed in both the EMTU model and BEC monocultures. However, the EMTU model maintained a higher TER at 6–24 h after challenge. With regards to cytokine secretion, poly (I:C) stimulation significantly induced pro-inflammatory (IL-6, IL-8, GM-CSF and IL-1 α) and anti-viral (IP-10) mediator release from BEC but not fibroblast monocultures. In the EMTU model, basolateral IL-6, IL-8, GM-CSF and IP-10 responses to poly (I:C) were significantly enhanced compared to BEC monocultures. In addition, basolateral pro-inflammatory (IL-6, IL-8 and GM-CSF) but not antiviral (IP-10) responses to poly (I:C) were abrogated in the EMTU model after pre-incubation with IL-1 receptor antagonist (IL-1ra). Furthermore, direct stimulation with IL-1 α induced IL-6 and IL-8 release in the EMTU model and fibroblast monocultures but not in BEC monocultures.

Conclusions Poly (I:C) activates inflammatory and anti-viral responses in BEC monocultures and fibroblast co-culture models of the EMTU. These responses were enhanced in the co-culture model suggesting that the EMTU is activated. Inflammatory but not anti-viral responses were mediated by epithelial-derived IL-1 α acting on the underlying fibroblasts. This may have important consequences in promotion of inflammation and airway remodelling in viral-induced exacerbations of asthma.

New asthma treatments

S91 ONCE-DAILY TIOTROPIUM RESPIMAT® ADD-ON TO ICS + LABA IMPROVES SYMPTOM CONTROL AND REDUCES EXACERBATIONS IN PATIENTS WITH SYMPTOMATIC ASTHMA

¹D Price, ²M Engel, ²P Moroni-Zentgraf, ²H Schmidt, ³R Dahl, ⁴P Paggiaro, ⁵M Vandewalker, ⁶HAM Kerstjens, ⁷A Kaplan. ¹Institute of Applied Health Sciences, University of Aberdeen, Aberdeen, UK; ²Boehringer Ingelheim Pharma GmbH and Co. KG, Biberach an der Riss, Germany; ³Odense University Hospital, Odense, Denmark; ⁴University of Pisa, Pisa, Italy; ⁵Clinical Research of the Ozarks, Colombia, USA; ⁶University Medical Center Groningen, Groningen, The Netherlands; ⁷Family Physician Airways Group of Canada, Ontario, Canada

10.1136/thoraxjnl-2014-206260.97