IMAGES IN THORAX

Cerebral air embolism during CT-guided lung biopsy

Rafael Dahmer Rocha, André Arantes Azevedo, Priscila Mina Falsarella, Antônio Rahal Jr, Rodrigo Gobbo Garcia

Department of Interventional Radiology, Hospital Israelita Albert Einstein, Avenida Albert Einstein, São Paulo, São Paulo, Brazil

Correspondence to

Dr Rafael Dahmer Rocha, Department of Interventional Radiology, Hospital Israelita Albert Einstein, Avenida Albert Einstein, 627/701 - Morumbi, São Paulo - São Paulo 05652-900, Brazil; rafaeldrocha@gmail.com

Received 21 April 2015

Accepted 14 May 2015

Published Online First

29 May 2015

A patient with cirrhosis and hepatocellular carcinoma, on the waiting list for liver transplantation, presented with lung nodule during initial evaluation. A CT-guided core needle lung biopsy under general anaesthesia was performed. Pneumothorax (figure 1) and a large amount of gas within the left ventricle (figure 2) were identified during the procedure. Neurological focal signs (drowsiness,

conjugate eye deviation and right hemiparesis) were observed in the anaesthesia-recovery period. A brain CT scan was immediately obtained,

Figure 1 CT-guided lung biopsy of the nodule was performed with a 20-gauge needle and the patient in supine position. A small pneumothorax was already seen in this image.

Figure 2 The repeated CT image showed a large amount of gas within the left ventricle.

Figure 3 Axial CT of the brain showed gas locules within the left parietal lobe.

Figure 4 Magnetic resonance image after 5 days revealed the real extension of the embolism. A massive stroke involving both sides of the brain can be seen.

CrossMark

To cite: Rocha RD, Azevedo AA, Falsarella PM, et al. *Thorax* 2015;**70**:1099–1100.

showing cerebral air embolism (figure 3). Initial resuscitation and hyperbaric oxygen therapy were performed. Five days later, MRI revealed a massive stroke (figure 4). At 3-month follow-up, the patient recovered motor functions, however, maintained with important cognitive deficits. Systemic air embolism is a rare but potentially life-threatening complication of percutaneous lung biopsy. The incidence ranges from 0.02% to 0.5%.^{1–3} Coughing during the procedure, cystic or cavitary lesion, positive pressure ventilation and needle-tip placement through a pulmonary vein are the most common predisposing factors. The management involves high-flow 100% oxygen and early hyperbaric oxygen therapy.

Competing interests None declared.

Provenance and peer review Not commissioned; internally peer reviewed.

REFERENCES

- 1 Ishii H, Hiraki T, Gobara H, *et al*. Risk factors for systemic air embolism as a complication of percutaneous CT-guided lung biopsy: multicenter case-control study. *Cardiovasc Intervent Radiol* 2014;37:1312–20.
- 2 Freund MC, Petersen J, Goder KC, *et al*. Systemic air embolism during percutaneous core needle biopsy of the lung: frequency and risk factors. *BMC Pulm Med* 2012;12:2.
- 3 Hare SS, Gupta A, Gonçalves ATC, *et al*. Systemic arterial air embolism after percutaneous lung biopsy. *Clin Radiol* 2011;66:589–96.