



MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton, UK

#### Correspondence to

Professor David Coggon, MRC Lifecourse Epidemiology Unit, Southampton General Hospital, Southampton SO16 6YD, UK; dnc@mrc.soton.ac.uk

Received 1 August 2014

Revised 15 August 2014

Accepted 20 August 2014

Published Online First

15 October 2014

## OPINION

# Pneumococcal vaccination for welders

David Coggon, E Clare Harris, Vanessa Cox, Keith T Palmer

Over the past 20 years, evidence has accumulated that occupations entailing exposure to metal fume (principally welders) are at increased risk of developing and dying from pneumococcal and unspecified lobar pneumonia.<sup>1–9</sup> The hazard is apparent in people working with ferrous metal,<sup>2–4</sup> may extend to other metals<sup>4</sup> and is reversible following cessation of exposure.<sup>1,4,5</sup> There are indications that the ultra-fine particles in welding fume promote adherence of pneumococci to bronchial epithelium,<sup>10</sup> but the underlying mechanisms are not yet well established. Nevertheless, in 2011, the independent Joint Committee on Vaccination at the English Department of Health (DH) judged that the evidence of increased risk was sufficient to recommend that employers offer welders vaccination against pneumococcus.<sup>11</sup> This was a new measure that had not previously been applied in other countries.

Subsequently, on the advice of the UK Health and Safety Executive (HSE), the DH weakened its recommendation, stating only that vaccination ‘should be considered for those at risk of frequent or continuous occupational exposure to metal fume (e.g. welders), taking into account the exposure control measures in place’.<sup>12</sup> Then after some delay, HSE published guidance for employers in which they were advised to prioritise controls on exposure and to take the efficacy of such controls into account when deciding whether or not to offer vaccination.<sup>13</sup> The guidance emphasised that vaccination was not a regulatory requirement, provided that an adequate risk assessment could demonstrate that fume was effectively controlled.

A fundamental flaw of this approach is that, as the HSE’s guidance acknowledges, ‘there are no research studies which specify how much exposure to welding or metal fume (dose) will result in an increased susceptibility to pneumonia (response)’.<sup>13</sup> It follows that there is no evidential basis for determining whether exposures are sufficiently controlled to eliminate the risk. When challenged on the matter, HSE responded that despite the admitted absence of evidence, ‘we do know what constitutes effective controls in relation to welding and those are intended to prevent all health-related end-points’ (J Delic, personal communication). They also suggested that exposures should have reduced over time, although they had no supporting empirical evidence.

New data have now become available on mortality by occupation in England and Wales during 2001–2010, and we used them to explore whether the excess risk of pneumonia in welders had persisted into the new century.

The Office for National Statistics provided us with the numbers of deaths in England and Wales

during 2001–2010 at ages 16–74 years, broken down by combinations of sex, age (in 5-year bands), occupation (in 143 categories), social class (eight categories) and underlying cause of death (229 categories). All of this information came originally from death certificates. The data were used to calculate proportional mortality ratios (PMRs) for pneumococcal and unspecified lobar pneumonia (categories J13 and J18.1 in the 10th revision of the International Classification of Diseases) in welders, standardised for age and social class. CIs for PMRs were based on the Poisson distribution. Results were compared with those derived previously by a similar method for the period 1991–2000 when pneumococcal and unspecified lobar pneumonia was classified to a single category (481) in the ninth revision of the International Classification of Diseases.<sup>14</sup>

During 2001–2010, there were 2863 male deaths from pneumococcal and unspecified pneumonia at ages 16–74 in all occupations combined compared with 3234 during 1991–2000. The corresponding numbers of deaths from all causes were 927 152 (including 8160 in welders) during 2001–2010 and 1 202 888 (10 455 in welders) during 1991–2000. As in 1991–2000, mortality from pneumococcal and unspecified lobar pneumonia during 2001–2010 was significantly elevated in welders of working age (table 1), with a particularly high PMR for pneumococcal pneumonia specifically (3.58, 95% CI 1.44 to 7.37). The estimated excess of deaths from pneumococcal and unspecified lobar during 2001–2010 was 10.9 compared with 18.8 during 1991–2000.

Our analysis indicates that welders continued to experience significant excess mortality from pneumococcal and unspecified lobar pneumonia during 2001–2010. As discussed in relation to earlier similar analyses, the excess is unlikely to be explained by bias or confounding.<sup>1–3,5,8</sup> The PMR during 2001–2010, although significantly elevated, was somewhat lower than in the preceding decade (1.64 vs 2.42), which may reflect reduced levels of exposure. Nevertheless, the hazard was responsible for approximately 11 excess deaths, and for every attributable fatality there are likely to have been many more non-fatal cases, including some associated with serious morbidity. An important proportion of such disease should be preventable by pneumococcal vaccination.

There is an urgent need for regulatory bodies internationally to consider the need for pneumococcal vaccination for welders and for the HSE to review its approach to the hazard. Similarly, clinicians advising welders should review their practice and the advice they give. Controls on exposure are important, but as currently applied in the UK, they



**To cite:** Coggon D, Harris EC, Cox V, et al. *Thorax* 2015;**70**:198–199.



**Table 1** Mortality of welders from pneumococcal and unspecified lobar pneumonia, England and Wales, 1991–2010

Cause of death	Deaths at age 16–64 years				Deaths at age 65–74 years			
	Observed	Expected	PMR	(95% CI)	Observed	Expected	PMR	(95% CI)
1991–2000								
Pneumococcal and unspecified lobar pneumonia	32	13.2	2.42	(1.66 to 3.42)	21	15.0	1.40	(0.86 to 2.13)
2001–2010								
Pneumococcal and unspecified lobar pneumonia	28	17.1	1.64	(1.09 to 2.36)	18	13.6	1.32	(0.78 to 2.09)
<i>Pneumococcal pneumonia</i>	7	1.96	3.58	(1.44 to 7.37)	0	0.82	0.00	(0.00 to 4.51)
<i>Unspecified lobar pneumonia</i>	21	15.2	1.38	(0.86 to 2.12)	18	12.8	1.40	(0.83 to 2.22)

PMR, proportional mortality ratio.

appear not to provide adequate protection. It is likely that the same applies in many other countries. An analogy can be drawn with hepatitis B immunisation for healthcare workers at risk of sharps injuries. Good working practices are necessary to minimise the occurrence of such injuries, but because in practice they are not fully effective, it is necessary to promote and provide additional protection by vaccination.

**Acknowledgements** We thank the Office for National Statistics, who provided us with the data on occupational mortality.

**Contributors** DC was the principal investigator, oversaw the analysis and wrote the first draft of the paper. ECH jointly carried out the analysis and contributed to the revision of the paper. VC also jointly carried out the analysis and contributed to the revision of the paper. KTP contributed to the design of the study and revision of the manuscript.

**Funding** Medical Research Council (MRC\_MC\_UP\_A620\_1018).

**Competing interests** None.

**Provenance and peer review** Not commissioned; internally peer reviewed.

**Data sharing statement** It is planned that data from the new national analysis of occupational mortality will be made publicly available when a full report of the study is published.

## REFERENCES

- Coggon D, Inskip H, Winter P, *et al.* Lobar pneumonia: an occupational disease in welders. *Lancet* 1994;344:41–4.
- Palmer K, Coggon D. Does occupational exposure to iron promote infection? *Occup Environ Med* 1997;54:529–34.
- Wergeland E, Iversen BG. Deaths from pneumonia after welding. *Scand J Work Environ Health* 2001;27:353.
- Palmer KT, Poole J, Ayres JG, *et al.* Exposure to metal fume and infectious pneumonia. *Am J Epidemiol* 2003;157:227–33.
- Palmer KT, Cullinan P, Rice S, *et al.* Mortality from infectious pneumonia in metal workers: a comparison with deaths from asthma in occupations exposed to respiratory sensitizers. *Thorax* 2009;64:983–6.
- Wong A, Marrie TJ, Garg S, *et al.* Welders are at increased risk for invasive pneumococcal disease. *Int J Infect Dis* 2010;14:e796–9.
- Torén K, Qvarfordt I, Bergdahl IA, *et al.* Increased mortality from infectious pneumonia after occupational exposure to inorganic dust, metal fumes and chemicals. *Thorax* 2011;66:992–6.
- Palmer KT, Cosgrove M. Vaccinating welders against pneumonia. *Occup Med* 2012;62:325–30.
- Palmer KT, Coggon D. Infectious pneumonia in workers exposed to metal fume. In: Venables KM, ed. *Current topics in occupational epidemiology*. Oxford University Press, 2013: 42–52, Chapter 4.
- Suri R, Palmer K, Ross J, *et al.* Exposure to welding fume and adhesion of *Streptococcus pneumoniae* to A549 alveolar cells. *Thorax* 2012; 67:A51.
- Department of Health. *Immunisation against infectious disease*. 3rd edn. 2006. (updated November 2011), Chapter 25 v2\_0,p305.
- Department of Health. *Immunisation against infectious disease*. 3rd edn. 2006. (updated October 2012), Chapter 25 v4\_0.
- Health and Safety Executive. *Pneumonia vaccination for employees exposed to welding and metal fume*. EIS44, December 2013. <http://www.hse.gov.uk/pubns/eis44.pdf> (accessed 1 Jul 2014).
- ICD CDC National Center for Health Statistics. *International Classification of Diseases, Ninth Revision*. <http://www.cdc.gov/nchs/icd/icd9cm.htm> (accessed 1 Jul 2014).

## **PRESS RELEASE**

### **THORAX**

#### **Health & Safety Executive (HSE) advice on pneumonia jobs for welders “flawed,” say experts**

*HSE disputed independent recommendations; new data show metal fumes still dangerous and no known safety threshold*

**[Pneumococcal vaccination for welders Online First doi 10.1136/thoraxjnl-2014-206129]**

The Health & Safety Executive’s advice to employers on pneumonia jobs for welders is flawed and needlessly risking lives, say occupational health experts in an opinion piece in the journal *Thorax*.

New data show that welders’ heightened risk of potentially lethal pneumonia, as a result of exposure to metal fumes, is still very real, they say.

Over the past 20 years, a growing body of evidence has linked exposure to metal fumes with a heightened risk of developing, and dying from, bacterial lobar pneumonia.

It is not clear exactly why this happens, but there are indications that the tiny particles in welding fumes boost the stickiness of pneumococcal bacteria to the cells lining the airways of the lungs.

The scientific evidence was sufficient to prompt the government’s independent advisory body, the Joint Committee on Vaccination and Immunisation (JCVI), to recommend in 2011 that employers offer welders a jab to protect them against the infection.

But, on the advice of the UK Health & Safety Executive, which aims to safeguard the health of the working population, the Department of Health in England decided that a blanket recommendation was not required.

Vaccination should be considered in the light of measures taken to curb the amount of exposure to metal fumes, it said.

Subsequently, the HSE published guidance for employers, urging them to prioritise controls on exposure, and emphasising that vaccination was not a regulatory requirement, provided that adequate steps had been taken to address fume levels.

But, write the authors: “A fundamental flaw of this approach is that, as the HSE’s guidance acknowledges, “there are no research studies which specify how much exposure to welding or metal fume (dose) will result in an increased susceptibility to pneumonia (response).”

As a result, it is impossible to tell what threshold of exposure is critical, and what limits are needed to curb the risk, irrespective of whether the HSE feels it knows what constitute effective controls on exposure, they say.

Using new data from the Office for National Statistics on deaths by occupation in England and Wales during 2001-2010, the authors checked whether welders still had a higher risk of

pneumonia than people in other jobs. They compared their calculations with a similar analysis for the period 1991-2000.

Between 2001 and 2010, 2863 men aged 16-74 died from pneumococcal and unspecified lobar pneumonia across all job categories compared with 3234 between 1991 and 2000.

But, just as in 1991-2000, deaths from pneumonia among welders in 2001-2010 were significantly more common than in other workers. In particular, welders of working age were more than 3.5 times as likely to die from pneumococcal pneumonia as men in other jobs.

Based on these figures, the authors estimate that there were 11 excess deaths nationally among welders from pneumonia in 2001-10. And while this is around half of what it was in 1991-2000 (19), it still means that “welders continued to experience significant excess mortality from pneumococcal and unspecified lobar pneumonia during 2001-2010,” they write.

“Deaths are only the tip of the iceberg,” points out lead author Professor David Coggon in an accompanying podcast. “For every excess death there will be many more cases of ill health, some of which will be quite serious,” he says, adding: “Pneumonia is not just a disease of old people, it can strike at all ages.”

The calculations prompt the authors to conclude that regulatory bodies around the globe should consider making the offer of vaccination for welders mandatory, and they suggest that the HSE should reconsider its position on the job.

“Controls on exposure are important, but as currently applied in the UK, they appear not to provide adequate protection,” they say, adding that the situation in the UK is unlikely to be unique.

**Contact:**

Professor David Coggon, or Professor Keith Palmer, MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton General Hospital, Southampton, UK

Tel: +44 (0)2380 777 624

Email: [dnc@mrc.soton.ac.uk](mailto:dnc@mrc.soton.ac.uk)

**Link to podcast:**

<https://soundcloud.com/bmjpodcasts/pneumococcal-vaccination-for-welders/s-T9KPR>