CORRESPONDENCE

Conventional bronchoscopic techniques in sarcoidosis: not too far behind

We read with interest the review by Rintoul *et al*¹ wherein the authors have summarised the usefulness of linear endobronchial ultrasound (EBUS) in a lucid manner. In the section on 'Investigation of Suspected Sarcoidosis', the authors state that the diagnostic yield to detect granulomas using endosonography (80%) is significantly higher than bronchoscopy (53%) using transbronchial lung biopsy (TBB) and endobronchial biopsy (EBB) as demonstrated by the GRANULOMA trial.² This gives an impression that conventional bronchoscopy techniques are significantly inferior to endosonography in the diagnosis of sarcoidosis. However, a recent randomised controlled trial presents evidence contrary to this notion.³ The trial compared the diagnostic yield of routine bronchoscopy techniques (conventional transbronchial needle aspiration (TBNA) plus EBB and TBB) with EBUS-TBNA (combined with EBB and TBB). The yield of EBUS-TBNA (74.5%) was significantly better than conventional TBNA (48.4%, $p=0.004$) in this study. However, when

conventional TBNA was combined with EBB and TBB, the yield (85.5%) was not significantly different as compared with EBUS-TBNA with EBB and TBB (92.7%, $p=0.34$).³ Thus, it is useful to have linear EBUS in one's armamentarium, but the bronchoscopist can achieve similar results with conventional bronchoscopic techniques in the diagnosis of sarcoidosis.⁴ This is in stark contrast to the situation in lung cancer staging where conventional TBNA (pooled sensitivity=39%) cannot match EBUS-TBNA (pooled sensitivity=88%) in terms of sensitivity.^{5,6}

Sahajal Dhooria, Ritesh Agarwal, Dheeraj Gupta

Department of Pulmonary Medicine, Postgraduate Institute of Medical Education and Research (PGIMER), Chandigarh, India

Correspondence to Dr Ritesh Agarwal, Department of Pulmonary Medicine, Postgraduate Institute of Medical Education and Research, Sector-12, Chandigarh 160012, India; agarwal.ritesh@live.com/riteshpgi@gmail.com

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

CrossMark

To cite Dhooria S, Agarwal R, Gupta D. *Thorax* 2015;**70**:587.

Received 15 January 2015

Accepted 2 March 2015

Published Online First 18 March 2015

- ▶ <http://dx.doi.org/10.1136/thoraxjnl-2014-205635>
- ▶ <http://dx.doi.org/10.1136/thoraxjnl-2015-206994>

Thorax 2015;**70**:587.

doi:10.1136/thoraxjnl-2015-206805

REFERENCES

- 1 Rintoul RC, Ahmed R, Dougherty B, *et al*. Linear endobronchial ultrasonography: a novelty turned necessity for mediastinal nodal assessment. *Thorax* 2014;**70**:175–80.
- 2 von Bartheld MB, Dekkers OM, Szlubowski A, *et al*. Endosonography vs conventional bronchoscopy for the diagnosis of sarcoidosis: the GRANULOMA randomized clinical trial. *JAMA* 2013;**309**:2457–64.
- 3 Gupta D, Dadhwal DS, Agarwal R, *et al*. Endobronchial ultrasound-guided transbronchial needle aspiration vs conventional transbronchial needle aspiration in the diagnosis of sarcoidosis. *Chest* 2014;**146**:547–56.
- 4 Agarwal R, Gupta D. Endobronchial ultrasound-guided transbronchial needle aspiration: Indian perspective. *J Bronchology Interv Pulmonol* 2014;**21**:284–7.
- 5 Holty JE, Kuschner WG, Gould MK. Accuracy of transbronchial needle aspiration for mediastinal staging of non-small cell lung cancer: a meta-analysis. *Thorax* 2005;**60**:949–55.
- 6 Adams K, Shah PL, Edmonds L, *et al*. Test performance of endobronchial ultrasound and transbronchial needle aspiration biopsy for mediastinal staging in patients with lung cancer: systematic review and meta-analysis. *Thorax* 2009;**64**:757–62.