CORRESPONDENCE

Response to: 'Conventional bronchoscopic techniques in sarcoidosis: not too far behind' by Dhooria *et al*

We thank Dhooria *et al*¹ for their complimentary comments on our review article about linear endobronchial ultrasound.² We congratulate them on their carefully performed randomised controlled trial comparing endobronchial ultrasound-guided transbronchial needle aspiration (EBUS-TBNA) in conjunction with transbronchial lung biopsies (TBLB) and endobronchial biopsies (EBB) with conventional non-ultrasound-guided transbronchial needle aspiration (cTBNA) in conjunction with TBLB and EBB for the diagnosis of sarcoidosis.³ In this study, they showed that EBUS-TBNA, as an individual test, has the highest diagnostic yield for granulomas in sarcoidosis (74.5%), a figure similar to that shown by other groups^{4,5} and considerably better than that achieved by cTBNA (48%).

Bronchoscopists now have a considerable armamentarium to choose from for investigating suspected sarcoidosis including EBUS-TBNA, cTBNA, endoscopic ultrasound, EBB, TBLB and bronchoalveolar lavage. Therefore, the question that arises is which is the best combination? To some extent, this will be defined by availability and expertise. Dhooria *et al* and others suggest that EBUS-TBNA in combination with TBLB optimises yield (90.9%), but it must be remembered that

headline figures achieved by expert practitioners may not be achievable by all. Furthermore, the additional yield provided by TBLB must be tempered by the risk of complication, specifically pneumothorax, which although admirably low in the study by Dhooria *et al* has been reported by others as occurring in 1–6% of cases.⁶ In situations where EBUS-TBNA is not available, cTBNA in combination with EBB and TBLB probably provides the highest accuracy but again is very dependent upon disease prevalence and expertise.⁷

Although it seems likely that EBUS-TBNA±TBLB will become the standard of care for investigating pulmonary sarcoidosis, further carefully designed comparative studies are warranted and these would benefit further from having a cost-effectiveness component.

Robert C Rintoul,¹ Rawya Ahmed,¹ Brendan Dougherty,¹ Nicholas R Carroll²

¹Department of Thoracic Oncology, Papworth Hospital, Cambridge, UK

²Department of Radiology, Cambridge University NHS Foundation Trust, Cambridge, UK

Correspondence to Dr Robert Rintoul, Department of Thoracic Oncology, Papworth Hospital, Cambridge CB23 3RE, UK; Robert.rintoul@nhs.net

Contributors RCR wrote the letter on behalf of the other authors who have reviewed it.

Competing interests None declared.

Provenance and peer review Not commissioned; internally peer reviewed.

CrossMark

To cite Rintoul Robert C, Ahmed R, Dougherty B, *et al*. *Thorax* 2015;**70**:588.

Received 1 March 2015

Accepted 3 March 2015

Published Online First 24 March 2015

► <http://dx.doi.org/10.1136/thoraxjnl-2015-206805>

► <http://dx.doi.org/10.1136/thoraxjnl-2014-205635>

Thorax 2015;**70**:588.

doi:10.1136/thoraxjnl-2015-206994

REFERENCES

- 1 Dhooria S, Agarwal R, Gupta D. Conventional bronchoscopic techniques in sarcoidosis: not too far behind. *Thorax* 2015;**70**:587.
- 2 Rintoul RC, Ahmed R, Dougherty B, *et al*. Linear endobronchial ultrasonography: a novelty turned necessity for mediastinal nodal assessment. *Thorax* 2015;**70**:175–80.
- 3 Gupta D, Dadhwal D, Agarwal R, *et al*. Endobronchial Ultrasound-guided transbronchial needle aspiration versus conventional transbronchial needle aspiration in the diagnosis of sarcoidosis. *Chest* 2014;**146**: 547–56.
- 4 Navani N, Booth HL, Kocjan G, *et al*. Combination of endobronchial ultrasound-guided transbronchial needle aspiration with standard bronchoscopic techniques for the diagnosis of stage I and II pulmonary sarcoidosis. *Respirology* 2011;**16**:467–72.
- 5 Tremblay A, Stather DR, Maceachern P, *et al*. A randomized controlled trial of standard vs endobronchial ultrasonography-guided transbronchial needle aspiration in patients with suspected sarcoidosis. *Chest* 2009;**136**:340–6.
- 6 Du Rand IA, Blaikley J, Booton R, *et al*. British Thoracic Society guideline for diagnostic flexible bronchoscopy in adults. *Thorax* 2013;**68**:i1–44.
- 7 Holty JE, Kushner WG, Gould MK. Accuracy of transbronchial needle aspiration for mediastinal staging of non-small cell lung cancer: a meta-analysis. *Thorax* 2005;**60**:949–55.