IMAGES IN THORAX

Teenager with chest pain and swollen neck: a leave-it-alone condition

Pen-Yuan Liao,^{1,2,3} Hung-Jung Wang³

¹Department of Emergency Medicine, Taipei Medical University Hospital, Taipei, Taiwan

²Imaging Research Center, Taipei Medical University, Taipei, Taiwan

³Department of Medical Imaging, Taipei Medical University Hospital, Taipei, Taiwan

Correspondence to

Dr Pen-Yuan Liao, Department of Emergency Medicine and Department of Medical Imaging, Taipei Medical University Hospital, 252 Wu-Shing Street, Taipei 110, Taiwan; pyliao100@hotmail.com

Received 25 February 2015
Accepted 28 February 2015
Published Online First
18 March 2015

ABSTRACT

A 19-year-old boy with shortness of breath and chest pain after strenuous exercise presented to emergency department. On physical examination, the neck and shoulders appeared to be swollen. There was crepitus on skin palpation. Chest X-ray disclosed diffuse subcutaneous emphysema and pneumomediastinum. CT showed additional finding of air in epidural space. The patient was discharged after 2 days of hospitalisation with conservative treatment uneventfully. Pneumorrhachis is usually caused by abrupt increase in intrathoracic pressure in instance of forceful vomiting, cough or asthma attack in an otherwise healthy young adult. It is usually accompanied with pneumomediastinum. The management of epidural pneumatosis should be tailored according to its primary cause. For most patients with pneumorrhachis associated to a spontaneous pneumomediastinum without neurological symptoms, this condition is generally self-limited. For epidural free air of large volume that causes neurological deficits, surgical laminectomy may be indicated.

A 19-year-old boy with shortness of breath and chest pain after strenuous exercise presented to emergency department (ED). He also had a common cold for 1 week prior to ED visit. On triage, he was afebrile, and haemodynamics were stable. On physical examination, the neck and shoulders appeared to be swollen, revealing crepitus on skin palpation. Chest X-ray disclosed diffuse subcutaneous emphysema and pneumomediastinum. CT showed additional finding of air in

epidural space (figure 1). The patient was discharged after 2 days of hospitalisation with conservative treatment uneventfully.

Pneumorrhachis is usually caused by abrupt increase in intrathoracic pressure in instance of forceful vomiting, cough or asthma attack in an otherwise healthy young adult. It is usually accompanied with pneumomediastinum, since the air from pulmonary alveoli has to travel across the mediastinum in order to reach the epidural space.¹ It is estimated that pneumorrhachis occurs in 9.5% of patients with spontaneous pneumomediastinum, which is already a rare disease entity and is responsible for 1 of 8005 to 1 of 42 000 hospital incidents and emergency admissions.^{1 2} Other possible origins of free air include retropharyngeal space, illicit drug use or direct trauma to the vertebral column. The presence of epidural free air usually does not cause symptoms; however, neurological deficits may occur if the volume of free air is large.³

The management of epidural pneumatosis should be tailored according to its primary cause. For most patients with pneumorrhachis associated with a spontaneous pneumomediastinum without neurological symptoms, this condition is generally self-limited if cough, emesis or asthma is under control. For pneumorrhachis of other aetiologies, such as oesophageal rupture, emphysematous infection and open trauma of the spine, more aggressive and specific treatment should be undertaken. For epidural free air of large volume that causes neurological deficits, surgical laminectomy may be indicated.

Figure 1 CT of the chest, showing extensive subcutaneous emphysema, pneumomediastinum and epidural pneumatosis. Free air dissected along the intervertebral foramina of upper thoracic vertebrae (arrow).

CrossMark

To cite: Liao P-Y, Wang H-J. *Thorax* 2015;**70**:707–708.

Contributors P-YL is the main writer and primary care physician of emergency department, and contributed to patient history. H-JW was the reporting radiologist and contributed to several parts of discussion.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; internally peer reviewed.

REFERENCES

- 1 Murayama S, Gibo S. Spontaneous pneumomediastinum and Macklin effect: overview and appearance on computed tomography. *World J Radiol* 2014;6:850–4.
- 2 Kono T, Kuwashima S, Fujioka M, *et al.* Epidural air associated with spontaneous pneumomediastinum in children: uncommon complication? *Pediatr Int* 2007;49:923–7.
- 3 Song KJ, Lee KB. Spontaneous extradural pneumorrhachis causing cervical myelopathy. *Spine J* 2009;9:e16–18.