Non-coding RNAs in the pathogenesis of COPD

Elise G De Smet,¹ Pieter Mestdagh,² Jo Vandesompele,² Guy G Brusselle,¹ Ken R Bracke¹

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/thoraxjnl-2014-206560>).

¹Laboratory for Translational Research in Obstructive Pulmonary Diseases, Department of Respiratory Medicine, Ghent University Hospital, Ghent, Belgium
²Center for Medical Genetics, Ghent University, Ghent, Belgium

Correspondence to

Dr Ken R Bracke, Department of Respiratory Medicine, Ghent University Hospital, De Pintelaan 185, 9000 Ghent, Belgium; ken.bracke@ugent.be

Received 19 November 2014

Revised 16 April 2015

Accepted 29 April 2015

Published Online First
20 May 2015

ABSTRACT

A large part of the human genome is transcribed in non-coding RNAs, transcripts that do not code for protein, including microRNAs (miRNAs) and long non-coding RNAs (lncRNAs). MiRNAs are short single-stranded RNA molecules that negatively regulate gene expression at the post-transcriptional level. They play an important regulatory role in many biological processes. Consequently, altered expression of these non-coding RNAs has been shown to lead to inflammation and disease. In contrast, lncRNAs, can both enhance or repress the expression of protein-coding genes. COPD is typically caused by tobacco smoking and leads to a progressive decline in lung function and a premature death. Exaggerated pulmonary inflammation is a hallmark feature in this disease, leading to obstructive bronchiolitis and emphysema. In this review, we discuss the miRNA expression patterns in lungs of patients with COPD and in mouse models and we highlight various miRNAs involved in COPD pathogenesis. In addition, we briefly discuss a specific lncRNA that is upregulated upon cigarette smoke exposure, providing a short introduction to this more recently discovered group of non-coding RNAs.

SEARCH STRATEGY

We searched PubMed with the search terms 'COPD' or 'emphysema' or 'smoke' in combination with one of the following terms: 'microRNA', 'miRNA', 'micro-RNA', 'long non-coding RNA', 'lncRNA' or 'lincRNA'. We also specifically searched for the combination: 'COPD/emphysema', 'miRNA/lncRNA' and 'biomarker'. We focused on lung and not on systemic disease manifestations of COPD and on *ex vivo/in vivo* studies of tissues or cells of patients with COPD and animal models instead of *in vitro* studies. The date of the last search was 10 April 2015. In addition, we consulted the reference lists of articles resulting from this search strategy and included those we judged relevant.

INTRODUCTION

COPD

COPD causes significant morbidity and mortality worldwide and is expected to become the third leading cause of death by 2020.¹ COPD is characterised by a progressive and irreversible airflow limitation, associated with an abnormal inflammatory response of the lungs to noxious particles and gasses.¹ Next to cigarette smoking, which is the main risk factor for COPD, exposure to smoke from biomass fuels and occupational exposure to chemicals or dust are also regarded as substantial threats.^{2,3}

Exaggerated inflammation in the airways and lung parenchyma is critical in the development and progression of COPD, causing small airway obstruction (obstructive bronchiolitis) and destruction of lung parenchyma (emphysema).¹ By triggering pattern recognition receptors (PRRs), noxious particles or gasses activate innate immune cells such as alveolar macrophages and airway epithelial cells. These cells subsequently release pro-inflammatory mediators such as chemokines and cytokines, mediating the recruitment of neutrophils and inflammatory monocytes from the blood circulation to the lung. The specialised antigen-presenting dendritic cells initiate adaptive immunity by promoting T helper 1 (T_H1) and T helper 17 (T_H17) cell differentiation and CD8⁺ cytotoxicity.

Obstructive bronchiolitis is caused by the accumulation of these immune cells in the airway lumen, airway wall and lung parenchyma and by structural alterations of the small airways (airway wall remodelling). Next to obstructive bronchiolitis, emphysema is the second pathological hallmark contributing to COPD. It compromises lung function and reduces the surface area available for gas exchange. Oxygen radicals and proteolytic enzymes are responsible for this lung destruction. Activated neutrophils and macrophages release such proteolytic enzymes, for example neutrophil elastase and matrix metalloproteinases (MMPs).⁴

The normal decline of lung function with age is accelerated in patients with COPD and slowly leads to disability and death.⁵ Furthermore, current pharmacological therapy is limited and is directed against the clinical manifestations, mainly leaving the underlying disease-causing process unaffected. The fact that only 20% of smokers develop COPD⁶ suggests that genetic and epigenetic factors contribute to the susceptibility for this disease.

Non-coding RNAs

Protein-coding genes represent only a small portion of the human genome (20 687 genes, <2%),⁷ whereas a large part is transcribed into non-coding RNAs (ncRNAs), including microRNAs (miRNAs) and long non-coding RNAs (lncRNAs). The purpose of this article is to establish a comprehensive overview of the literature on non-coding RNAs in COPD, focusing on miRNAs and lncRNAs. Literature on other non-coding RNAs in COPD (such as small nuclear RNAs, small nucleolar RNAs and piwi-interacting RNAs) is limited and beyond the scope of this review. For definition of terms see Table 1.

MiRNAs are the most widely studied class of ncRNAs. They are small RNA molecules (approximately 21–25 nucleotides long) that negatively regulate gene expression post-transcriptionally, by

CrossMark

To cite: De Smet EG, Mestdagh P, Vandesompele J, et al. *Thorax* 2015;**70**:782–791.

means of mRNA degradation, inhibition of protein translation, or by a combination of both mechanisms.⁸ MiRNAs are predicted to regulate approximately 60% of all human protein-coding genes.⁹ Dysregulation of miRNAs is reported to be involved in the pathogenesis of many diseases, including pulmonary diseases such as COPD.¹⁰

Many forms of small ncRNAs are relatively strongly conserved across species.¹¹ lncRNAs, in contrast, do not show such conservation and were originally classified as non-functional.¹² They comprise a heterogeneous class of RNAs, varying in nucleotide length (200 nucleotides to over 100 kilobases), cellular location (nucleus, cytoplasm or both), splicing pattern and functional and structural role in various biological processes.

The tertiary structure of lncRNAs allows them to bind to and interact with proteins, explaining their involvement in many regulatory functions. They can prevent or facilitate the recruitment of RNA polymerase II, transcription factors and/or cofactors to gene promoters, thereby controlling transcription of target genes. They also function as molecular scaffolds, bringing together multiple protein partners. Hundreds of lncRNAs have been found to form ribonucleic-protein interactions with multiple protein partners, supporting the concept of a molecular scaffold.¹³ They can also guide specific regulatory protein complexes to the correct chromosomal locations in *cis* or in *trans* providing target specificity to these complexes.

Besides their predominantly transcriptional modes of actions described so far, lncRNAs can also base-pair with mRNA molecules and affect their stability or translation.¹⁴ Finally, lncRNAs can compete for miRNA binding, and in that way prevent the miRNA from carrying out its function.¹⁵ This is their so-called sponge function as competing endogenous RNA.

MIRNA EXPRESSION UPON CIGARETTE SMOKE EXPOSURE AND IN COPD

As mentioned above, cigarette smoke (CS) is the uttermost important risk factor for the development of COPD, making it a relevant risk factor when studying COPD pathogenesis. It remains unknown by which mechanism CS is able to dysregulate miRNAs. Most likely, the high toxicity and mutagenicity of CS plays an important role, since CS contains radicals that can interact with DNA *in vitro*.¹⁶ Next to directly affecting the miRNA, disrupting miRNA regulatory mechanisms might be an alternative mechanism by which CS interferes with miRNA expression. An overview of the biological functions of miRNAs in the pathogenesis of COPD is illustrated in figure 1.

MiRNAs as potent regulators of gene and protein expression

Given the relative redundancy of complementary sequences between miRNAs and their target mRNAs, a single miRNA is capable of targeting several genes simultaneously. Moreover, by targeting transcription factors or genes involved in epigenetic regulation, miRNAs can indirectly alter expression of hundreds of genes at once. In addition, miRNAs have a tissue- and cell type-specific expression pattern, making them excellent fine-tuners of gene regulation. Consequently, dysregulation of miRNAs can have widespread effects on gene and protein expression, affecting a multitude of biological processes.

For example, miR-218 was downregulated in smokers relative to never-smokers in human bronchial epithelium,¹⁷ in induced sputum of current smokers with and without COPD in comparison to never-smokers¹⁰ and in the bronchoalveolar lavage (BAL) fluid of patients with COPD compared with control subjects.¹⁸ Amongst its targets, the transcription factor MAFG (v-maf

musculoaponeurotic fibrosarcoma oncogene homologue G, avian) represents a gene of particular interest, since its binding sites were significantly overrepresented among genes altered in smokers.¹⁷ In this way, miR-218 is capable of indirectly regulating a multitude of genes in response to CS by modulating the expression of the transcription factor MAFG.

Although proteins typically have half-lives in the range of a couple of hours to a few weeks,¹⁹ changes in gene expression were found to persist in the bronchial epithelium of ex-smokers years after smoking cessation, indicating the presence of persisting alterations in gene regulation upon CS exposure.²⁰ Likewise, genes differentially expressed between current and never-smokers were categorised based on their expression in former smokers. They were classified as rapidly reversible, slowly reversible and irreversible. Although the mRNA expression of most genes (79%) in this study was rapidly reversible upon smoking cessation, those genes classified as slowly reversible or irreversible still presented with a 1.5-fold change between former and never-smokers for at least 78 months after smoking cessation.²¹ In line with this, CS-induced changes in miRNA expression in murine lung are irreversible if a certain threshold is reached, depending on the duration and dose of smoke exposure.²² Taken together, these findings support the idea that persistent dysregulated gene expression, found in the lungs of smokers and ex-smokers, might be attributable to irreversible changes in miRNA expression induced by exposure to a sufficiently high dose of smoke, for a sufficiently long period of time.

MiRNA expression profiling upon CS exposure and in COPD

In general, most studies report a global downregulation of miRNA abundance in response to cigarette smoking.

Graff *et al*²³ found that cigarette smoking decreases global miRNA expression in human alveolar macrophages, possibly by modification of DICER, an RNA endonuclease involved in the maturation of miRNAs.²⁴ Interestingly, the decrease in abundance was more pronounced in individuals with a heavy smoking history in comparison to light smokers, suggesting a correlation between the number of downregulated miRNAs and smoking history.²³ Likewise, Izzotti *et al*²⁵ also reported a downregulation of miRNA expression in the lungs of smoke-exposed rats. Notably, further research by the same group demonstrated that both age and gender influenced smoke-related miRNA dysregulation in murine lungs.²⁶ In line with the results of smoke-exposed rat lungs²⁵ and of human alveolar macrophages,²³ the majority of differentially expressed miRNAs (23 out of 28) in human bronchial airway epithelium were downregulated in smokers compared with never-smokers.¹⁷

A global decrease in miRNA expression upon smoking can be explained by different possible mechanisms, such as a deficiency in miRNA processing (eg, due to Dicer inhibition), changes in primary miRNA transcription, an increase in miRNA degradation or localisation of miRNAs to stress granules (aggregates of mRNAs in conjunction with proteins that appear when the cell is under stress) and processing bodies (cytoplasmic foci containing aggregates of specific proteins and mRNAs), or relative upregulation of other RNA fractions, such as ribosomal RNAs or mRNAs. In the process of COPD pathogenesis, downregulation of miRNAs can provide a plausible mechanism for the observed upregulation of inflammatory mediators.

In contrast, Ezzie *et al*²⁷ analysed lung tissue of smokers with or without COPD and found the expression of 57 miRNAs to be increased in patients with COPD, while only 13 miRNAs showed a decreased expression.

Figure 1 The biological functions of microRNAs and long non-coding RNAs in the pathogenesis of COPD. Illustration of the hallmarks of COPD, with chronic inflammation, airway remodelling and emphysema as pathological hallmarks and oxidative stress and imbalance in fluid homeostasis as pathogenetic mechanisms. CFTR, cystic fibrosis transmembrane conductance regulator; Cox-2, cyclooxygenase-2; IL-1R1, interleukin 1 receptor 1; let-7c, lethal-7c; miR, microRNA; MMP12, matrix metalloproteinase 12; MPK-1, mitogen-activated protein kinase phosphatase 1; SCAL1, smoke and cancer-associated lncRNA-1; SMAD7, mothers against decapentaplegic homologue 7; TNFR-II, tumour necrosis factor receptor II.

The discrepancy between this study and previous studies reporting a global downregulation of miRNAs^{10 17 23} might be due to the fact that miRNAs have a cell type-specific expression pattern and the global analysis of miRNA expression in whole lung tissue, where multiple cell types are present, might not reflect the situation in those cell types important for disease pathogenesis. Alternatively, differences in gene expression analysis workflow in general or normalisation procedure more specifically might also account for the observed discrepancies. Microarrays, reverse transcription quantitative PCR (RT-qPCR) and small RNA sequencing are widely used miRNA expression evaluation platforms. Nonetheless, these methods do not always deliver the same results.^{28 29} Microarray analysis, for example, can result in false positives (non-specific hybridisation) or false negatives (when hybridisation signals do not exceed the background level).

In addition, patient characteristics (such as COPD stage, age, gender, number of pack-years smoked, duration of smoking cessation) and sample size of the study population or differences in study design may also have a substantial influence on the results (for an overview of these parameters, see online supplemental table S1). Given that miRNA expression values are highly context dependent, an adequate experimental design and a sound statistical analysis are of high importance.

MiRNAs modulate the chronic pulmonary inflammation in COPD

Exaggerated and imbalanced inflammatory responses play an important role in COPD pathogenesis. These inflammatory responses are mediated by inflammatory proteins that are released upon CS exposure by structural cells of the lungs and

by inflammatory cells. Pro-inflammatory transcription factors, such as nuclear factor κ B (NF- κ B), regulate the expression of these inflammatory proteins. Amplification of this inflammatory response upon CS exposure might lead to the development of COPD. For an overview of the expression pattern of the discussed miRNAs in patients with COPD, smokers and never-smokers see table 2. In addition, table 2 contains the accession number and the most recent annotation of each miRNA, as checked by miRBase Tracker.³⁰ For an overview of miRNA profiling studies in rodent models discussed in this review see online supplemental table S2.

MiR-146a

Increased expression of cyclooxygenase-2 (COX-2) protein is a hallmark of chronic inflammation in CS-related diseases such as COPD and lung cancer.^{31 32} Sato *et al* showed that while interleukin (IL)-1 β and tumour necrosis factor (TNF)- α , two potent inducers of the NF- κ B pathway, induced the expression of miR-146a in both COPD and control fibroblasts, this was less pronounced in COPD fibroblasts. They suggested that this diminished miR-146a expression in COPD fibroblasts resulted in a reduced degradation and a prolonged half-life of COX-2 mRNA, a predicted target of miR-146a (TargetScan, miRBase), ultimately leading to overexpression of COX-2 protein.³³ COX-2 is a key enzyme in prostaglandin E₂ (PGE₂) biosynthesis, a promoter of neutrophil recruitment.³⁴ PGE₂ also inhibits the repair functions of lung fibroblasts³⁵ and is increased in the lungs of patients with COPD and in COPD fibroblasts in culture.^{33 36}

Likewise, miR-146a was downregulated in sputum of patients with COPD and smokers compared with never-smokers and in

Table 1 Definition of terms

Conditional knockout	An organism where gene disruption occurs in a tissue- or time-specific manner
Dicer	An enzyme involved in the processing of precursor microRNAs (pre-miRNAs) into mature miRNAs
Epigenetics	All changes in gene function that can be transmitted from one generation to the next, but that are not coded in the DNA sequence itself. Examples are DNA methylation or histone acetylation
Gene promoter	A region of DNA where transcription of genes is initiated
Gene set enrichment analysis	A computational method used to determine whether an a priori defined set of genes shows statistically significant differences between two biological states
Hypermethylation	High degree of DNA methylation, resulting in a decreased gene expression
Introns	Nucleotide sequences within a transcribed gene that are removed (spliced out) when the mature RNA product of a gene is formed
Long non-coding RNA (lncRNA)	Non-coding RNA with a length greater than 200 nucleotides
Messenger RNA (mRNA)	A protein-coding RNA molecule
MicroRNA (miRNA)	Small RNA molecule (approximately 21–25 nucleotides long) that negatively regulates gene expression post-transcriptionally
miRNA mimic	A synthetic RNA molecule designed to resemble an endogenous miRNA, typically used to study its function
miRNA processing	The conversion of an immature miRNA into a mature miRNA by RNA endonucleases
Post-transcriptional	An event that occurs after the transcription of DNA to RNA
Post-translational modification	A step in protein biosynthesis where polypeptide chains undergo modifications in order to form the mature protein product
Primary miRNA	RNA molecule that is cleaved in the nucleus to form a shorter double stranded hairpin (pre-miRNA), which, in his turn, is cleaved to generate the mature miRNA
Ribosomal RNA	The predominant part of the ribosome, essential for protein synthesis
RNA endonuclease	Enzyme that is able to cleave a RNA chain
RNA polymerase	Enzyme that catalyses the transcription from DNA to RNA
RNA splicing	Process where introns are removed and exons are joined either shortly after or concurrent with transcription, in order to form a mature RNA
Small interfering (si)RNA knockdown	The use of a short complementary RNA molecule in order to reduce the expression of a specific gene
Transcription	The process where a particular DNA segment is transcribed into RNA by the enzyme RNA polymerase
Transcription factor	Protein that can activate or repress the transcription of a gene, by binding to a regulatory DNA sequence (eg, in the gene promoter)
Translation	The process in which mRNA is converted to proteins

smokers with COPD in comparison to smokers without COPD.¹⁰ In contrast, miR-146a expression was increased in lung tissue from patients with COPD in comparison to smokers.²⁷

MiR-135b

MiR-135b seems to be involved in a negative feedback mechanism attempting to curtail excessive inflammatory responses (figure 2). MiR-135b was found to be significantly upregulated in the BAL fluid of CS-exposed mice and upon *in vitro* activation of the IL-1 receptor 1 (IL-1R1) by its agonist IL-1 α .³⁷ Further experiments indicated that CS induces an increase in pulmonary expression of miR-135b and that this increase is dependent on IL-1 α /IL-1R1 signalling.³⁷

Furthermore, the authors demonstrated that IL-1R1 and caspase-1 are likely targeted by miR-135.³⁷ Activated caspase-1 generates the mature form of IL-1 β by cleaving its precursor, pro-IL-1 β .³⁸ Both mature IL-1 β and IL-1 α activate the IL-1R1. In the post-acute phase, miR-135b might thus function as a negative feedback regulator by binding on its own regulator (IL-1R1) and its downstream effector, caspase-1, in an attempt to halt the progression of inflammation into the chronic phase (figure 2). In accordance with this hypothesis, IL-1 β has, in general, been described to be upregulated and implicated in early responses upon CS exposure,³⁹ but not upon chronic CS exposure.⁴⁰

An appropriate immune response is required to protect an organism against harmful attacks from various pathogens and other damaging events such as CS. However, when the activated immune response is too excessive, this can lead to pathological manifestations causing inflammatory disorders such as COPD. In this respect, negative feedback regulation is an important

mechanism keeping inflammation in check. As certain miRNAs might be important negative feedback regulators, the feedback control may be lost due to reduced expression and concomitant exacerbated inflammation may occur.

Let-7c

Analysis of miRNA expression in induced sputum supernatant in the study of Van Pottelberge *et al*¹⁰ revealed that the expression of let-7c (lethal-7c) was significantly lower in patients with COPD in comparison with never-smokers and smokers without COPD. Moreover, let-7c was significantly associated with FEV₁ (% predicted) and was previously reported to be downregulated in rat and mouse lungs upon CS exposure.^{22–25} The protein level of TNF receptor II (TNFR-II) in sputum, a predicted target gene of let-7c (MiRgen database), was inversely correlated with the level of let-7c. Gene set enrichment analysis revealed that target genes of let-7c are significantly enriched in patients with severe and very severe COPD in comparison to patients with moderate COPD. Furthermore, TNFR-II was enriched in the sputum of patients with severe and very severe COPD.¹⁰ TNFR-II plays an important role in COPD pathogenesis, evidenced by the fact that TNFR-II knockout mice are protected against CS-induced inflammation and emphysema.⁴¹ Let-7 mimics have been investigated as cancer therapeutics in mice⁴² and, given the demonstrated importance of let-7c in COPD pathogenesis, it might also be an interesting therapeutic target to investigate in preclinical COPD models.

MiR-101 and miR-144

The cystic fibrosis transmembrane conductance regulator (CFTR) protein, a chloride channel involved in the maintenance of fluid homeostasis in the lung, is suppressed in the lungs of

Table 2 Expression pattern of miRNAs in patients with COPD, smokers and never-smokers

miRNA	Most recent annotation*	Accession number	Disease stage	Smoker vs never-smoker	COPD vs smoker	COPD vs never-smoker	Increasing emphysema severity	Tissue/cell-type	Targets	References
COPD										
miR-218	hsa-miR-218-5p	MIMAT0000275	no COPD	↓				Bronchial airway epithelium	MAFG (validated)	17
			GOLD I and II	↓		↓		Induced sputum		10
			GOLD I to IV		↓			BAL fluid		18
miR-146a	hsa-miR-146a-5p	MIMAT0000449	GOLD I to IV		↓			Fibroblasts	COX-2 (predicted)	33
			GOLD I and II	↓	↓	↓		Induced sputum		10
			GOLD I, II and IV		↑			Lung tissue		27
let-7c	hsa-let-7c-5p	MIMAT0000064	GOLD I and II	↓	↓	↓		Induced sputum	TNFR-II (predicted)	10
miR-101	hsa-miR-101-3p	MIMAT0000099	GOLD IV			↑↑		Lung tissue	CFTR (validated), MPK-1 (predicted)	44
miR-15b	hsa-miR-15b-5p	MIMAT0000417	GOLD I, II and IV		↑			Lung tissue	SMAD7 (validated)	27
miR-144	hsa-miR-144-3p	MIMAT0000436	GOLD I, II and IV		↑			Lung tissue	CFTR (validated)	27
miR-199a-5p	hsa-miR-199a-5p	MIMAT0000231	not specified			↓↑		Peripheral blood monocytes	ATF6, NF-κB1, RELA, GRP78, IRE1, HIF-1α (validated)	49
			GOLD I to IV			↑		Lung tissue		57
			GOLD I to IV		↓			T-regulatory cells		50
miR-34b	hsa-miR-34b-3p	MIMAT0004676	GOLD I and II	↓		↓		Induced sputum		10
miR-452	hsa-miR-452-5p	MIMAT0001635	No COPD	↓				Alveolar macrophages	MMP12 (predicted)	23
miR-34c	hsa-miR-34c-5p	MIMAT0000686	GOLD I and II	↓		↓		Induced sputum		10
Emphysema										
miR-34c	hsa-miR-34c-5p	MIMAT0000686	GOLD I and II: Mild versus moderate emphysema (measured by KCO)				↓	Lung tissue		63
miR-638	hsa-miR-638	MIMAT0003308	GOLD IV (measured by Lm on CT)				↑	Lung tissue		52

*MiRBase release 21.

†Patients with COPD were compared with controls (the article does not specify whether the controls were smokers or non-smokers).

ATF6, activation of transcription factor 6; BAL, bronchoalveolar lavage; CFTR, cystic fibrosis transmembrane conductance regulator; COX-2, cyclooxygenase-2; GOLD, Global Initiative for Chronic Obstructive Pulmonary Disease; GRP78, glucose response protein 78; HIF-1α, hypoxia inducible factor 1α; IRE1, inositol requiring enzyme 1α; KCO, lung carbon monoxide transfer coefficient; Lm, mean linear intercept; MAFG, v-maf musculoaponeurotic fibrosarcoma oncogene homologue G, avian; miR, microRNA; MMP-12, matrix metalloproteinase 12; MPK-1, mitogen-activated protein kinase phosphatase 1; NF-κB1, nuclear factor κB1; RELA, V-Rel avian reticuloendotheliosis viral oncogene homologue A; SMAD7, mothers against decapentaplegic homologue 7; TNFR-II, tumour necrosis factor receptor II.

Figure 2 Proposed negative feedback mechanism involving miR-135b. Cigarette smoke induces an increase in pulmonary expression of miR-135b in mice and this increase is dependent on IL-1 α /IL-1R1 signalling.³⁷ Activated caspase-1 generates the mature form of IL-1 β by cleaving its precursor, pro-IL-1 β . Both mature IL-1 β and IL-1 α activate the IL-1R1, which transduces its signal through myeloid differentiation primary response 88 protein (MyD88), ultimately resulting in the activation of the transcription factors NF- κ B and activator protein 1 (AP-1, not shown here). NF- κ B activation induces pro-IL-1 β transcription and might also regulate the transcription of miR-135b (dashed arrow, observed in lung cancer cells⁸²). IL-1R1 and caspase-1 are likely targeted by miR-135b.³⁷ In the post-acute phase, miR-135b might thus function as a negative feedback regulator by binding on its own regulator (IL-1R1) and its downstream effector, caspase-1, in an attempt to halt the progression of inflammation into the chronic phase. Figure adapted from Halappanavar *et al*.³⁷ IL-1 α , interleukin 1 α ; IL-1RI, interleukin 1 receptor 1; IL-1RAcP, IL-1R accessory protein; miRNA, microRNA; mRNA, messenger RNA; MyD88, myeloid differentiation primary response protein 88; NF- κ B, nuclear factor κ B.

patients with COPD with a history of smoking.⁴³ Hassan *et al*⁴⁴ showed that expression of miR-144 and miR-101 suppresses CFTR protein in human bronchial epithelial cells and that they target the 3' untranslated region of CFTR *in vitro*. Both miRNAs are upregulated by CS and cadmium.⁴⁴

In addition, miR-144 was found to be upregulated in lung tissue from patients with COPD in comparison to smokers without COPD.²⁷ Chronic cigarette smoking might thus upregulate miR-101 and miR-144, and this may result in the suppression of CFTR in the lungs of patients with COPD, causing an imbalance in fluid homeostasis (figure 3).

MiR-15b

MiR-15b is significantly upregulated in COPD tissue and is expressed in bronchial epithelium and alveolar wall in type II pneumocytes.²⁷ SMAD7 (mothers against decapentaplegic homologue 7) is involved in the pathogenesis of COPD⁴⁵ and is a predicted target of miR-15b (TargetScan V5.1, PicTar, miRanda) (figure 3). SMAD7 has an inhibitory function in transforming growth factor β (TGF- β) signalling and was markedly downregulated in bronchial biopsies of patients with COPD.²⁷ Moreover, knockdown of miR-15b in Beas2B bronchial epithelial cells resulted in increased SMAD7 protein expression and, consequently, a reduced early phosphorylation of SMAD3 in response to TGF- β . In line with this, overexpression of miR-15b caused an increased early phosphorylation of SMAD3.²⁷

MiRNA-199a-5p

CS causes proteins to misfold, leading to ER stress and the activation of the unfolded protein response.⁴⁶ GRP78 (glucose response protein 78), IRE1 (inositol requiring enzyme 1 α), ATF6 (activation of transcription factor 6), NF- κ B1 and RELA (V-Rel avian reticuloendotheliosis viral oncogene homologue A) are all involved in the unfolded protein response and are confirmed targets of miR-199a-5p.^{47–49}

By studying isolated peripheral blood monocytes from individuals with or without COPD, a mechanism was put forward whereby hypermethylation of the miR-199a-5p promoter induces miR-199a-5p silencing, followed by derepression of proteins involved in the unfolded protein response and consequent unfolded protein response activation in patients with COPD.⁴⁹

Recently, Chatila *et al*⁵⁰ demonstrated that miR-199a-5p is significantly higher expressed in T-regulatory (T_{reg}) cells compared with T-effector (T_{eff}) cells in peripheral blood. Moreover miR-199a-5p was approximately fourfold lower in T_{reg} cells of patients with COPD compared with healthy smokers.⁵⁰ The role of T_{reg} cells in the pathogenesis in COPD is still undetermined. These cells are CD4⁺ T cells with an immunoregulatory function, mediating a suppressive effect on other T cells and antigen-presenting dendritic cells.

The role of miRNAs in emphysema

COPD is a heterogeneous disease, with patients suffering from varying degrees of obstructive bronchiolitis and emphysema.

Figure 3 Schematic overview of hypothetical functions of miR-144, miR-101 and miR-15b in bronchial epithelial cells. The CFTR protein is suppressed in the lungs of patients with COPD with a history of smoking.⁴³ MiR-144 and miR-101 are upregulated by cigarette smoke and in patients with COPD.^{27, 44} Expression of miR-144 and miR-101 suppresses CFTR protein in HBE cells and both miRNAs target the 3' untranslated region of CFTR in vitro.⁴⁴ Chronic cigarette smoking might upregulate miR-101 and miR-144, and this may result in the suppression of CFTR in the lungs of patients with COPD, causing an imbalance in fluid homeostasis. MiR-101 targets MPK-1.⁵³ High expression of this miRNA could lead to diminished dephosphorylation by MPK-1 and a sustained activation of ERK 1/2, seen in patients with COPD.⁵⁴ Notably, ERK1/2 signalling is also required for the increased expression of MMP-1 in emphysematous lung tissue of chronic smokers.⁵⁵ SMAD7 is a predicted target of miR-15b. It has an inhibitory function in transforming growth factor β (TGF- β) signalling and was markedly downregulated in bronchial biopsies of patients with COPD.²⁷ Overexpression of miR-15b in Beas2B bronchial epithelial cells resulted in decreased SMAD7 protein expression and, consequently, an increased early phosphorylation of SMAD3 in response to TGF- β .²⁷ CFTR, cystic fibrosis transmembrane conductance regulator; Cl⁻, chloride; ERK 1/2, extracellular signal-regulated kinase 1/2; miR, microRNA; MMP1, matrix metalloproteinase 1; MPK1, mitogen-activated protein kinase phosphatase 1; SMAD, mothers against decapentaplegic; TGF- β , transforming growth factor β .

Importantly, there is only a weak correlation between these two disease manifestations, where the same FEV₁ value (as measured by spirometry) can be accompanied by a low or high degree of emphysema (as evidenced by pathological examination of lung tissue resection specimens, CT scan of the lungs (imaging), or measurement of the diffusing capacity of the lungs) in different patients.⁵¹ However, most studies on miRNA expression in COPD use a disease classification solely based on the degree of airflow obstruction. For an overview of the expression pattern of the discussed miRNAs in patients with COPD, smokers and never-smokers see [table 2](#).

MiR-638

Notably, miRNA expression seems to be correlated with regional emphysema severity.⁵² MiR-638, for example, was positively correlated with emphysema severity and had more than 50 positively or negatively regulated targets. Inhibition of this miRNA in lung fibroblasts resulted in overexpression of its predicted targets, including genes involved in the oxidative stress response and extracellular matrix remodelling.⁵²

MiR-101

Hassan and colleagues found miR-101 to be highly expressed in the lungs of patients with COPD and in the lungs of mice

subjected to CS.⁴⁴ MiR-101 targets mitogen-activated protein kinase (MAPK) phosphatase 1 (MPK-1), an inhibitor of MAPKs.⁵³ High expression of this miRNA could lead to reduced dephosphorylation of extracellular signal-regulated kinase (ERK) 1/2 by MPK-1 and a sustained activation of ERK 1/2, seen in patients with COPD⁵⁴ ([figure 3](#)). Notably, ERK1/2 signalling is also required for the increased expression of the protease MMP-1 in emphysematous lung tissue of chronic smokers.⁵⁵

MiR-452

MiR-452 is highly downregulated in alveolar macrophages from active smokers compared with non-smokers.²³ Upon transfection of a miR-452 inhibitor in human monocytic THP-1 cells that are induced to differentiate toward a macrophage phenotype, the expression of matrix metalloproteinase 12 (MMP12) mRNA (a predicted target, MicroCosm database) was increased. MMP12 is secreted by macrophages and plays an important role in the pathogenesis of COPD by degrading elastin, thereby causing emphysema.⁵⁶

MiR-199a-5p

In contrast to Hassan *et al*,⁴⁹ Mizuno and colleagues reported an increased miR-199a-5p expression in lung tissue samples of

patients with COPD in comparison to control samples of never-smokers with normal lung function.⁵⁷

The authors propose a mechanism whereby oxidative stress mediates an upregulation of TP53, which in turn upregulates miR-34a, followed by regulation of miR-199a-5p, possibly through AKT inactivation.⁵⁷

Transfection of a miR-199a-5p precursor in human pulmonary microvascular endothelial cells resulted in a decreased hypoxia inducible factor 1 α (HIF-1 α) expression,⁵⁷ in line with decreased expression in lungs from patients with COPD.⁵⁸

Furthermore, HIF-1 α positively regulates vascular endothelial growth factor (VEGF) gene expression,⁵⁹ which is of importance, since conditional knockout of the VEGF ligand⁶⁰ and inhibition of its receptor⁶¹ in mouse lungs resulted in emphysema. In addition, in a genome-wide gene expression profiling study of lung tissue samples from current and ex-smokers with and without COPD, VEGF was one of the most downregulated genes in COPD samples.⁶²

MiR-34c and miR-34b

In a recent study miRNAs differentially expressed between mild and moderate emphysematous lung were identified.⁶³ Five miRNAs were significantly downregulated in lungs from patients with moderate emphysema, with miR-34c exhibiting the largest fold change. Supporting these findings, a more than threefold downregulation of miR-34c was observed in sputum from smokers with or without COPD compared with never-smokers.¹⁰ Moreover, miR-34c was significantly downregulated in the lungs of rats exposed to CS.²⁵

MiR-34b originates from the same primary transcript as miR-34c and its expression is also reduced in the sputum from smokers with or without COPD compared with never-smokers¹⁰ and in the lungs of mice²² and rats²⁵ exposed to CS.

MiR-34c acts as a tumour suppressor and is downregulated in many types of cancer, including lung cancer.⁶⁴ A synthetic mimic of miR-34 (MRX34) is the first miRNA mimic that has reached a phase I clinical trial status,⁶⁵ aiming to restore its function in a range of cancers.

Further, given the downregulation of miR-34c in serum of patients with COPD compared with healthy controls, it was suggested that this miRNA might serve in the clinic as a biomarker in COPD.⁶⁶ These studies open a window to the potential for certain miRNAs to serve as diagnostic and therapeutic agents in the combat against COPD.

Use of non-coding RNAs as biomarkers in COPD

Since miRNAs are remarkably stable in biofluids⁶⁷ and can be easily and reproducibly detected, they make excellent candidates for risk prediction and minimally invasive diagnosis of COPD.

For example, levels of serum miR-21 and miR-181a and their ratio are suggested to have potential biomarker value for predicting the development of COPD in heavy asymptomatic smokers.⁶⁸

Also in serum, five miRNAs (miR-34c-5p, miR-20a, miR-28-3p, miR-100 and miR-7) were differentially expressed between patients with COPD and controls.⁶⁶ This suggests that these miRNAs might have a biological function in the pathogenesis of COPD and in the future might serve as biomarkers of COPD.

In sputum, an easily accessible body fluid, detection of both miRNAs and lncRNAs has also been reported.^{10 69} In addition, by using sampling of exhaled breath condensates, another minimally invasive method, it is also possible to detect differential expression of miRNAs.⁷⁰

DYSREGULATION OF LONG NON-CODING RNAs BY CS EXPOSURE AND IN PATIENTS WITH COPD

Unlike miRNAs, lncRNAs are much less characterised. The function of some individual lncRNAs (such as HOTAIR (LNCipedia ID: lnc-SMUG1-7^{71 72})) has been elucidated, but the function of most lncRNAs remains unknown.

Recently, a novel lncRNA was characterised, named the smoke and cancer-associated lncRNA-1 (SCAL1, lnc-ARRDC3-1), which is upregulated by CS extract *in vitro*⁷³ and is elevated in the airway epithelia of cigarette smokers versus non-smokers.^{74 75} As its name suggests, SCAL1 is also elevated in many lung cancer cell lines. Small interfering RNA knockdown experiments demonstrated that SCAL1 is transcriptionally regulated by nuclear factor erythroid 2 related factor (NRF2). NRF2 is involved in the regulation of protective antioxidant genes⁷⁶ and NRF2 loss of function sensitises lung cells to COPD-like changes.⁷⁷ Furthermore, NRF2 expression is reduced in pulmonary macrophages of patients with COPD.⁷⁸

Knockdown of SCAL1 in bronchial epithelial cells resulted in a significant potentiation of CS-induced cytotoxicity, indicating that it might function as a protector against oxidative stress (see figure 1). It is suggested that SCAL1 mediates some of the cytoprotective functions of NRF2 toward CS-induced stress.⁷³

The authors also analysed several publically available databases and found that CS is able to alter the expression of numerous lncRNAs in airway epithelial cells.⁷³ lncRNAs, similar to miRNAs, might function as regulators of large sets of genes that protect against environmental stresses such as CS exposure.

Interestingly, lncRNAs are expressed in a more pronounced tissue-specific manner than protein-coding genes,⁷⁹ making them attractive future diagnostic and therapeutic targets.

Recently, a study was published comparing the lncRNA expression profile in lung tissue from non-smokers and smokers with or without COPD.⁸⁰ Upon comparison of patients with COPD and smokers without COPD, upregulation of 120 and downregulation of 43 lncRNAs was reported. Given the small sample size and the inclusion of only male subjects, additional larger studies with both genders are warranted.

FUTURE DIRECTIONS

COPD is a heterogeneous disease, with patients suffering from varying degrees of obstructive bronchiolitis and emphysema. As it stands, most studies on non-coding RNA expression profiling in COPD characterise patients solely by measurements of airflow obstruction, not taking the degree of emphysema into account (see online supplemental table S1). Nonetheless, emphysema is independently associated with poor clinical outcome and might also be the predominant disease manifestation in certain patients with COPD.⁸¹ Given this limitation, future studies should make an accurate assessment of emphysema extent and severity to produce a more complete classification of patients with COPD, allowing a more precise comparison between different groups.

As discussed in this review, great difficulties arise when trying to find overlap between miRNA profiling studies. Next to taking differences in patient characteristics into account, it remains important that findings are replicated in independent validation cohorts, by different studies and by different laboratory measurement methods. Since RT-qPCR and microarray can only quantify known targets, (small) RNA sequencing forms an interesting method for discovery of novel tissue- and disease-specific ncRNAs.

Upon differential expression analysis of ncRNAs, the resulting (top) hits should be functionally validated. Exhaustive profiling experiments result in lists of statistically significant differentially expressed ncRNAs, but little is known about the biological significance of these findings. This can be achieved for example by *in vivo* experiments, using either transgenic/knockout animals or perturbation experiments.

CONCLUSION

Current therapy for COPD has only limited disease-modifying effects. Hence, there is a great need for novel therapeutic interventions and preventive strategies. Research concerning the pathogenetic mechanisms causing COPD would greatly contribute to an improved therapy for patients suffering from this disease. The above-mentioned studies highlight the involvement of ncRNAs in the host response towards environmental exposures. In doing so, they are capable of influencing the susceptibility towards lung diseases such as COPD, linking environmental exposures to disease development and progression. Identification of ncRNAs important in these processes could provide new clinical targets as well as diagnostic and prognostic tools for COPD.

Contributors EGDS performed the literature search and wrote the manuscript. PM, JV, GGB and KRB reviewed and edited the manuscript.

Funding Presented work within the Department of Respiratory Medicine of Ghent University is funded by grants from the Fund for Scientific Research in Flanders (FWO Vlaanderen, G.OA99.13), the Concerted Research Action of the Ghent University (BOF/GOA, 01G02714) and the Interuniversity Attraction Poles Program (IUAP, P7/30).

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- Global Initiative for Chronic Obstructive Pulmonary Disease. *Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease*.
- Ezzati M. Indoor air pollution and health in developing countries. *Lancet* 2005;366:104–6.
- Trupin L, Earnest G, San Pedro M, *et al*. The occupational burden of chronic obstructive pulmonary disease. *Eur Respir J* 2003;22:462–9.
- Demedts IK, Morel-Montero A, Lebecque S, *et al*. Elevated MMP-12 protein levels in induced sputum from patients with COPD. *Thorax* 2006;61:196–201.
- Fletcher C PR, Tinker C, Speizer FE. *The natural history of chronic bronchitis and emphysema*. Oxford: Oxford University Press, 1976.
- Pauwels RA, Rabe KF. Burden and clinical features of chronic obstructive pulmonary disease (COPD). *Lancet* 2004;364:613–20.
- Consortium EP, Bernstein BE, Birney E, *et al*. An integrated encyclopedia of DNA elements in the human genome. *Nature* 2012;489:57–74.
- Lim LP, Lau NC, Garrett-Engle P, *et al*. Microarray analysis shows that some microRNAs downregulate large numbers of target mRNAs. *Nature* 2005;433:769–73.
- Friedman RC, Farh KK, Burge CB, *et al*. Most mammalian mRNAs are conserved targets of microRNAs. *Genome Res* 2009;19:92–105.
- Van Pottelberge GR, Mestdagh P, Bracke KR, *et al*. MicroRNA expression in induced sputum of smokers and patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2011;183:898–906.
- Bentwich I, Aniel A, Karov Y, *et al*. Identification of hundreds of conserved and nonconserved human microRNAs. *Nat Genet* 2005;37:766–70.
- Brosius J. Waste not, want not—transcript excess in multicellular eukaryotes. *Trends Genet* 2005;21:287–8.
- Tsai MC, Manor O, Wan Y, *et al*. Long noncoding RNA as modular scaffold of histone modification complexes. *Science* 2010;329:689–93.
- Carrieri C, Cimatti L, Biagioli M, *et al*. Long non-coding antisense RNA controls Uchl1 translation through an embedded SINEB2 repeat. *Nature* 2012;491:454–7.
- Wang J, Liu X, Wu H, *et al*. CREB up-regulates long non-coding RNA, HULC expression through interaction with microRNA-372 in liver cancer. *Nucleic Acids Res* 2010;38:5366–83.
- Church DF, Pryor WA. Free-radical chemistry of cigarette smoke and its toxicological implications. *Environ Health Perspect* 1985;64:111–26.
- Schembri F, Sridhar S, Perdomo C, *et al*. MicroRNAs as modulators of smoking-induced gene expression changes in human airway epithelium. *Proc Natl Acad Sci USA* 2009;106:2319–24.
- Molina-Pinelo S, Pastor MD, Suarez R, *et al*. MicroRNA clusters: dysregulation in lung adenocarcinoma and COPD. *Eur Respir J* 2014;43:1740–9.
- Schwanhauser B, Busse D, Li N, *et al*. Global quantification of mammalian gene expression control. *Nature* 2011;473:337–42.
- Spira A, Beane J, Shah V, *et al*. Effects of cigarette smoke on the human airway epithelial cell transcriptome. *Proc Natl Acad Sci USA* 2004;101:10143–8.
- Beane J, Sebastiani P, Liu G, *et al*. Reversible and permanent effects of tobacco smoke exposure on airway epithelial gene expression. *Genome Biol* 2007;8:R201.
- Izzotti A, Larghero P, Longobardi M, *et al*. Dose-responsiveness and persistence of microRNA expression alterations induced by cigarette smoke in mouse lung. *Mutat Res* 2011;717:9–16.
- Graff JW, Powers LS, Dickson AM, *et al*. Cigarette smoking decreases global microRNA expression in human alveolar macrophages. *PLoS One* 2012;7:e44066.
- Gross TJ, Powers LS, Boudreau RL, *et al*. A microRNA processing defect in smokers' macrophages is linked to SUMOylation of the endonuclease DICER. *J Biol Chem* 2014;289:12823–34.
- Izzotti A, Calin GA, Arrigo P, *et al*. Downregulation of microRNA expression in the lungs of rats exposed to cigarette smoke. *FASEB J* 2009;23:806–12.
- Izzotti A, Calin GA, Steele VE, *et al*. Relationships of microRNA expression in mouse lung with age and exposure to cigarette smoke and light. *FASEB J* 2009;23:3243–50.
- Ezzie ME, Crawford M, Cho JH, *et al*. Gene expression networks in COPD: microRNA and mRNA regulation. *Thorax* 2012;67:122–31.
- Chen Y, Gelfond JA, McManus LM, *et al*. Reproducibility of quantitative RT-PCR array in miRNA expression profiling and comparison with microarray analysis. *BMC Genomics* 2009;10:407.
- Mestdagh P, Hartmann N, Baeriswyl L, *et al*. Evaluation of quantitative miRNA expression platforms in the microRNA quality control (miRQC) study. *Nat Methods* 2014;11:809–15.
- Van Peer G, Lefever S, Anckaert J, *et al*. miRBase Tracker: keeping track of microRNA annotation changes. *Database (Oxford)* 2014;2014.
- Martey CA, Pollock SJ, Turner CK, *et al*. Cigarette smoke induces cyclooxygenase-2 and microsomal prostaglandin E2 synthase in human lung fibroblasts: implications for lung inflammation and cancer. *Am J Physiol Lung Cell Mol Physiol* 2004;287:L981–91.
- Montuschi P, Kharitonov SA, Ciabattini G, *et al*. Exhaled leukotrienes and prostaglandins in COPD. *Thorax* 2003;58:585–8.
- Sato T, Liu X, Nelson A, *et al*. Reduced miR-146a increases prostaglandin E(2) in chronic obstructive pulmonary disease fibroblasts. *Am J Respir Crit Care Med* 2010;182:1020–9.
- Profita M, Sala A, Bonanno A, *et al*. Chronic obstructive pulmonary disease and neutrophil infiltration: role of cigarette smoke and cyclooxygenase products. *Am J Physiol Lung Cell Mol Physiol* 2010;298:L261–9.
- Huang SK, Wettlaufer SH, Chung J, *et al*. Prostaglandin E2 inhibits specific lung fibroblast functions via selective actions of PKA and Epac-1. *Am J Respir Cell Mol Biol* 2008;39:482–9.
- Montuschi P, KS, Carpagnano E, Culpitt S, *et al*. Exhaled prostaglandin E2: a new biomarker of airway inflammation in COPD. *Am J Respir Crit Care Med* 2000;161(A821).
- Halappanavar S, Nikota J, Wu D, *et al*. IL-1 receptor regulates microRNA-135b expression in a negative feedback mechanism during cigarette smoke-induced inflammation. *J Immunol* 2013;190:3679–86.
- Dinarello CA. Immunological and inflammatory functions of the interleukin-1 family. *Annu Rev Immunol* 2009;27:519–50.
- Doz E, Noulin N, Boichot E, *et al*. Cigarette smoke-induced pulmonary inflammation is TLR4/MyD88 and IL-1R1/MyD88 signaling dependent. *J Immunol* 2008;180:1169–78.
- Chung A, Zhou S, Wang X, *et al*. The role of interleukin-1beta in murine cigarette smoke-induced emphysema and small airway remodeling. *Am J Respir Cell Mol Biol* 2009;40:482–90.
- D'Hulst AI, Bracke KR, Maes T, *et al*. Role of tumour necrosis factor-alpha receptor p75 in cigarette smoke-induced pulmonary inflammation and emphysema. *Eur Respir J* 2006;28:102–12.
- Barh D, Malhotra R, Ravi B, *et al*. MicroRNA let-7: an emerging next-generation cancer therapeutic. *Curr Oncol* 2010;17:70–80.
- Hassan F Jr GN, ME Ezzie, P Diaz, E, *et al*. Cigarette smoke regulates the expression of the CFTR chloride channel in bronchial epithelial cells. 2011.
- Hassan F, Nuovo GJ, Crawford M, *et al*. MiR-101 and miR-144 regulate the expression of the CFTR chloride channel in the lung. *PLoS One* 2012;7:e50837.
- Zandvoort A, Postma DS, Jonker MR, *et al*. Altered expression of the Smad signalling pathway: implications for COPD pathogenesis. *Eur Respir J* 2006;28:533–41.
- Jorgensen E, Stinson A, Shan L, *et al*. Cigarette smoke induces endoplasmic reticulum stress and the unfolded protein response in normal and malignant human lung cells. *BMC Cancer* 2008;8:229.
- Dai BH, Geng L, Wang Y, *et al*. microRNA-199a-5p protects hepatocytes from bile acid-induced sustained endoplasmic reticulum stress. *Cell Death Dis* 2013;4:e604.

- 48 Su SF, Chang YW, Andreu-Vieyra C, *et al.* miR-30d, miR-181a and miR-199a-5p cooperatively suppress the endoplasmic reticulum chaperone and signaling regulator GRP78 in cancer. *Oncogene* 2013;32:4694–701.
- 49 Hassan T, Carroll TP, Buckley PG, *et al.* miR-199a-5p silencing regulates the unfolded protein response in chronic obstructive pulmonary disease and alpha1-antitrypsin deficiency. *Am J Respir Crit Care Med* 2014;189:263–73.
- 50 Chatila WM, Criner GJ, Hancock WW, *et al.* Blunted expression of miR-199a-5p in regulatory T cells of patients with chronic obstructive pulmonary disease compared to unaffected smokers. *Clin Exp Immunol* 2014;177:341–52.
- 51 Gelb AF, Hogg JC, Muller NL, *et al.* Contribution of emphysema and small airways in COPD. *Chest* 1996;109:353–9.
- 52 Christenson SA, Brandsma CA, Campbell JD, *et al.* miR-638 regulates gene expression networks associated with emphysematous lung destruction. *Genome Med* 2013;5:114.
- 53 Zhu QY, Liu Q, Chen JX, *et al.* MicroRNA-101 targets MAPK phosphatase-1 to regulate the activation of MAPKs in macrophages. *J Immunol* 2010;185:7435–42.
- 54 Mercer BA, Kolesnikova N, Sonett J, *et al.* Extracellular regulated kinase/mitogen activated protein kinase is up-regulated in pulmonary emphysema and mediates matrix metalloproteinase-1 induction by cigarette smoke. *J Biol Chem* 2004;279:17690–6.
- 55 Mercer BA, Wallace AM, Brinckerhoff CE, *et al.* Identification of a cigarette smoke-responsive region in the distal MMP-1 promoter. *Am J Respir Cell Mol Biol* 2009;40:4–12.
- 56 Hautamaki RD, Kobayashi DK, Senior RM, *et al.* Requirement for macrophage elastase for cigarette smoke-induced emphysema in mice. *Science* 1997;277:2002–4.
- 57 Mizuno S, Bogaard HJ, Gomez-Arroyo J, *et al.* MicroRNA-199a-5p is associated with hypoxia-inducible factor-1alpha expression in lungs from patients with COPD. *Chest* 2012;142:663–72.
- 58 Yasuo M, Mizuno S, Kraskauskas D, *et al.* Hypoxia inducible factor-1alpha in human emphysema lung tissue. *Eur Respir J* 2011;37:775–83.
- 59 Iyer NV, Kotch LE, Agani F, *et al.* Cellular and developmental control of O2 homeostasis by hypoxia-inducible factor 1 alpha. *Genes Dev* 1998;12:149–62.
- 60 Tang K, Rossiter HB, Wagner PD, *et al.* Lung-targeted VEGF inactivation leads to an emphysema phenotype in mice. *J Appl Physiol (1985)* 2004;97:1559–66; discussion 49.
- 61 Kasahara Y, Tuder RM, Taraseviciene-Stewart L, *et al.* Inhibition of VEGF receptors causes lung cell apoptosis and emphysema. *J Clin Invest* 2000;106:1311–9.
- 62 Brandsma CA, van den Berge M, Postma DS, *et al.* A large lung gene expression study identifying fibulin-5 as a novel player in tissue repair in COPD. *Thorax* 2015;70:21–32.
- 63 Savarimuthu Francis SM, Davidson MR, Tan ME, *et al.* MicroRNA-34c is associated with emphysema severity and modulates SERPINE1 expression. *BMC Genomics* 2014;15:88.
- 64 Liu X, Sempere LF, Galimberti F, *et al.* Uncovering growth-suppressive MicroRNAs in lung cancer. *Clin Cancer Res* 2009;15:1177–83.
- 65 Bouchie A. First microRNA mimic enters clinic. *Nat Biotechnol* 2013;31:577.
- 66 Akbas F, Coskunpinar E, Aynaci E, *et al.* Analysis of serum micro-RNAs as potential biomarker in chronic obstructive pulmonary disease. *Exp Lung Res* 2012;38:286–94.
- 67 Weber JA, Baxter DH, Zhang S, *et al.* The microRNA spectrum in 12 body fluids. *Clin Chem* 2010;56:1733–41.
- 68 Xie L, Wu M, Lin H, *et al.* An increased ratio of serum miR-21 to miR-181a levels is associated with the early pathogenic process of chronic obstructive pulmonary disease in asymptomatic heavy smokers. *Mol Biosyst* 2014;10:1072–81.
- 69 Vencken SF, Greene CM, McKiernan PJ. Non-coding RNA as lung disease biomarkers. *Thorax* 2015;70:501–3.
- 70 Pinkerton M, Chinchilli V, Banta E, *et al.* Differential expression of microRNAs in exhaled breath condensates of patients with asthma, patients with chronic obstructive pulmonary disease, and healthy adults. *J Allergy Clin Immunol* 2013;132:217–19.
- 71 Volders PJ, Helsens K, Wang X, *et al.* LNCipedia: a database for annotated human lncRNA transcript sequences and structures. *Nucleic Acids Res* 2013;41(Database issue):D246–51.
- 72 Rinn JL, Kertesz M, Wang JK, *et al.* Functional demarcation of active and silent chromatin domains in human HOX loci by noncoding RNAs. *Cell* 2007;129:1311–23.
- 73 Thai P, Statt S, Chen CH, *et al.* Characterization of a novel long noncoding RNA, SCAL1, induced by cigarette smoke and elevated in lung cancer cell lines. *Am J Respir Cell Mol Biol* 2013;49:204–11.
- 74 Beane J, Vick J, Schembri F, *et al.* Characterizing the impact of smoking and lung cancer on the airway transcriptome using RNA-Seq. *Cancer Prev Res (Phila)* 2011;4:803–17.
- 75 Hackett NR, Butler MW, Shaykhiev R, *et al.* RNA-Seq quantification of the human small airway epithelium transcriptome. *BMC Genomics* 2012;13:82.
- 76 Itoh K, Wakabayashi N, Katoh Y, *et al.* Keap1 regulates both cytoplasmic-nuclear shuttling and degradation of Nrf2 in response to electrophiles. *Genes Cells* 2003;8:379–91.
- 77 Rangasamy T, Cho CY, Thimmulappa RK, *et al.* Genetic ablation of Nrf2 enhances susceptibility to cigarette smoke-induced emphysema in mice. *J Clin Invest* 2004;114:1248–59.
- 78 Suzuki M, Betsuyaku T, Ito Y, *et al.* Down-regulated NF-E2-related factor 2 in pulmonary macrophages of aged smokers and patients with chronic obstructive pulmonary disease. *Am J Respir Cell Mol Biol* 2008;39:673–82.
- 79 Cabili MN, Trapnell C, Goff L, *et al.* Integrative annotation of human large intergenic noncoding RNAs reveals global properties and specific subclasses. *Genes Dev* 2011;25:1915–27.
- 80 Bi H, Zhou J, Wu D, *et al.* Microarray analysis of long non-coding RNAs in COPD lung tissue. *Inflamm Res* 2015;64:119–26.
- 81 Makita H, Nasuhara Y, Nagai K, *et al.* Characterisation of phenotypes based on severity of emphysema in chronic obstructive pulmonary disease. *Thorax* 2007;62:932–7.
- 82 Lin CW, Chang YL, Chang YC, *et al.* MicroRNA-135b promotes lung cancer metastasis by regulating multiple targets in the Hippo pathway and LZTS1. *Nat Commun* 2013;4:1877.