of AEs in >2% of patients was comparable in the tiotropium Respimat[®] 5 µg, tiotropium Respimat[®] 2.5 µg and placebo Respimat[®] groups (Table 1). No deaths occurred. 110 (5.7%) and 55 (4.4%) patients receiving tiotropium Respimat[®] and placebo Respimat[®], respectively, reported drug-related AEs (cardiac AEs were rare: tiotropium Respimat[®], 7 [0.4%]; placebo Respimat[®], 3 [0.2%]). One drug-related serious AE (asthma) was reported with tiotropium Respimat[®].

Abstract P149 Table 1 Frequency of AEs occurring in >2% of patients

	AEs occurring in >2% of patients, n (%) ^a		
	Tiotropium Respimat [®] 5 µg (n = 1256)	Tiotropium Respimat [®] 2.5 µg (n = 673)	Placebo Respimat [®] (n = 1260)
Exposure, patient-years	705.42	271.08	708.04
Any AE	732 (58.3)	350 (52.0)	772 (61.3)
Serious AEs	51 (4.1)	12 (1.8)	56 (4.4)
AEs by preferred term			
Asthma	326 (26.0)	106 (15.8)	384 (30.5)
Decreased PEF rate	158 (12.6)	58 (8.6)	207 (16.4)
Nasopharyngitis	98 (7.8)	51 (7.6)	118 (9.4)
Upper respiratory tract infection	49 (3.9)	29 (4.3)	67 (5.3)
Bronchitis	43 (3.4)	9 (1.3)	27 (2.1)
Headache	41 (3.3)	19 (2.8)	49 (3.9)
Sinusitis	31 (2.5)	17 (2.5)	33 (2.6)
Influenza	29 (2.3)	1 (0.1)	25 (2.0)

^aTreated set. PrimoTinA-asthma[®]: NCT00776984/NCT00772538; MezzoTinA-asthma[®]: NCT01172808/NCT01172821; GraziaTinA-asthma[®]: NCT01316380; Study 342: NCT00350207.

Conclusion Once-daily tiotropium Respimat[®] add-on to at least ICS maintenance therapy in adult patients demonstrates a safety profile comparable with that of placebo and is well tolerated across severities of symptomatic asthma.

P150 ONCE-DAILY TIOTROPIUM RESPIMAT[®] REDUCES RISK OF SEVERE ASTHMA EXACERBATION AND ASTHMA WORSENING IN SYMPTOMATIC ASTHMA, INDEPENDENT OF ALLERGIC AND INFLAMMATORY STATUS

¹R Dahl, ²T Casale, ³M Vandewalker, ⁴H Schmidt, ⁵M Engel, ⁵P Moroni-Zentgraf, ⁶HAM Kerstjens. ¹Allergy Centre, Odense University Hospital, Odense, Denmark; ²University of South Florida Morsani College of Medicine, Tampa, Florida, USA; ³Clinical Research of the Ozarks, Columbia, Missouri, USA; ⁴Global Biometrics and Clinical Applications, Boehringer Ingelheim Pharma GmbH & Co. KG, Biberach an der Riss, Germany; ⁵TA Respiratory Diseases, Boehringer Ingelheim Pharma GmbH & Co. KG, Ingelheim am Rhein, Germany; ⁶University of Groningen, Department of Pulmonary Medicine, University Medical Center Groningen, Groningen, The Netherlands

10.1136/thoraxjnl-2015-207770.287

Background Four trials explored whether tiotropium Respimat[®] add-on to at least ICS is effective in the T_H2 phenotype, determined by high serum immunoglobulin E (IgE) and blood eosinophil values, in reducing risk of severe asthma exacerbation and asthma worsening in adult patients with moderate or severe symptomatic asthma.

Methods Four Phase III, double-blind, placebo-controlled, parallel-group trials. PrimoTinA-asthma[®] (two 48-week trials;

NCT00776984/NCT00772538; n = 912); tiotropium Respimat[®] 5 µg or placebo Respimat[®] add-on to ICS + LABA (≥800 µg budesonide or equivalent); MezzoTinA-asthma[®] (two 24-week trials; NCT01172808/NCT01172821; n = 2100): tiotropium Respimat[®] 5 µg, tiotropium Respimat[®] 2.5 µg or placebo add-on to ICS (400–800 µg budesonide or equivalent). Patients had symptomatic asthma requiring treatment with at least ICS for ≥4 weeks before screening; COPD was excluded. Subgroups of allergic and inflammatory status (IgE and eosinophils) were used to analyse risk of severe exacerbation and asthma worsening, *post hoc*. Cox regression modelling analyses, adjusted for treatment, IgE or eosinophils and treatment by IgE or eosinophil interaction, were applied to calculate hazard ratios and 95% confidence intervals across IgE (2–2000 µg/L) and eosinophil (0.05–7.00 × 10⁹/L) values.

Results Severe exacerbation: in PrimoTinA-asthma[®], tiotropium Respimat[®] 5 µg reduced risk in terms of hazard ratio versus placebo Respimat[®] up to an IgE level of ~1000 µg/L, and consistently across all eosinophil values. In MezzoTinA-asthma[®], tiotropium Respimat[®] 5 µg and 2.5 µg reduced risk versus placebo consistently across all IgE and eosinophil levels. Asthma worsening: in PrimoTinA-asthma[®], tiotropium Respimat[®] 5 µg reduced risk in terms of hazard ratio versus placebo Respimat[®], independent of IgE and eosinophils. In MezzoTinA-asthma[®], tiotropium Respimat[®] 5 µg reduced risk versus placebo across all IgE and eosinophil values. Tiotropium Respimat[®] 2.5 µg reduced risk versus placebo across all IgE values and at eosinophil values <3.00 × 10⁹/L.

Conclusion Tiotropium Respimat[®] add-on to ICS ± LABA reduces risk of severe exacerbation and asthma worsening in patients across severities of symptomatic asthma and a broad range of IgE and eosinophil values, suggesting efficacy independent of underlying allergic/eosinophilic inflammation. Once-daily tiotropium Respimat[®] may have potential as add-on to at least ICS maintenance therapy in patients with symptomatic asthma, independent of T_H2 phenotype.

P151 TIOTROPIUM RESPIMAT[®] ADD-ON THERAPY REDUCES EXACERBATION RISK IN PATIENTS WITH MODERATE OR SEVERE SYMPTOMATIC ASTHMA, INDEPENDENT OF TH2 STATUS

¹T Casale, ²R Dahl, ³JC Virchow, ⁴M Engel, ⁴P Moroni-Zentgraf, ⁵R Lühmann, ⁶HAM Kerstjens. ¹Division of Allergy and Immunology, University of South Florida Morsani College of Medicine, Tampa, Florida, USA; ²Odense University Hospital, Odense, Denmark; ³Department of Pneumology, Intensive Care Medicine, Zentrum Für Innere Medizin, Klinik I, University Clinic Rostock, Rostock, Germany; ⁴TA Respiratory Diseases, Boehringer Ingelheim Pharma GmbH & Co. KG, Ingelheim am Rhein, Germany; ⁵Global Biometrics and Clinical Applications, Boehringer Ingelheim Pharma GmbH & Co. KG, Biberach an der Riss, Germany; ⁶University of Groningen, Department of Pulmonary Medicine, University Medical Center Groningen, Groningen, The Netherlands

10.1136/thoraxjnl-2015-207770.288

Background Phase III studies have demonstrated reduced exacerbation rates with tiotropium Respimat[®] (tioR) add-on to ICS + LABA in patients with symptomatic asthma (Kerstjens *et al.* NEJM 2012;367:1198–207). There are currently no reported specific treatments for asthma that work equally well in both T_H2-low and T_H2-high phenotypes. We explored, in patients with moderate or severe symptomatic asthma, whether T_H2 status influenced tioR responses, assessed by time to first exacerbation.

Methods In two 48-week trials (PrimoTinA-asthma[®]: NCT00776984/NCT00772538), patients on ICS + LABA (≥800

µg budesonide or equivalent) received once-daily tioR 5 µg or placebo Respimat®. In two 24-week trials (MezzoTinA-asthma®: NCT01172808/NCT01172821), patients on ICS (400–800 µg budesonide or equivalent) received once-daily tioR 5 µg or 2.5 µg, twice-daily salmeterol 50 µg via hydrofluoroalkane metered-dose inhaler (active comparator) or placebo (identical devices in a double-dummy protocol). Pre-planned analyses (pooled data) of time to first severe exacerbation and time to first episode of asthma worsening were performed in T_H2-low and T_H2-high subgroups: total serum immunoglobulin (IgE) ≤ or >430 µg/L (179.2 IU/L); blood eosinophils ≤ or >0.6 × 10⁹/L (600/µL).

Results 912 patients with severe asthma received tioR 5 µg or placebo Respimat®: 205/182 were reported with IgE >430 µg/L and 99/87 with an eosinophil count of >0.6 × 10⁹/L. 2100 patients with moderate asthma received tioR 5 µg or 2.5 µg, salmeterol or placebo: 319/320/319/326 were reported with IgE >430 µg/L and 104/103/111/107 with an eosinophil count of >0.6 × 10⁹/L. Time to first severe exacerbation was longer with tioR versus placebo (Table 1) in patients with severe or moderate asthma, independent of IgE and eosinophils (interaction p values [Cox regression] 0.169 and 0.754, respectively, for PrimoTinA-asthma®; analyses not performed for MezzoTinA-asthma® because of low incidence of severe exacerbations). Time to first asthma worsening was longer with tioR versus placebo (Table 1) in patients with moderate or severe asthma, independent of IgE (interaction p values 0.998 [PrimoTinA-asthma®] and 0.041 [MezzoTinA-asthma®]) and eosinophils (interaction p values 0.251 [PrimoTinA-asthma®] and 0.125 [MezzoTinA-asthma®]).

Abstract P151 Table 1 Risk of severe asthma exacerbation and asthma worsening in PrimoTinA-asthma® and MezzoTinA-asthma®

	All comparisons versus placebo Respimat® or placebo HFA-MDI, hazard ratio			
	Serum IgE (µg/L)		Eosinophils (×10 ⁹ /L)	
	≤430	>430	≤0.6	>0.6
Time to first severe asthma exacerbation				
PrimoTinA-asthma® (ICS + LABA)				
TioR 5 µg	0.75	1.07	0.81	0.75
QD	p = 0.162	p = 0.692	p = 0.162	p = 0.218
MezzoTinA-asthma® (ICS)				
TioR 5 µg	0.86	0.61	0.65	1.04
QD ^b	p = 0.691	p = 0.107	p = 0.099	p = 0.953
TioR 2.5 µg	0.51	0.50	0.41	1.25
QD ^b	p = 0.119	p = 0.033	p = 0.003	p = 0.715
Salmeterol ^c	0.82	0.68	0.62	1.76
	p = 0.594	p = 0.189	p = 0.066	p = 0.310
Time to first episode of asthma worsening^a				
PrimoTinA-asthma® (ICS + LABA)				
TioR 5 µg	0.73	0.73	0.65	0.85
QD	p = 0.030	p = 0.017	p < 0.001	p = 0.360
MezzoTinA-asthma® (ICS)				
TioR 5 µg	0.88	0.83	0.90	0.70
QD ^b	p = 0.495	p = 0.220	p = 0.410	p = 0.170
TioR 2.5 µg	0.45	0.81	0.60	0.91
QD ^b	p < 0.001	p = 0.157	p < 0.001	p = 0.691
Salmeterol ^c	0.60	0.84	0.71	0.92
	p = 0.009	p = 0.231	p = 0.009	p = 0.714

^aDefined as either a progressive increase in symptoms or a decline of ≥30% in best morning PEF for ≥2 consecutive days; ^bPlus placebo HFA-MDI BID; ^cSalmeterol HFA-MDI 50 µg BID plus placebo Respimat® QD.
BID, twice-daily; HFA-MDI, hydrofluoroalkane metered-dose inhaler; QD, once-daily; tioR, tiotropium Respimat®.

Conclusion Once-daily tiotropium Respimat® add-on to at least ICS reduced the risk of severe exacerbation and asthma worsening in patients with moderate or severe symptomatic asthma, independent of T_H2 phenotype.

P152 FLUTICASONE FUROATE (FF)/VILANTEROL (VI) ONCE DAILY REDUCES ASTHMA SYMPTOMS BOTH DAY AND NIGHT

¹DA Leather, ²R Forth, ¹L Yates, ³LA Jacques. ¹GlaxoSmithKline Respiratory Global Franchise, London, UK; ²PAREXEL International Research Triangle Park, Durham, USA; ³Respiratory Medicines Development Centre, GlaxoSmithKline, London, UK

10.1136/thoraxjnl-2015-207770.289

Introduction and objectives FF/VI is the first once daily inhaled corticosteroid/long-acting b₂-agonist combination available for the treatment of asthma. Results from five phase III studies that have previously been presented demonstrated a sustained 24 h improvement in lung function and improvement in symptom-free 24 h periods.

Methods Post-hoc analyses of diary card data from these studies were performed to examine whether there was any difference in the contribution of the day and night time symptom-free period to the 24 h symptom-free period. The diary card scale used is described below.

Day-time Symptom Score:

0 = No symptoms during the day

1 = Symptoms for one short period during the day

2 = Symptoms for two or more short periods during the day

3 = Symptoms for most of the day which did not affect my normal daily activities

4 = Symptoms for most of the day which did affect my normal daily activities

5 = Symptoms so severe that I could not go to work or perform normal daily activities

Night-time Symptom Score:

0 = No symptoms during the night

1 = Symptoms causing me to wake once (or wake early)

2 = Symptoms causing me to wake twice or more (including waking early)

3 = Symptoms causing me to be awake for most of the night

4 = Symptoms so severe that I did not sleep at all

To be counted as symptom-free during the day or night the patient needed to record a score of 0.

Results The post-hoc analyses demonstrated that the improvements in day and night time symptom-free periods were similar to the 24 h symptom free periods. See Figure 1 below.

Conclusions In general benefits in symptom free days and symptom free nights contributed to the benefit of FF/VI over comparator groups in terms of 24 h symptom free periods.

P153 FLUTICASONE PROPIONATE/FORMOTEROL PRESSURISED METERED-DOSE INHALER '2-3-4' TRAINING PARADIGM AIDS CORRECT INHALER TECHNIQUE

¹D Bell, ²L Mansfield, ²M Lomax, ²S Dissanayake. ¹Bio-Kinetic Europe Limited, Belfast, UK; ²Mundipharma Research Limited, Cambridge, UK

10.1136/thoraxjnl-2015-207770.290

Background Inhaler technique is crucial to effective disease control. Amongst the most frequent mistakes made with all inhalers are the failure to exhale adequately, an insufficient breath-hold