IMAGES IN THORAX

Nitinol stent insertion in tracheomalacia

Carmen Pizarro,¹ Darius Dabir,² Georg Nickenig,¹ Dirk Skowasch¹

¹Department of Internal Medicine II, Cardiology, Pneumology and Angiology, University Hospital Bonn, Bonn, Germany
²Department of Radiology, University Hospital Bonn, Bonn, Germany

Correspondence to

Dr Carmen Pizarro,
 Department of Internal Medicine II, Cardiology,
 Pneumology and Angiology,
 University Hospital Bonn,
 Sigmund-Freud-Straße 25,
 Bonn 53105, Germany;
 carmen.pizarro@ukb.uni-bonn.de

Received 23 January 2016
 Accepted 15 February 2016
 Published Online First
 4 March 2016

A 69-year-old man was admitted to our hospital with progressive dyspnoea accompanied by inspiratory stridor. He presented a history of surgical aortic valve replacement 3 years previously, complicated by prolonged postoperative mechanical ventilation requiring percutaneous tracheostomy. Presently, bronchoscopy evidenced tracheomalacia (TM) with subtotal tracheal stenosis. CT scan confirmed long segment narrowing >20 mm of the upper trachea (figure 1). Operative retracheostomy

resulted in repeated tube dislocation incl. necessity for cardiopulmonary resuscitation. As cardiac comorbidities additionally impeded extensive surgical repair, insertion of an uncovered, self-expanding nitinol stent (stent diameter: 18 mm, stent length: 40 mm, individually manufactured by Leufen Medical GmbH, Berlin, Germany) was performed (figure 2, left). No intraprocedural complications occurred post stent deployment. For 2 years by now, clinical and bronchoscopic patient follow-up

Figure 1 CT scan images in coronal (A) and axial (B) projection showing 20 mm subtotal stenosis of the upper trachea.

CrossMark

To cite: Pizarro C, Dabir D, Nickenig G, et al. *Thorax* 2016;**71**:770–771.

Figure 2 Endotracheal imaging during bronchoscopy on the day following nitinol stent implantation (left) and 25 months thereafter (right). Though formation of granulation tissue has meanwhile occurred, tracheomalacia-driven tracheal renarrowing is persistently prevented.

revealed durable adequate stent expansion and stenting of the upper trachea (figure 2, right).

TM is a condition characterised by excessive diffuse or segmental tracheal collapsibility that can be classified as congenital or acquired. Among the latter, tracheostomy may provoke tracheal cartilage injury and thereby induce TM. As to the scarce therapeutic approaches in symptomatic TM, insertion of silicone stents is preferentially employed, though stent-related complications like migration, plugging or formation of granulation tissue are frequent.¹ In contrast, metal stents are currently not recommended for alleviation of benign tracheal narrowing due to their potential risk of breakage. However, we presently report

sustained efficiency of airway stabilisation by long-term nitinol stenting that might evolve into an equivalent treatment option in TM.

Contributors CP, DD, GN and DS drafted, conducted and coreported this article.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCE

- 1 Murgu SD, Colt HG. Complications of silicone stent insertion in patients with expiratory central airway collapse. *Ann Thorac Surg* 2007;84:1870–7.