cell surface activation similar to other agents. Comparing challenges in chronic cough patients shows that whilst inter-patient variability is low, intra-patient variability is high. Therefore, in contrast to healthy subjects, cough challenges are unlikely to be a useful measure of determining individual improvement in chronic cough patients.

Beyond FEV₁ in COPD

S32 EPIDEMIOLOGY OF CHRONIC OBSTRUCTIVE PULMONARY DISEASE (COPD) IN THE UK: FINDINGS FROM THE BRITISH LUNG FOUNDATION'S 'RESPIRATORY HEALTH OF THE NATION' PROJECT

¹N Snell, ²D Strachan, ³R Hubbard, ³J Gibson, ⁴K Gruffydd-Jones, ¹I Jarrold. ¹British Lung Foundation, London, UK; ²St George's, University of London, London, UK; ³University of Nottingham, Nottingham, UK; ⁴Box Surgery, Corsham, UK

10.1136/thoraxjnl-2016-209333.38

Background We performed an analysis of UK respiratory disease epidemiology covering 2004–2012. Findings pertaining to COPD are presented here.

Methods Prevalence and incidence rates were estimated from The Health Intelligence Network database representing ~5 per cent of the population. Mortality figures came from official government statistics. WHO data were used for international mortality comparisons and numbers of hospital admissions/inpatient bed-days.

Results An estimated 1.2 million people (2% of the population) have diagnosed COPD – considerably more than the 835,000 estimated by the Department of Health in 2011 – making COPD the second most common lung disease in the UK, after asthma. Prevalence has increased by 27% in the last decade. Incidence fell 2004–2008 but has been stable since with just under 115,000 new diagnoses in 2012. Men are more likely to be diagnosed with COPD and to die from it than women. COPD is rare under 40 and becomes commoner with age, affecting 9% of those aged >70. COPD prevalence, incidence and mortality rates are highest in Scotland and the north of England. Prevalence and incidence are over twice as great in the most deprived population quintile than in the least. Nearly 30,000 people die from COPD each year, making it the second greatest cause of death from lung disease and the UK's fifth biggest killer. Mortality increased from 2004–2012. The UK COPD mortality rate ranks third in Europe. COPD accounts for over 140,000 hospital admissions and over a million bed days each year across the UK (1.7% of all hospital admissions and bed days). 97% of these admissions are for emergency care. London has notably more hospital admissions for COPD than other regions with similar prevalence.

Conclusions Gender, location, and deprivation differences in COPD epidemiology probably largely reflect differences in smoking behaviour. COPD continues to represent an enormous burden on UK health services.

S33

PHYSICAL ACTIVITY INTERVENTION VERSUS PULMONARY REHABILITATION IN COPD: THE LIVELY COPD PROJECT

¹B O'Neill, ¹OM O'Shea, ²SM McDonough, ¹L McGarvey, ¹I Bradbury, ³MA Arden, ⁴T Troosters, ⁵D Cosgrove, ⁶T McManus, ⁷TJ McDonnell, ⁸JM Bradley. ¹Ulster University, Belfast, UK; ²Ulster University and UK CRC, Centre for Excellence for Public Health, Belfast, UK; ³Sheffield Hallam University, Sheffield, UK; ⁴Katholieke University, Leuven, Belgium; ⁵Northern Irish Clinical Research Network, Belfast, UK; ⁶Department of Respiratory Medicine, Western Health and Social Care Trust, Enniskillen, UK; ⁷St Vincent's University Hospital, Dublin, Ireland; ⁸Queen's University, Belfast, UK

10.1136/thoraxjnl-2016-209333.39

Introduction and objectives The format of pulmonary rehabilitation (PR) may not meet the needs of all patients with COPD or lead to improved physical activity (PA) levels. Drop outs from PR can be high. A pedometer driven physical activity intervention (PAI) may offer patients an alternative method for increasing their PA. The aim of this study was to assess the feasibility of a 12 week clinician facilitated PAI versus PR in people with COPD. **Methods** The design was a multicenter-randomised, parallel-group, feasibility study. Patients with COPD referred for PR were included. Spirometry and demographics were recorded. The following were assessed at baseline, post-intervention and follow up (12 weeks): PA using an ActiGraph GT3X+ accelerometer, sealed Yamax Digiwalker pedometer and the International Physical Activity Questionnaire (IPAQ) (long form); exercise capacity (Incremental Shuttle Walk Test (ISWT)); COPD Assessment Test (CAT). Recruitment, retention and completion/rates were recorded. Descriptive statistics and mean differences were used to analyse the data.

Results 50 patients (mean (SD) age 64 (8) years, 24M, FEV₁ 1.44 (0.63)) were recruited and randomised: PR n = 26, PAI n = 24. Of those screened 50/651, 13% were recruited. One participant randomised to the PAI started PR, a per protocol analysis was conducted; PR n = 27 and PAI n = 23. Completion of the PAI was 74% (17/23) and PR was 48% (13/27). Retention overall was 74% post-intervention (n = 18 PAI; n = 19 PR), and 66% at follow up (n = 15 PAI; n = 18 PR). There was a mean (95% confidence interval (CI)) change of 972 (–1080 to 3024) steps/day and 4 (–441 to 449) in the PAI and PR group respectively; results are found in Table 1. There were 4 minor adverse events (PAI n = 3 PR n = 1).

Conclusions This study will inform a future large scale randomised control trial (RCT). The LIVELY PAI intervention appears to be feasible and safe within this preliminary study, and enhanced physical activity in people with COPD. While the results require confirmation in a fully powered RCT, the mean increase in step count is in line with a recently published minimally clinically important difference.¹

REFERENCE

- Demeyer H, et al. The Minimal Important Difference in Physical Activity in Patients with COPD. *PLoS One*. 2016;11:0154587.