Method Prospective patients (2013–15 cohort) were identified from ward lists. Retrospective patients (2009–12 cohort) were identified from haemato-oncology patients who had had blood cultures and were co-incidentally found to be neutropenic. Medical notes, drug charts, discharge letters and microbiology results were reviewed.

Results The 2 cohorts were well matched in terms of haemato-oncology diagnosis. The 2009/12 cohort included 561 episodes, with 333 CT chests and 62 bronchoscopies compared to 508 episodes, 288 CT chests and 86 bronchoscopies in 2013/15 group. 12 month mortality was 42% in 2009/12 versus 34% in 2013/15 cohort. 30 day mortality was 11% for both cohorts. There was no significant difference in length of antifungal treatments, although 24% switched to voriconazole following positive biomarkers.

Concordance between serum and BAL GM was 14.8% and APCR was 6.3%. Concordance between serum GM and APCR was 7% and between BAL GM and PCR was 41%.

Confidence in diagnosis of IFD increased with the novel pathway See Figure 1.

Conclusion The introduction of a novel NF pathway was found to be non-inferior in terms of 12 month and 30 day mortality. Although there was increased confidence in the diagnosis of IFD, this did not translate to reduced antifungal treatment, although it did influence switching to voriconazole and secondary chemoprophylaxis. Negative serum GM and PCR did not rule out the diagnosis of IFD and BAL biomarkers were more sensitive than serum ones.

REFERENCE

- 1 De Pauw B, et al. *Clin Infect Dis* 2008;**46**(12).

P267 DLCO PREDICTS DISEASE SEVERITY AND MORTALITY IN BRONCHIECTASIS

MJ McDonnell, M O'Mahony, D Breen, JJ Gilmartin, A O'Regan, RM Rutherford. *Galway University Hospitals, Galway, Ireland*

10.1136/thoraxjnl-2016-209333.410

Background Few studies have assessed the role of lung diffusing capacity (DLCO%) in bronchiectasis. We sought to examine the relationship between DLCO% and clinical and radiological variables in a well-defined population of bronchiectasis patients to determine its potential prognostic significance and compare with FEV₁%.

Methods Of 312 consecutive bronchiectasis patients attending our institution over a 3-year period, 204 patients were suitable for study inclusion. Exclusion criteria consisted of patients with cystic fibrosis or traction bronchiectasis, patients with missing data and patients with absent radiological evidence of bronchiectasis on independent expert thoracic radiologist review. Univariate analyses was performed using Pearson's correlation. Backwards stepwise logistic regression analysis was subsequently performed to determine independent associations of DLCO% and FEV₁%.

Results DLCO% strongly correlated with all measured lung function parameters FEV₁%, FVC%, FEF 25–75% and FEV₁/FVC ratio (all $p < 0.001$). Negative correlations were noted with age at diagnosis ($p = 0.047$), body mass index ($p < 0.001$) and number of comorbidities ($p = 0.002$). Significant symptom associations included SOB ($p = 0.001$) and fatigue ($p < 0.001$).

Reduced DLCO% was associated with higher MRCD scores ($p < 0.001$), higher number of hospitalisations on follow up ($p = 0.015$), higher BSI scores ($p < 0.001$) and increased mortality ($p = 0.028$). No correlations were noted with gender, smoking history, aetiology or bacterial colonisation of any form. Reduced DLCO% was associated with increased number of lobes ($p = 0.004$), and higher total Reiff and modified Bhalla HRCT scores ($p = 0.001$ and $p < 0.001$ respectively). The modified Bhalla was excluded from stepwise regression as unavailable in most clinical settings. Backwards elimination showed DLCO% to be significant in predicting BSI ($p < 0.001$), number of lobes (0.017) and mortality ($p = 0.028$). Comparatively, backwards stepwise regression of FEV₁% showed significance in predicting BSI ($p < 0.001$), number of hospitalisations on follow up ($p = 0.008$), and mortality ($p = 0.024$). Separate regression models of DLCO% and FEV₁% using modified Bhalla components as cofactors showed DLCO% to be associated with disease extent ($p = 0.002$), bronchial wall thickness ($p < 0.001$), bronchial wall dilatation ($p < 0.001$) and reduced parenchymal attenuation ($p = 0.015$); FEV₁% was associated with bronchial wall dilatation only ($p < 0.001$).

Conclusion DLCO% predicts radiological disease, disease severity and mortality in bronchiectasis, independently of aetiology, and may identify patients with advanced disease who could benefit from intensive management.

P268 IS BRONCHIECTASIS SEVERITY INFLUENCED BY AETIOLOGY OR CO-MORBID AIRWAYS DISEASE?

TM Quinn, AT Hill. *Royal Infirmary and University of Edinburgh, Edinburgh, UK*

10.1136/thoraxjnl-2016-209333.411

Background There is increased interest whether aetiology and co-morbid airways disease influence bronchiectasis disease severity.

Methods We conducted a retrospective study of 400 patients attending a specialist bronchiectasis clinic in NHS Lothian, Edinburgh, UK between May 2013 and September 2014 and using multivariable models we identified independent risk factors that influenced bronchiectasis disease severity using the Bronchiectasis Severity Index. We adjusted for age, sex, smoking history, aetiology and presence of co-morbid airways disease (asthma and COPD).

Results 400 patients were included in this study. The mean age was 66.0 (13.9) years. 253 (63.2%) were female. The majority (77%) had idiopathic (53%) or post infective bronchiectasis (24%). Other aetiologies were: allergic bronchopulmonary aspergillosis 8%; immune/auto-immune 6%; interstitial lung disease 3%; ciliary defects 3%; and inflammatory bowel disease 3%. Co-existing airways disease was common but not the predominant diagnosis (36% had asthma and 19% COPD).

Independent risk factors for severe bronchiectasis (BSI ≥ 9) were age 70–79 (OR 6.3, $p = 0.003$), age 80 and above (OR 7.3, $p = 0.003$) and smoking (OR 1.02, $p = 0.002$). It was not influenced by presence of airways disease or aetiology.

Conclusion In conclusion, neither aetiology nor presence of airways disease was independent risk factors for severe bronchiectasis severity. Age was the strongest independent predictor for severe bronchiectasis severity.