were recorded at baseline, 2 weeks and 3 months. Feasibility outcomes included recruitment, retention and study form completion rates. A patient and carer focus group was held to discuss the intervention and the trial process.

Results Recruitment rate was 50% of those assessed for eligibility (See Figure). Questionnaire completion rates were excellent with very few missing data. 24 participants were male and 38 female. Median age was 63 years (range 18–82).

Participants reported using the information provided, and feedback was positive, particularly highlighting the usefulness of the video clips and self-management information. The most popular pages of the website included diet and lifestyle advice, advice for carers, and symptoms and prognosis. Focus group data reinforced users' positive experiences of the resource and the trial process. Full analysis is ongoing.

Discussion and conclusions We have co-developed a multi-format, accessible information resource that could be made widely available outside of the specialist clinic setting. The BRIEF study suggests larger, definitive studies using interventions to improve understanding, compliance and self-management are warranted. Full results of the feasibility study are expected by December 2016.

Abstract M15 Figure 1

REFERENCES

- Hester KLM, Newton J, De Soyza A, *et al.* Living your life with bronchiectasis: an exploration of patients and carers information needs informing development of a novel information resource. *Thorax* 2015;**70**(suppl. 3):A178.
- Hester KLM, Newton J, Rapley T, *et al.* Evaluation of a novel information resource for patients with bronchiectasis: study protocol for a randomised controlled trial. *Trials* 2016;**17**:210.

M16 CONSTRUCT VALIDITY OF THE NEEDS ASSESSMENT TOOL PROGRESSIVE DISEASES FOR INTERSTITIAL LUNG DISEASE (NAT: PD-ILD) PATIENTS

¹C Reigada, ²C Fairhurst, ³J Yorke, ⁴J Ross, ¹J Boland, ⁵S Hart, ⁶D Currow, ³G Grande, ⁷S Bajwah, ⁴A Wells, ¹U Macleod, ²M Bland, ¹M Johnson. ¹Hull York Medical School, Hull, UK; ²University of York, York, UK; ³University of Manchester, Manchester, UK; ⁴Royal Brompton and Harefield NHS Foundation Trust, London, UK; ⁵Hull and East Yorkshire Hospitals NHS Foundation Trust, Hull, UK; ⁶Flinders University, Australia; ⁷Cicely Saunders Institute, London, UK

10.1136/thoraxjnl-2016-209333.458

Background People with ILD, currently have less access to SPC and there is no validated needs assessment tool (NAT). We

adapted the NAT:PD-cancer for use in ILD and conducted psychometric testing.

Aim To test the construct validity of NAT:PD-ILD.

Methods ILD clinicians in four hospitals were trained to use the NAT:PD-ILD. After a consultation, the clinician completed the NAT:PD-ILD, patients completed the St. George's Respiratory Questionnaire (SGRQ-I) and carers completed the Carer Strain Index (CSI) and Carer Support Needs Assessment Tool (CSNAT).

Kendall's Tau-b correlation coefficient (and associated p-value) was calculated to determine the correlation between the NAT: PD-ILD items relating to patient wellbeing, and a total score for a subset of SGRQ-I questions identified *a priori* as measuring similar constructs. The prevalence and bias adjusted kappa (PABAK), Cohen's kappa and percentage of agreement were used to assess whether responses were similar between the NAT: PD-ILD items relating to the ability and wellbeing of the carer and appropriate CSI and CSNAT items which were considered to measure similar concerns/support needs.

Results A total of 68 patients were recruited. The average age of participating patients was 66 years (range 34 to 87) and 62% were male. Forty-five (66%) patients had a carer of whom 27 completed the CSI (mean 4.4, SD 3.0, median 4, range 0–11) and 29 completed at least one item of the CSNAT.

Items 2, 3, 5 and 6 of the NAT: PD-ILD statistically significantly positively correlated with their comparator SGRQ-I scores (ρ range 0.24 to 0.36, $p < 0.05$). PABAK values comparing the NAT: PD-ILD items with appropriate CSI and CSNAT items show most items have PABAK positive values (range from 0.04 to 0.57, with a minimum of 52% agreement). However, NAT:PD-ILD items 11 and 13 have negative PABAK values (Inter-personal relationships and Grief topics – Psychosocial Dimension).

Conclusion The NAT: PD-ILD has adequate construct validity for most domains. However, agreement is poor for physical symptoms and spiritual concerns. This may indicate that clinicians identify concerns with symptoms less well unless they are severe.

M17 LIMITED VALUE OF BASELINE CHEST RADIOGRAPHY IN ADULTS WITH NON-TUBERCULOUS MYCOBACTERIA

¹ME Murphy, ²NM Shah, ³T Bharucha, ³C Cash, ³JR Cleverley, ³IM Cropley, ³S Hopkins, ¹MCI Lipman. ¹University College London, London, UK; ²Kings College London, London, UK; ³Royal Free London NHS Foundation Trust, London, UK

10.1136/thoraxjnl-2016-209333.459

Chest radiographic changes are associated with mycobacterial burden, treatment response and outcome in patients with tuberculosis. There is a paucity of similar data for non-tuberculous mycobacteria (NTM). We describe the chest radiology (CXR) findings in a cohort of adults without cystic fibrosis.

Methods Patients with NTM isolated from respiratory specimens between 2010–2013 at our centre were reviewed. Chest X-rays (CXR) nearest the date of positive NTM culture were read independently by two consultant Radiologists for 5 categories of abnormality (nodules, cavities, bronchiectasis, bronchial wall thickness [BWT] and consolidation) in each of 6 zones. A consensus result was agreed where discrepant. CXR results were recorded as "normal" or "abnormal" overall and for each category per zone. The total number of zones affected in all categories was summed to provide a measure of radiological extent of disease (with a maximum score of 30), e.g. a patient with cavitation in 2 zones and bronchiectasis in 3 would score 5/30. Results