Methods We conducted a retrospective review of patients diagnosed with new PE at Milton Keynes hospital between August 2014 and 2015 to determine the proportion of patients found to have malignancy after CT. Selected patient notes were interrogated for clinical and laboratory findings at the time of diagnosis, and for details of subsequent management.

Results 177 patients were included in our study. 102 received a CT A/P, 88 of whom did not have an established diagnosis of cancer. Out of the 88, 5 new diagnoses of cancer were made. In 10 cases, CT revealed incidental findings. 8 patients received further imaging, and 2 investigated with invasive procedures. 4 of the 5 new cancer diagnoses had abnormal findings after basic screening.

Standardised incidence ratios were calculated to assess the probability of presence of undiagnosed cancer in patients presenting with PE. Our data showed no significant increase in the incidence of cancer in patients presenting with PE compared to national cancer statistics (SIR of 0.75 in males (CI: 0.2–0.64) and 1.0 in females (CI: 0.23–0.56) aged 70–79).

Discussion Our data suggests that in patients presenting with acute PE, clinical acumen (as outlined by NICE) can be used to identify patients with potential malignancy. Our data would support limiting CT A/P to patients with significant clinical features or deranged tests.

REFERENCES

- 1 National Institute for Health and Clinical Excellence. Venous thromboembolic diseases: diagnosis, management and thrombophilia testing. CG144. NICE 2015.
- 2 Carrier M, Lazo-Langner A, Shivakumar S, et al. Screening for occult cancer in unprovoked venous thromboembolism. *New England Journal of Medicine* 2015;**373**(8):697–704.

P26 CT ABDOMEN AND PELVIS FOR UNPROVOKED PULMONARY EMBOLISM – WHAT IS THE BEST PRACTICE?

R Wahida, S Ahmad, B Saunders, N How, M Anwar. *Princess Alexandra Hospital NHS Foundation Trust, Harlow, United Kingdom*

10.1136/thoraxjnl-2016-209333.169

Background NICE guidelines advocate further investigations for cancer with an abdomino-pelvic CT scan (CT A/P) in all patients aged over 40 years with a first unprovoked deep vein thrombosis (DVT) or pulmonary embolism (PE) who do not have signs or symptoms of cancer based on initial investigation.¹ Recent prospective study showed no difference between 'limited' screening and CT A/P in diagnosing occult cancer.² We aimed to establish whether the number of new malignancies detected justified the risk of radiation exposure from performing a CT A/P in patient with unprovoked PE.

Methods We performed a retrospective analysis of all CT Pulmonary Angiograms (CTPA) performed during a one year period (2014–2015) in a district general hospital. Records of those patients with confirmed pulmonary embolism on CTPA (n = 254) were examined to ascertain whether performing CT A/P increased the detection rate of occult malignancy.

Results 124 (49%) out of a total 254 patients had an acute unprovoked PE. Of these, 6 patients were under the age of 40 years. Out of the remaining 118 patients, 80 (68%) patients underwent CT A/P. Unexpected malignancy was found in 3 (4%) of these 80 patients. No evidence of malignancy has been found

in those patients that did not undergo CT A/P (n = 38) so far based on the follow up clinical encounters – both as an in-patient and outpatient.

Conclusion Our data support the finding that the routine use of CT A/P in patients with unprovoked PE doesn't detect significant number of occult malignancies.

REFERENCES

- 1 Singh B, et al. *National institute for health and care excellence*. Venous thromboembolic diseases: diagnosis, management and thrombophilia testing. Nov 2012.
- 2 Carrier M, et al. Screening for occult cancer in unprovoked venous thromboembolism. *NEJM* 2015;**373**:8.

P27 EVALUATION AND BASELINE CHARACTERISTICS OF PATIENTS WITH CHRONIC THROMBOEMBOLIC DISEASE IN A SINGLE REFERRAL CENTRE

EM Swietlik, D Taboada, A Ruggiero, E Bales, L Harlow, A Fletcher, JE Cannon, K Sheares, DP Jenkins, J Pepke-Zaba, M Toshner. *Papworth Hospital NHS Foundation Trust, Cambridge, UK*

10.1136/thoraxjnl-2016-209333.170

Introduction Chronic thromboembolic disease (CTED) is a consequence of failure of thrombus resolution following pulmonary embolism. Thrombotic material becomes fibrosed, resulting in chronic vascular occlusion without pulmonary hypertension. The prevalence and incidence of the condition is unknown and the mechanisms behind exercise intolerance are poorly understood. Surgical management in selected cases may significantly improve symptoms and patient functioning.¹

Methods We prospectively evaluated baseline characteristics of patients with CTED in a single referral centre between January 2015 and June 2016. Newly referred patients with suspected CTED underwent a standard assessment as delineated in international guidelines with a minimum of 2 imaging modalities, resting and exercise right heart catheterisation and additionally incremental cardiopulmonary exercise testing (CPET). All patients were assessed in a pulmonary endarterectomy (PEA) MDT.

Results 128 patients were diagnosed with CTEPH or CTED from our referral centre. 28 patients were referred with suspected CTED due to ECHO findings. Of these 21 patients were confirmed to have CTED at right heart catheterization and 16 underwent full investigation protocol and were analysed. Patients with CTED were younger than contemporary cohorts of CTEPH² and were more likely to have a past medical history of VTE (94%). Patients with CTED had normal resting haemodynamics, preserved RV function at rest and normal NT-proBNP (Table1). After careful review of each patient's investigations only 5 of the 21 patients with CTED were offered PEA.

Conclusions Patients with CTED represent a significant proportion of the new referrals to our specialist centre. Surgery is deemed an appropriate therapeutic approach in a small subset of patients with significant functional and symptomatic impairment. The natural history of CTED is unclear so any discussion of surgery needs to carefully consider surgical risk of death and morbidity against the potential for symptomatic improvement.