

Supplementary Table II. Baseline comparison of smokers who did/did not complete T₂ questionnaires

| | | T₂ completers (n=865) n (%) or mean (SD) | T₂ non-completers (n=661)[‡] n (%) or mean (SD) | P value |
|--|-----------------------------|--|--|----------------|
| Trial allocation | Intervention | 488 (56%) | 263 (40%) | <0.001 |
| | Control | 377 (44%) | 398 (60%) | |
| Site | Liverpool | 445 (51%) | 397 (60%) | <0.001 |
| | Cambridge | 420 (49%) | 264 (40%) | |
| Age | Up to 65 years | 341 (39%) | 270 (41%) | 0.69 |
| | 66 – 70 years | 379 (44%) | 275 (42%) | |
| | Over 70 years | 145 (17%) | 116 (18%) | |
| Gender | Male | 608 (70%) | 463 (70%) | 0.92 |
| | Female | 257 (30%) | 198 (30%) | |
| Marital group | Married/cohabiting | 593 (69%) | 420 (64%) | 0.05 |
| | Not married/cohabiting | 271 (31%) | 239 (36%) | |
| IMD | Quintile 1 (most deprived) | 269 (31%) | 252 (38%) | <0.01 |
| | Quintile 2 | 108 (12%) | 80 (12%) | |
| | Quintile 3 | 133 (15%) | 119 (18%) | |
| | Quintile 4 | 149 (17%) | 92 (14%) | |
| | Quintile 5 (least deprived) | 206 (24%) | 118 (18%) | |
| Lung cancer experience | No | 520 (60%) | 342 (52%) | <0.001 |
| | Yes | 342 (40%) | 316 (48%) | |
| Cancer distress (T₀)[†] | | 2.26 (0.28) <i>9.58</i> | 2.24 (0.33) <i>9.39</i> | 0.40 |

Note: percentages were calculated based on available data.

[‡] Non-completers included non-responders and ineligible responders.

[†] Log_n scores in normal text, original scores in italics (analyses performed using log_n scores).